M1.(a) add excess copper carbonate (to dilute hydrochloric acid)

accept alternatives to excess, such as 'until no more reacts'

1

filter (to remove excess copper carbonate)

reject heat until dry

1

heat filtrate to evaporate some water **or** heat to point of crystallisation

accept leave to evaporate or leave in evaporating basin

1

leave to cool (so crystals form)

until crystals form

1

*must be in correct order to gain **4** marks*

(b) M_r $CuCl_2$ = 134.5

*correct answer scores **4** marks*

1

moles copper chloride = (mass / M_r = 11 / 134.5) = 0.0817843866

1

M_r $CuCO_3$= 123.5

1

Mass $CuCO_3$ (=moles × M_2= 0.08178 × 123.5) = 10.1(00)

1

*accept 10.1 with no working shown for **4** marks*

(c) $\frac{79.1}{100} \times 11.0$

or

11.0 × 0.791

1

8.70 (g)

1

*accept 8.70(g) with no working shown for **2** marks*

(d) Total mass of reactants = 152.5

1

$\frac{134.5}{152.5}$

allow ecf from step 1

1

88.20 (%)

1

*allow 88.20 with no working shown for **3** marks*

(e) atom economy using carbonate lower because an additional product is made **or** carbon dioxide is made as well

allow ecf

1

[14]

M2.(a) (delivery) tube sticks into the acid

1

the acid would go into the water **or** the acid would leave the flask or go up the delivery tube

ignore no gas collected

1

(b) any **one** from:

- bung not put in firmly / properly
- gas lost before bung put in
- leak from tube

1

(c) all of the acid has reacted

1

(d) take more readings in range 0.34 g to 0.54 g

1

take more readings is insufficient

ignore repeat

(e) $\frac{95}{24000}$

1

0.00396

or

3.96×10^{-3}

1

*accept 0.00396 or 3.96×10^{-3} with no working shown for **2** marks*

(f) use a pipette / burette to measure the acid

1

because it is more accurate volume than a measuring cylinder
or
greater precision than a measuring cylinder
or
use a gas syringe to collect the gas

so it will not dissolve in water

or
use a flask with a divider

accept description of tube suspended inside flask

so no gas escapes when bung removed

1

(g) they should be collected because carbon dioxide is left in flask at end

1

and it has the same volume as the air collected / displaced

1

[11]

M3.(a) **X:**

Fe^{2+} / iron(II), SO_4^{2-} / sulfate

allow iron(II) sulfate
or *$FeSO_4$*

1

Y:

Na^+ / sodium, I^- / iodide

allow sodium iodide
or *NaI*

1

Z:

Fe^{3+} / iron(III), Br^- / bromide

allow iron(III) bromide
or *$FeBr_3$*
correct identification of any two ions = one mark
correct identification of any four ions = two marks

1

(b) any **five** from:

allow converse arguments

method 1

- weighing is accurate
- not all barium sulfate may be precipitated
- precipitate may be lost
- precipitate may not be dry
- takes longer
- requires energy

allow not all the barium hydroxide has reacted

method 2

- accurate
- works for low concentrations

allow reliable / precise

5

[8]

M4.(a) copper has delocalised electrons

*accept copper has free electronsignore sea of electrons **or** mobile electrons*

1

(electrons) which can move through the metal / structure

allow (electrons) which can carry a charge through the metal / structure

1

(b) (i) (M_r $FeCl_3$ =) 162.5

*correct answer with or without working gains **3** marks*

*can be credited from correct substitution in step **2***

1

or

2 (moles of) $FeCl_3$ = 325

or

112 → 325

$$\frac{11.20}{56} \times 162.5$$

allow ecf from step 1

accept $\frac{325}{112} \times 11.2$

1

= 32.5

accept 32.48

1

(ii) 74.8

accept 74.77 - 75

accept ecf from (b)(i)

if there is no answer to part(i)

or

if candidate chooses not to use their answer then accept 86.79 - 87

1

[6]