M1.(a) both water <u>vapour</u> and ethanol will condense

allow steam for water vapour

allow they both become liquids

allow ethane condenses at a lower temperature

allow some of the steam hasn't reacted

allow it is a reversible reaction / equilibrium

1

(b) amount will decrease

1

because the equilibrium will move to the left

1

(c) more ethanol will be produced

1

because system moves to least / fewer molecules

1

[5]

M2.(a) because sulfur dioxide causes <u>acid rain</u>

1

which kills fish / aquatic life **or** dissolves / damages statues / stonework **or** kills / stunts growth of trees

if no other mark awarded then award 1 mark for sulfur dioxide is toxic or causes breathing difficulties.

1

(b) (i) <u>electrons</u> are lost

1

(ii) $Cu^{2+} + 2e^- \rightarrow Cu$

allow $Cu^{2+} \rightarrow Cu - 2e^-$

ignore state symbols

1

(iii) copper sulfate

allow any ionic copper compound

1

(c) (lattice of) positive ions

1

delocalised electrons

accept sea of electrons

1

(electrostatic) attraction between the positive ions and the electrons

1

electrons can move through the metal / structure **or** can flow

allow electrons can carry charge through the metal / structure

if wrong bonding named or described or attraction between oppositely charged ions then do not award M1 or M3 − MAX 2

1

(d) (copper compounds are absorbed / taken up by) plants

allow crops

1

which are burned

1

the ash contains the copper compounds

do not award M3 if the ash contains copper (metal)

1

(e)

/ A_r	55.6 / 63.5	16.4 / 56	28.0 / 32
moles	0.876	0.293	0.875
ratio	3	1	3
formula	Cu_3FeS_3		

*award **4** marks for Cu_3FeS_3 with some correct working*

*award **3** marks for Cu_3FeS_3 with **no** working*

*if the answer is not Cu_3FeS_3 award up to **3** marks for correct steps from the table apply ecf*

*if the student has inverted the fractions award **3** marks for an answer of $CuFe_3S$*

4

[16]

M3.(a) (i) the products are at a lower energy level than the reactants

accept products have less energy / less energy at the end than the beginning

1

(ii) because a catalyst provides an alternative / different pathway / mechanism / reaction route

accept adsorption or 'increases concentration at the surface'

ignore absorption

1

(that has) <u>lower activation energy</u>

allow weakens bonds

allow idea of increased successful collisions.

DO NOT ALLOW answers stating catalysts provide energy for M1 and M2


1

(b) one pair of electrons in each overlap (8 pairs in total)

allow any combination of dots, crosses or other symbols

1

the rest of the diagram correct with four non-bonding electrons on the oxygen giving a total of eight electrons in oxygen outer energy level.



*gains **2** marks*

1

(c) (i) ±3024 (J)

*correct answer with or without working gains **3** marks*

*if the answer is incorrect, award up to **2** marks for the following steps:*

- *ΔT = 14.4(°C)*
- *50 x 4.2 x 14.4*

allow ecf for incorrect ΔT

3

(ii) 0.015(2173913)

*correct answer with or without working gains **3** marks*

if answer is incorrect, allow 1 mark each for any of the following steps up to a max of 2.

- *0.70g*
- *M_r of ethanol = 46*
- *0.70 / 46*

allow ecf in final answer for arithmetical errors

3

(iii) ±198 720(J / mole)

c(i) ÷ c(ii)

*allow ecf from **(c)(i)** and **(c)(ii)***

0.015 gives 201600

0.0152 gives 198947

0.01522 gives 198686

1

(d) (as the molecules get bigger **or** the number of carbon atoms increases) the intermolecular forces

allow intermolecular bonds

1

(intermolecular forces) increase

allow more / stronger (intermolecular forces)

1

and therefore require more (heat) energy to overcome

*breaking covalent bonds or unspecified bonds max **1** mark (M3)*

1

[15]

M4.(a) (i) silver nitrate

allow $AgNO_3$

1

(ii) potassium carbonate **or**

allow K_2CO_3

sodium carbonate

allow Na_2CO_3

1

(b) base

allow ionic

ignore insoluble or soluble

ignore alkali

1

(c) (i) evaporate

or

crystallise

allow heat or boil or leave (to evaporate)

allow cool

ignore filtration unless given as an alternative

*do **not** accept freeze or solidify*

1

(ii) 2 (HNO_3)

accept multiples

1

(iii) 9

accept nine

1

(d) 6.21 / 207 0.72 / 16

***1** mark for dividing mass by A_r*

1

= 0.03 = 0.045

***1** mark for correct proportions (allow multiples)*

1

2 3

1 *mark for correct whole number ratio (allow multiples). Can be awarded from formula.*

1

Pb_2O_3

allow O_3Pb_2

ecf *allowed throughout if sensible attempt at step 1*

correct formula with no working gains ***1*** *mark*

1

[10]

M5.(a) lattice / giant structure

*max **3** if incorrect structure or bonding or particles*

1

ionic **or** (contains) ions

1

Na^+ **and** Cl^-

accept in words or dot and cross diagram: must include type and magnitude of charge for each ion

1

electrostatic attraction

allow attraction between opposite charges

1

(b) hydrogen

allow H_2

1

sodium hydroxide

allow NaOH

1

(c) any **one** from, eg:

- people should have the right to choose
- insufficient evidence of effect on individuals
- individuals may need different amounts.

allow too much could be harmful

ignore religious reasons


ignore cost

ignore reference to allergies

1

(d) (i) one bonding pair of electrons

accept dot, cross or e or − or any combination, eg



1

6 unbonded electrons on each atom

1

(ii) simple molecules

*max **2** if incorrect structure or bonding or particles*

accept small molecules

accept simple / small molecular structure

1

with intermolecular forces

accept forces between molecules

must be no contradictory particles

1

which are weak **or** which require little energy to overcome − must be linked to second marking point

reference to weak covalent bonds negates second and third marking points

1

(iii) iodine has no delocalised / free / mobile electrons or ions

1

so cannot carry charge

*if no mark awarded iodine molecules have no charge gains **1** mark*

1

[14]