M1.(a) (i) calcium oxide

in either order

1

carbon dioxide

accept correct formulae

1

(ii) $C(s) + CO_2(g) \rightarrow \mathbf{2}CO(g)$

allow multiples

1

(iii) 210 (tonnes)

*award **3** marks for the correct answer with or without working*

allow ecf for arithmetical errors

*if answer incorrect allow up to **2** marks for any of the steps below:*

160 → 112

300 → 112 / 160 × 300

or

moles Fe_2O_3 = 1.875 (× 10^6) or 300 / 160

moles of Fe = 3.75 (× 10^6) or 2 × moles Fe_2O_3

mass Fe = moles Fe × 56

105 (tonnes) scores 2 (missing 1:2 ratio)

420 (tonnes) scores 2 − taken M_r of iron as 112

3

(b) (i) aluminium is more reactive than carbon **or** carbon is less reactive than aluminium

must have a comparison of reactivity of carbon and aluminium

accept comparison of position in reactivity series.

1

(ii) (because) aluminium ions are positive

ignore aluminium is positive

1

and are attracted / move / go to the negative electrode / cathode

1

where they gain electrons / are reduced / $Al^{3+} + 3e^- \rightarrow Al$

accept equation or statements involving the wrong number of electrons.

1

(iii) (because) the anodes **or** (positive) electrodes are made of carbon / graphite

1

oxygen is produced (at anode)

1

which reacts with the electrodes / anodes

*do **not** accept any reference to the anodes reacting with oxygen from the air*

equation $C + O_2 \rightarrow CO_2$ *gains **1** mark (M3)*

1

[13]

M2.(a) left hand: (conical) flask

*do **not** accept round bottomed flask or container which is not a flask*

1

right hand: beaker / trough

accept plastic box

1

(b) (i) 157

1

(ii) all calcium carbonate used up **or** reaction stopped

*do **not** accept all acid used up*

1

(c) (i) 0.007(272727…)

*correct answer with or without working gains **2** marks*

*if answer incorrect, allow (0.32 / 44) for **1** mark*

2

(ii) 0.007(272727…)

*allow ecf from **(c)(i)***

1

(iii) (M_r = mass / moles = 1 / 0.00727…) = 137.5 or 138

*allow ecf from **(c)(ii)***

if use 0.00943 moles then = 106

if use 0.007 allow 143 (142.857)

1

(iv) (138) – 60 (= 78)

23 / 85

1

(78 / 2) = 39

1

potassium

sodium / rubidium

*identity of metal ecf on A_r, but **must** be Group 1*

*If no working max **1** mark*

1

(d) (i) (relative atomic mass) would decrease

1

because the mass lost greater

1

so moles carbon dioxide larger **or** moles metal carbonate greater

1

(ii) no change

1

because the acid (already) in excess

1

so the amount carbon dioxide lost is the same

1

[17]

M3.(a) copper has delocalised electrons

*accept copper has free electronsignore sea of electrons **or** mobile electrons*

1

(electrons) which can move through the metal / structure

allow (electrons) which can carry a charge through the metal / structure

1

(b) (i) (M_r $FeCl_3$ =) 162.5

*correct answer with or without working gains **3** marks*

*can be credited from correct substitution in step **2***

1

or

2 (moles of) $FeCl_3$ = 325

or

112 → 325

$$\frac{11.20}{56} \times 162.5$$

allow ecf from step 1

accept $\frac{325}{112} \times 11.2$

1

= 32.5

accept 32.48

1

(ii) 74.8

accept 74.77 - 75

accept ecf from (b)(i)

if there is no answer to part(i)

or

if candidate chooses not to use their answer then accept 86.79 - 87

1

[6]

M4.(a) (i) CH_4

allow H_4C

*do **not** allow lower-case h*

*do **not** allow superscript*

1

(ii) single

1

(iii) alkanes

1

(b) (i) carbon / C

any order

1

hydrogen / H

allow phonetic spelling

1

sulfur / sulphur / S

1

(ii) air / atmosphere

1

(iii) acid rain

1

damages trees / plants **or** kills aquatic organisms **or** damages buildings / statues **or** causes respiratory problems

allow harmful to living things

1

(c) carbon / C

accept soot / particulates / charcoal

1

(d) any **four** from:

- (supports hypothesis) because when the fuel contained more carbon the temperature of the water went up more / faster (in 2 minutes)
- (does not support hypothesis as) temperature change per gram decreases as the number of carbons increases
- (does not support hypothesis) because the more carbon in the fuel the more smoke **or** the dirtier / sootier it is
- only tested hydrocarbons / alkanes / fuels with between 5 and 12 carbon atoms
- valid, justified, conclusion

accept converse statements

4

(e) (i) 0.15

*correct answer with or without working gains **2** marks*

*if answer incorrect, M_r carbon dioxide = 44 gains **1** mark*

*allow 0.236 / 0.24 / 0.2357142 (ecf from M_r of 28) for **1** mark*

2

(ii) 0.4(0)

1

(iii) C_3H_8

*correct formula with or without working scores **2** marks*

0.15 / 0.05 = 3

*allow ecf from **(e)(i)***

and

0.4 / 0.05 = 8 (1)

*allow ecf from **(e)(ii)***

*allow **1** mark for correct empirical formula from their values*

If use 'fall-back-values:

0.50 / 0.05 = 10

and

0.20 / 0.05 = 4

1 mark

C_4H_{10}

1 mark

if just find ratio of C to H using fall-back values, get C_2H_5
*allow **1** mark*

2

[19]