Q1. This question is about the reaction of ethene and bromine.

The equation for the reaction is:


$$C_2H_4 + Br_2 \rightarrow C_2H_4Br_2$$

(a) Complete the reaction profile in **Figure 1**.

Draw labelled arrows to show:

- The energy given out (ΔH)
- The activation energy.

Figure 1



(3)

(b) When ethene reacts with bromine, energy is required to break covalent bonds in the molecules.

Explain how a covalent bond holds two atoms together.

.....

.....


.....

.....

(2)

(c) **Figure 2** shows the displayed formulae for the reaction of ethene with bromine.

Figure 2



The bond enthalpies and the overall energy change are shown in the table below.

	C=C	**C–H**	**C–C**	**C–Br**	**Overall energy change**
Energy in kJ / mole	612	412	348	276	–95

Use the information in the table above and **Figure 2** to calculate the bond energy for the Br–Br bond.

.....

.....


.....

Bond energy ..... kJ / mole

(3)

(d) **Figure 3** shows the reaction between ethene and chlorine and is similar to the reaction between ethene and bromine.

Figure 3



"The more energy levels (shells) of electrons an atom has, the weaker the covalent bonds that it forms."

Use the above statement to predict and explain how the overall energy change for the reaction of ethene with chlorine will differ from the overall energy change for the reaction

of ethene with bromine.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(6)
(Total 14 marks)


Q2. This question is about ethanol.

(a) Ethanol is produced by the reaction of ethene and steam:

$$C_2H_4 + H_2O \longrightarrow C_2H_5OH$$

(i) **Figure 1** shows the energy level diagram for the reaction.

Figure 1



How does the energy level diagram show that the reaction is exothermic?

.....

.....

(1)

(ii) A catalyst is used for the reaction.

Explain how a catalyst increases the rate of the reaction.

.....

.....


.....

.....

(2)

(b) **Figure 2** shows the displayed structure of ethanol.


Figure 2



Complete the dot and cross diagram in **Figure 3** to show the bonding in ethanol.

Show the outer shell electrons only.

Figure 3




(2)

(c) A student burned some ethanol.

Figure 4 shows the apparatus the student used.

Figure 4



(i) The student recorded the temperature of the water before and after heating.

His results are shown in **Table 1**.

Table 1

Temperature before heating	20.7 °C
Temperature after heating	35.1 °C

Calculate the energy used to heat the water.

Use the equation $Q = m \times c \times \Delta T$

The specific heat capacity of water = 4.2 J / g / °C

.....

.....

.....

.....

.....

Energy used = ... J

(3)

(ii) **Table 2** shows the mass of the spirit burner before the ethanol was burned and after the ethanol was burned.

Table 2

Mass of spirit burner before ethanol was burned	72.80 g
Mass of spirit burner after ethanol was burned	72.10 g

Calculate the number of moles of ethanol (C_2H_5OH) that were burned.

Relative atomic masses (A_r): H = 1; C = 12; O = 16

.....

.....

.....

.....

.....

Number of moles burned = ...

(3)

(iii) Calculate the energy released in joules per mole.

You should assume that all the energy from the ethanol burning was used to heat the water.




.....

Energy = ... J / mole

(1)

(d) The names, structures and boiling points of ethanol and two other alcohols are shown in **Table 3**.

Table 3

Name	**Methanol**	**Ethanol**	**Propanol**
Structure			
Boiling point in °C	65	78	97

Use your knowledge of structure and bonding to suggest why the boiling points increase as the number of carbon atoms increases.

.....

.....

.....

.....

.....

.....

.....

(3)
(Total 15 marks)

Q3. Dilute nitric acid reacts with potassium hydroxide solution.

The equation for the reaction is:

$$HNO_3 + KOH \longrightarrow KNO_3 + H_2O$$

A student investigated the temperature change in this reaction.


This is the method the student used.

Step 1 Put 25 cm^3 of dilute nitric acid in a polystyrene cup.
Step 2 Use a thermometer to measure the temperature of the dilute nitric acid.
Step 3 Use a burette to add 4 cm^3 of potassium hydroxide solution to the dilute nitric acid and stir the mixture.
Step 4 Use a thermometer to measure the highest temperature of the mixture.
Step 5 Repeat steps 3 and 4 until 40 cm^3 of potassium hydroxide solution have been added.

The dilute nitric acid and the potassium hydroxide solution were both at room temperature.

(a) **Figure 1** shows part of the thermometer after some potassium hydroxide solution had been added to the dilute nitric acid.

Figure 1



What is the temperature shown on the thermometer?

The temperature shown is ..... °C

(1)

(b) Errors are possible in this experiment.

(i) Suggest **two** causes of random error in the experiment.

.....

.....

.....

.....

(2)

(ii) Another student used a glass beaker instead of a polystyrene cup.

This caused a systematic error.

Why does using a glass beaker instead of a polystyrene cup cause a systematic error?

.....


.....

.....

(1)

(c) The results of the student using the polystyrene cup are shown in **Figure 2**.

Figure 2



(i) How do the results in **Figure 2** show that the reaction between dilute nitric acid and potassium hydroxide solution is exothermic?

.....

.....

(1)

(ii) Explain why the temperature readings decrease between 28 cm^3 and 40 cm^3 of potassium hydroxide solution added.

.....

.....

.....

(2)

(iii) It is difficult to use the data in **Figure 2** to find the exact volume of potassium hydroxide solution that would give the maximum temperature.

Suggest further experimental work that the student should do to make it easier to find the exact volume of potassium hydroxide solution that would give the maximum temperature

.....

.....

.....

.....

(2)

(d) The student did further experimental work and found that 31.0 cm^3 of potassium hydroxide solution neutralised 25.0 cm^3 of dilute nitric acid.

The concentration of the dilute nitric acid was 2.0 moles per dm^3.

$$HNO_3 + KOH \longrightarrow KNO_3 + H_2O$$

Calculate the concentration of the potassium hydroxide solution in moles per dm^3.

.....

.....

.....

.....

.....

.....

Concentration = ... moles per dm^3

(3)

(e) The student repeated the original experiment using 25 cm^3 of dilute nitric acid in a polystyrene cup and potassium hydroxide solution that was twice the original concentration.

She found that:

- a smaller volume of potassium hydroxide solution was required to reach the maximum temperature
- the maximum temperature recorded was higher.

Explain why the maximum temperature recorded was higher.

.....

.....


.....

.....

(2)
(Total 14 marks)

Q4. Methane (CH_4) is used as a fuel.

(a) The displayed structure of methane is:



Draw a ring around a part of the displayed structure that represents a covalent bond.

(1)

(b) Why is methane a compound?

Tick (✓) **one** box.

Methane contains atoms of two elements, combined chemically. ☐

Methane is not in the periodic table. ☐

Methane is a mixture of two different elements. ☐


(1)

(c) Methane burns in oxygen.

(i) The diagram below shows the energy level diagram for the complete combustion of methane.

Draw and label arrows on the diagram to show:

- the activation energy
- the enthalpy change, ΔH.



(2)

(ii) Complete and balance the symbol equation for the complete combustion of methane.

CH_4 + ... ⟶ CO_2 + ...

(2)

(iii) Explain why the **incomplete** combustion of methane is dangerous.

.....

.....

.....

.....

(2)

(iv) Explain why, in terms of the energy involved in bond breaking and bond making, the combustion of methane is exothermic.

.....

.....

.....

.....

.....

.....

(3)

(d) Methane reacts with chlorine in the presence of sunlight.

The equation for this reaction is:



Some bond dissociation energies are given in the table.

Bond	**Bond dissociation energy in kJ per mole**
C–H	413
C–Cl	327
Cl–Cl	243
H–Cl	432

(i) Show that the enthalpy change, ΔH, for this reaction is −103 kJ per mole.

.....

.....

.....


.....

.....

.....

(3)

(ii) Methane also reacts with bromine in the presence of sunlight.



This reaction is less exothermic than the reaction between methane and chlorine.

The enthalpy change, ΔH, is −45 kJ per mole.

What is a possible reason for this?

Tick (✓) **one** box.

CH_3Br has a lower boiling point than CH_3Cl ☐

The C−Br bond is weaker than the C−Cl bond. ☐

The H−Cl bond is weaker than the H−Br bond. ☐

Chlorine is more reactive than bromine. ☐

(1)

(Total 15 marks)