Oxford Cambridge and RSA

Friday 6 June 2025 – Afternoon

AS Level Further Mathematics B (MEI)

Y414/01 Numerical Methods

Time allowed: 1 hour 15 minutes

You must have:
- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

QP

1869651726

INSTRUCTIONS
- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined page at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION
- The total mark for this paper is **60**.
- The marks for each question are shown in brackets **[ ]**.
- This document has **8** pages.

ADVICE
- Read each question carefully before you start your answer.

© OCR 2025 [R/508/5557]
DC (PQ/SG) 351558/4
OCR is an exempt Charity

1 The method of interval bisection is used to find a sequence of approximations to one of the roots of the equation $e^x - x^2 - 3x = 0$.

The table shows the associated spreadsheet output.

	A	B	C	D	E	F
1	a	f(a)	b	f(b)	c	f(c)
2	2	−2.61094	3	2.085537	2.5	−1.56751
3	2.5	−1.56751	3	2.085537	2.75	−0.16987
4	2.75	−0.16987	3	2.085537	2.875	0.834799
5	2.75	−0.16987	2.875	0.834799	2.8125	0.303839
6						

(a) Write down a suitable formula for cell E2. **[1]**

The formula in cell A3 is $\boxed{= \text{IF}(\text{F2} < 0, \text{E2}, \text{A2})}$.

(b) Write down a similar formula for cell C3. **[1]**

(c) Complete row 6 of the table in the **Printed Answer Booklet**. **[2]**

(d) **Without** doing any more calculations, write down the value of the root correct to the maximum number of decimal places which seems justified. You must explain the precision quoted. **[1]**

2 The table gives three values of x and the associated values of y.

x	−1	2	3
y	−3.39	0.18	0.45

Use Lagrange's form of the interpolating polynomial to construct a polynomial of degree 2 for the values in the table. Give your answer in the form

$$y = ax^2 + bx + c,$$

where a, b and c are constants to be determined. **[4]**

3 **(a)** Find the relative error when π is **chopped** to **3** decimal places. **[2]**

(b) Find the relative error when π is **rounded** to **3** decimal places. **[2]**

You are given that $y = \pi^2 - 5$ and $z = (\pi - 5)^2$.

You are also given the following information.

- Y is an approximation to y.
- Z is an approximation to z.
- Y and Z are found by using $\pi = 3.14$.

A student states that the relative error in using Y to approximate y is exactly the same as the relative error in using Z to approximate z, because in each case the calculation involves squaring and subtracting 5.

(c) **Without** doing any calculations, explain whether the student is correct. **[1]**

4 A student is using a spreadsheet to investigate the curve $y = \mathrm{f}(x)$ at the point where $x = 2$. Some of the spreadsheet output is shown below.

Table 4.1 shows some values of x and the associated values of $\mathrm{f}(x)$.

Table 4.2 shows **two** approximations to $\mathrm{f}'(2)$ found using the same method with $h = 0.001$ and $h = 0.01$.

	E	F	G	H	I	J
2						
3		**Table 4.1**				
4		x	2	2.001	2.01	
5		$\mathrm{f}(x)$	0.30103	0.30125	0.3032	
6						
7						
8		**Table 4.2**				
9		h	0.001	0.01		
10		$\mathrm{f}'(2)$	0.21709	0.21661		
11						

(a) The formula in cell G10 is $\boxed{= (\mathrm{H5} - \mathrm{G5})/\mathrm{G9}}$.

The formula in cell H10 is $\boxed{= (\mathrm{I5} - \mathrm{G5})/\mathrm{H9}}$.

State the method being used to find these approximations to $\mathrm{f}'(2)$. **[1]**

(b) • Calculate $\dfrac{0.30125 - 0.30103}{0.001}$.

• Explain why your answer is not the same as the value displayed in cell G10. **[3]**

(c) State the value of $\mathrm{f}'(2)$ as accurately as you can. You must justify the precision quoted. **[1]**

(d) Hence determine an estimate for the error when $\mathrm{f}(2)$ is used as an approximation to $\mathrm{f}(2.1)$. **[2]**

(e) Explain whether your answer to part **(d)** is likely to be an over-estimate or an under-estimate. **[1]**

5 The spreadsheet output shows a sequence of approximations to $\int_{0.5}^{1.5} \mathrm{f}(x)\,\mathrm{d}x$ using the trapezium rule and Simpson's rule, together with some further analysis of the approximations found using Simpson's rule.

	F	G	H	I	J
21	n	T_n	S_n	difference	ratio
22	1	0.84557616			
23	2	0.88144672	0.89340357		
24	4	0.89066651	0.89373977	0.00034	
25	8	0.89299453	0.89377054	3.1E-05	0.092
26	16	0.89357822	0.89377279	2.2E-06	0.073
27	32	0.89372426	0.89377294	1.5E-07	0.066
28	64	0.89376077	0.89377295	9.4E-09	0.063
29	128	0.8937699	0.89377295	5.9E-10	0.063

(a) Write down the value displayed in cell I25 using standard mathematical notation. **[1]**

(b) State the value of $\int_{0.5}^{1.5} \mathrm{f}(x)\,\mathrm{d}x$ as accurately as you can. You must justify the precision quoted. **[1]**

(c) Write down a suitable formula for cell H23. **[2]**

(d) **(i)** Explain what the values in column J tell you about the order of Simpson's rule in this case. **[2]**

(ii) Explain whether this is usual. **[1]**

6 The equation $0.5\ln x + x^2 - x - 2 = 0$ has a root α, such that $1 < \alpha < 2$.

A student attempts to find α using the iterative formula

$x_{n+1} = \mathrm{g}(x_n)$ with $x_0 = 1$, where $\mathrm{g}(x) = 0.5\ln x + x^2 - 2$.

The associated spreadsheet output is shown in the table.

	K	L
1	n	x_n
2	0	1
3	1	-1
4	2	#NUM!

(a) Explain why #NUM! is displayed in cell L4. **[1]**

The graph shows part of the graphs of $y = x$ and $y = \mathrm{g}(x)$.



(b) On the copy of the graph in the **Printed Answer Booklet** show how the iterative formula $x_{n+1} = \mathrm{g}(x_n)$ with $x_0 = 2$ finds x_1 and x_2. **[1]**

(c) Explain why the iterative formula $x_{n+1} = \mathrm{g}(x_n)$ may **not** be used to successfully find α, whatever the value of x_0. **[1]**

(d) Use the relaxed iteration

$$x_{n+1} = (1-\lambda)x_n + \lambda \mathrm{g}(x_n),$$

with $\lambda = -0.328$ and $x_0 = 1$, to determine the value of α correct to **6** decimal places. **[3]**

7 A student makes a cup of tea and investigates how long it takes to reach room temperature. The student places the cup of tea on a table in the kitchen and records the temperature T°C of the tea t minutes after it was made, at **5** minute intervals starting with $t = 5$. The data collected are shown in the table.

Time after tea was made t (minutes)	5	10	15	20
Temperature of tea T (°C)	78.5	68.1	58.7	50.3

(a) Show that a quadratic polynomial would be a reasonable model for these data. **[3]**

(b) Use Newton's forward difference interpolation method to determine a suitable quadratic polynomial for these data. **[4]**

(c) The temperature in the kitchen is 19°C. It takes one hour for the tea to cool to room temperature.

(i) Use the model to find an estimate of the temperature of the tea after 1 hour. **[1]**

(ii) Use the model to find an estimate of the temperature of the tea after an hour and a half. **[1]**

(iii) Hence identify a limitation of the model. **[1]**

8 **Table 8.1** shows some values of x and the associated values of $f(x)$.

Table 8.1

x	0.5	0.9	1.3
$f(x)$	1.41421	1.09947	0.71101

(a) Use the central difference method to determine an approximation to $f'(0.9)$. **[2]**

Table 8.2 shows a sequence of approximations to $f'(0.9)$ for different values of the step length, h, together with some further analysis.

Table 8.2

h	0.2	0.1	0.05	0.025	0.0125
$f'(0.9)$	−0.957844	−0.977203	−0.982021	−0.983225	−0.983525
difference		−0.019360	−0.004818	−0.001203	−0.000301
ratio			0.248882	0.249719	0.249930

(b) Use extrapolation to determine the value of $f'(0.9)$ as accurately as you can. You must justify the precision quoted. **[5]**

Turn over for question 9

9 In this question you must show detailed reasoning

The equation $0.2x^4 + 0.8x - 0.2 = 0$ has a root α such that $0 < \alpha < 1$.

(a) Use the Newton-Raphson method with $x_0 = 1$ to determine the value of α correct to **6** decimal places. **[5]**

The equation $0.2x^4 + 0.8x - 0.2 = 0$ has another root β such that $-2 < \beta < -1$.

The Newton-Raphson method is used to find a sequence of approximations to β.
The associated spreadsheet output is shown in the table below, together with some further analysis.

A	B	C	D
r	x_r	difference	ratio
0	−2		
1	−1.75	0.25	
2	−1.670923	0.079077	0.3163082
3	−1.663319	0.007604	0.096159
4	−1.663252	6.7E-05	0.0088147
5	−1.663252	5.18E-09	7.724E-05

(b) (i) Explain what the values in column D suggest about the order of convergence of this sequence of approximations. **[2]**

(ii) Explain whether this is usual. **[1]**

END OF QUESTION PAPER

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

© OCR 2025