

Please check the examination details below before entering your candidate information

Candidate surname		Other names	
Centre Number		Candidate Number	

Pearson Edexcel Level 3 GCE

Friday 7 June 2024

Afternoon (Time: 1 hour 30 minutes) **Paper reference** **9FM0/3C**

Further Mathematics
Advanced
PAPER 3C: Further Mechanics 1

You must have:
 Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P75322A

©2024 Pearson Education Ltd.
F:1/1/



1. [In this question, $\mathbf{i}$ and $\mathbf{j}$ are horizontal perpendicular unit vectors.]

A particle A has mass 3 kg and a particle B has mass 2 kg.

The particles are moving on a smooth horizontal plane when they collide directly.

Immediately **before** the collision, the velocity of A is $(3\mathbf{i} - \mathbf{j})\text{ m s}^{-1}$ and the velocity of B is $(-6\mathbf{i} + 2\mathbf{j})\text{ m s}^{-1}$

Immediately **after** the collision the velocity of A is $\left(-2\mathbf{i} + \frac{2}{3}\mathbf{j}\right)\text{ m s}^{-1}$

- (a) Find the total kinetic energy of the two particles **before** the collision. (3)
- (b) Find, in terms of $\mathbf{i}$ and $\mathbf{j}$, the impulse exerted on A by B in the collision. (3)
- (c) Find, in terms of $\mathbf{i}$ and $\mathbf{j}$, the velocity of B immediately **after** the collision. (3)



DO NOT WRITE IN THIS AREA

Question 1 continued

Lined area for writing answers.

(Total for Question 1 is 9 marks)



2. A rough plane is inclined to the horizontal at an angle θ , where $\tan \theta = \frac{3}{4}$

A particle P of mass m is at rest at a point on the plane.

The particle is projected **up** the plane with speed $\sqrt{2ag}$

The particle moves up a line of greatest slope of the plane and comes to instantaneous rest after moving a distance d .

The coefficient of friction between P and the plane is $\frac{1}{7}$

- (a) Show that the magnitude of the frictional force acting on P as it moves up the plane

is $\frac{4mg}{35}$ (3)

Air resistance is assumed to be negligible.

Using the work-energy principle,

- (b) find d in terms of a . (4)



DO NOT WRITE IN THIS AREA

Question 2 continued

Handwriting practice area with horizontal lines.

(Total for Question 2 is 7 marks)



3. A car of mass 1000 kg moves in a straight line along a horizontal road at a constant speed of 72 km h^{-1}

- The resistance to the motion of the car is modelled as a constant force of magnitude 900 N

The engine of the car is working at a constant rate of $P\text{ kW}$.

Using the model,

- (a) find the value of P .

(3)

The car now travels in a straight line up a road which is inclined to the horizontal at an

angle α , where $\sin \alpha = \frac{2}{49}$

- In a refined model, the resistance to the motion of the car from non-gravitational forces is now modelled as a force of magnitude $20v$ newtons, where $v\text{ m s}^{-1}$ is the speed of the car

At the instant when the engine of the car is working at a constant rate of 30 kW and the car is moving up the road at 10 m s^{-1} , the acceleration of the car is $a\text{ m s}^{-2}$

Using the refined model,

- (b) find the value of a .

(4)

Later on, when the engine of the car is again working at a constant rate of 30 kW , the car is moving up the road at a constant speed $U\text{ m s}^{-1}$

Using the refined model,

- (c) find the value of U .

(5)



DO NOT WRITE IN THIS AREA

Question 3 continued

Lined area for writing the answer to Question 3.



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 3 continued

Lined area for writing answers.

(Total for Question 3 is 12 marks)



DO NOT WRITE IN THIS AREA

Question 4 continued

Lined area for writing the answer to Question 4.



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 4 continued

Handwriting practice area with horizontal lines.

(Total for Question 4 is 15 marks)



5. A light elastic string has natural length $2a$ and modulus of elasticity $2mg$. One end of the string is attached to a fixed point A on a horizontal ceiling. The other end is attached to a particle P of mass m .

The particle P hangs in equilibrium at the point E , where $AE = 3a$.

The particle P is then projected vertically downwards from E with speed $\frac{3}{2}\sqrt{ag}$

Air resistance is assumed to be negligible.

Find the elastic energy stored in the string, when P first comes to instantaneous rest. Give your answer in the form $kmg a$, where k is a constant to be found.

(7)



DO NOT WRITE IN THIS AREA

Question 5 continued

Handwriting practice area with horizontal lines.

(Total for Question 5 is 7 marks)



DO NOT WRITE IN THIS AREA

Question 6 continued

Lined area for writing the answer to Question 6.



Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 6 continued

Handwriting practice area with horizontal lines.

(Total for Question 6 is 10 marks)



7.




Figure 1

A smooth uniform sphere A of mass m is moving with speed U on a smooth horizontal plane. The sphere A collides obliquely with a smooth uniform sphere B of mass $3m$ which is at rest on the plane. The two spheres have the same radius.

Immediately before the collision, the direction of motion of A makes an angle α , where $0^\circ < \alpha < 90^\circ$, with the line joining the centres of the spheres.

Immediately after the collision, the direction of motion of A is **perpendicular** to its original direction, as shown in Figure 1.

The coefficient of restitution between the spheres is e .

(a) Show that the speed of B immediately after the collision is

$$\frac{1}{4}(1+e)U\cos\alpha \quad (6)$$

(b) Show that $e > \frac{1}{3}$ (4)

(c) Show that $0 < \tan\alpha \leq \frac{1}{\sqrt{2}}$ (5)



DO NOT WRITE IN THIS AREA

Question 7 continued

Lined area for writing the answer to Question 7.



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 7 continued

Handwriting practice area with horizontal lines.



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 is 15 marks)

TOTAL FOR PAPER IS 75 MARKS

