



CANDIDATE  
NAME

|  |
|--|
|  |
|--|

CENTRE  
NUMBER

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

CANDIDATE  
NUMBER

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

## 9231/21

October/November 2020

**2 hours**

You must answer on the question paper.

You will need: List of formulae (MF19)

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.

- 1 (a)** By differentiating  $e^{-x^2}$ , find the Maclaurin's series for  $e^{-x^2}$  up to and including the term in  $x^2$ . [5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- (b) Deduce an approximation to  $\int_0^{\frac{1}{5}} e^{-x^2} dx$ , giving your answer as a rational fraction in its lowest terms. [2]

[illegible]

**2** The variables  $x$  and  $y$  are related by the differential equation

$$9\frac{d^2y}{dx^2} + 6\frac{dy}{dx} + y = 3x^2 + 30x.$$

(a) Find the general solution for  $y$  in terms of  $x$ .

[6]

This image shows a full page of white paper with horizontal dotted lines, typical of primary-ruled notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

**(b)** State an approximate solution for large positive values of  $x$ .

[1]

.....

.....

.....

.....

- 3 (a) Show that the system of equations

$$\begin{aligned}x - 2y - 4z &= 1, \\x - 2y + kz &= 1, \\-x + 2y + 2z &= 1,\end{aligned}$$

where  $k$  is a constant, does not have a unique solution.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Given that  $k = -4$ , show that the system of equations in part (a) is consistent. Interpret this situation geometrically.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Given instead that  $k = -2$ , show that the system of equations in part (a) is inconsistent. Interpret this situation geometrically. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) For the case where  $k \neq -2$  and  $k \neq -4$ , show that the system of equations in part (a) is inconsistent. Interpret this situation geometrically. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....


$$\int_0^1 (1-x^3) dx \leq \frac{3n^2+2n-1}{4n^2}. \quad [4]$$
[illegible]

- (b) Use a similar method to find, in terms of  $n$ , a lower bound for  $\int_0^1 (1 - x^3) dx$ . [4]

**5** It is given that

$$x = \sinh^{-1} t, \quad y = \cos^{-1} t,$$

where  $-1 < t < 1$ .

(a) By differentiating  $\cos y$  with respect to  $t$ , show that  $\frac{dy}{dt} = -\frac{1}{\sqrt{1-t^2}}$ . [4]

[illegible]

- (b)** Find  $\frac{d^2y}{dx^2}$  in terms of  $t$ , simplifying your answer. [5]

[illegible]

- 6 (a)** Use de Moivre's theorem to show that  $\sin^4 \theta = \frac{1}{8}(\cos 4\theta - 4 \cos 2\theta + 3)$ . [5]

[illegible]

**(b)** Find the solution of the differential equation

$$\frac{dy}{d\theta} + y \cot \theta = \sin^3 \theta$$

for which  $y = 0$  when  $\theta = \frac{1}{2}\pi$ .

[6]

[illegible]

**7** The matrix  $\mathbf{P}$  is given by

$$\mathbf{P} = \begin{pmatrix} 1 & 4 & 2 \\ 0 & -1 & 1 \\ 0 & 0 & 2 \end{pmatrix}.$$

(a) State the eigenvalues of  $\mathbf{P}$ . [1]

.....

.....

.....

.....

.....

(b) Use the characteristic equation of  $\mathbf{P}$  to find  $\mathbf{P}^{-1}$ . [4]

This image shows a full page of a document template designed for handwriting practice. It features a series of evenly spaced, horizontal dashed lines across the entire width of the page. The background is plain white, providing a clear guide for letter height and placement. There are no margins, text, or other markings present.



- 8 (a)** Sketch the graph of  $y = \coth x$  for  $x > 0$  and state the equations of the asymptotes. [2]

- (b)** Starting from the definitions of  $\coth$  and  $\operatorname{cosech}$  in terms of exponentials, prove that

$$\coth^2 x - \operatorname{cosech}^2 x = 1. \quad [3]$$

[illegible]

The curve  $C$  has equation  $y = \ln \coth\left(\frac{1}{2}x\right)$  for  $x > 0$ .

- (c) Show that  $\frac{dy}{dx} = -\operatorname{cosech} x$ . [3]

---

---

---

---

---

---

- (d)** It is given that the arc length of  $C$  from  $x = a$  to  $x = 2a$  is  $\ln 4$ , where  $a$  is a positive constant.

Show that  $\cosh a = 2$  and find, in logarithmic form, the exact value of  $a$ . [7]

[illegible]

[illegible]

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.