Pearson Edexcel

Mark Scheme (Results)

January 2021

Pearson Edexcel International Advanced Level
In Further Pure Mathematics F1
Paper WFM01/01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2021
Publications Code WFM01_01_2101_MS
All the material in this publication is copyright
© Pearson Education Ltd 2021

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75

2. The Edexcel Mathematics mark schemes use the following types of marks:

- **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.

3. Abbreviations

   These are some of the traditional marking abbreviations that will appear in the mark schemes.

   - bod – benefit of doubt
   - ft – follow through
   - the symbol $\sqrt{}$ will be used for correct ft
   - cao – correct answer only
   - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
   - isw – ignore subsequent working
   - awrt – answers which round to
   - SC: special case
   - o.e. – or equivalent (and appropriate)
   - d… or dep – dependent
   - indep – independent
   - dp decimal places
   - sf significant figures
   - * The answer is printed on the paper or ag- answer given
   - ⌊ or d… The second mark is dependent on gaining the first mark

4. All A marks are ‘correct answer only’ (cao), unless shown, for example, as A1ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

6. If a candidate makes more than one attempt at any question:
    - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
    - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.

7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Further Pure Mathematics Marking

(*But note that specific mark schemes may sometimes override these general principles*)

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to $x = \ldots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to $x = \ldots$

2. Formula

Attempt to use the correct formula (with values for *a*, *b* and *c*).

3. Completing the square

Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0,\ \ q \neq 0$, leading to $x = \ldots$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. $(x^n \to x^{n-1})$

2. Integration

Power of at least one term increased by 1. $(x^n \to x^{n+1})$

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Question Number	Scheme	Marks
1.(a)	$f(0.2)=...$ and $f(0.6)=...$	M1
	$f(0.2)=-0.5973...$ and $f(0.6)=0.2707...$	
	Continuous function with change of sign so root (in given interval)	A1 (2)
(b)	$f(0.4)=-0.1788$	B1
	$f(0.5)=...$	M1
	$f(0.5)=0.04114... \Rightarrow 0.4 \leqslant \alpha \leqslant 0.5$	A1 (3)
		[5]
	Notes	
	Must see correct values for the accuracy marks. But allow signs only for attempts at values for the method marks.	
(a)		
M1	Attempts both values, accept f(0.2)=... and f(0.6) =… with any values. (NB f(0.2) = −0.2069…, f(0.6) = 1.379 are the values with calculator in degrees mode.)	
A1	Both values correct (rounded or truncated to 1d.p.) and a correct conclusion (continuous and sign change). Allusion to continuity must be mentioned somewhere in the solution. Allow other ways to show sign change e.g. <0, >0 etc.	
(b)		
B1	Correct value of f (0.4); may be rounded or truncated to 1 dp	
M1	Attempt value of f (0.5) or attempt value of f(0.3) if relevant for their sign of f(0.4).	
A1	Correct value of f (0.5) which may be rounded to 2 dp and correct interval. Allow as open or closed interval. Accept any valid notation for the interval. Accept e.g. $0.4 < x < 0.5$	

Question Number	Scheme	Marks
2		
(a)	$\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}$	B1 (1)
(b)	$\left(x-\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}\right)\left(x-\frac{3}{8}+\frac{\sqrt{71}}{8}\mathrm{i}\right)\ \left((x-4)=0\right)$	
	$\left(x^2-\frac{3}{4}x+\frac{5}{4}\right)\ \left((x-4)=0\right)$	M1A1
	$x^3-\frac{19}{4}x^2+\frac{17}{4}x-5\ \ (=0)$	dM1
	$4x^3-19x^2+17x-20\ \ (=0)\qquad p=17,\ q=-20$	A1 (4)
		[5]
(a) **B1**	Correct answer only	
(b) **M1**	Attempt the multiplication of the 2 brackets with the complex terms. Allow $(x\pm\text{root})$ for the brackets. Allow "invisible" brackets.	
A1	Correct quadratic obtained - may have multiplied by the 4 (or other constant factor) and this is fine. (Need not be fully simplified but must have real terms)	
dM1	Attempt to multiply their quadratic by $(x-4)$ or may divide their quadratic into the cubic or other full method leading to at least one of p or q.	
A1	Correct values. Values of p and q need not be shown explicitly but may been seen in a cubic, provided the cubic starts $4x^3-19x^2$ (isw after a correct cubic)	
	Note if a candidate uses a hybrid method, mark under main scheme unless an Alt scores more marks.	
Alt 1 (b)	$-\frac{q}{4}=4\times\left(\frac{3}{8}+\frac{\sqrt{71}}{8}\mathrm{i}\right)\times\left(\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}\right)=\ldots\rightarrow q=..$ **or**	
	$\frac{p}{4}=4\left(\frac{3}{8}+\frac{\sqrt{71}}{8}\mathrm{i}\right)+4\left(\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}\right)+\left(\frac{3}{8}+\frac{\sqrt{71}}{8}\mathrm{i}\right)\left(\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}\right)\rightarrow p=\ldots$	M1
	$\Rightarrow q=-16\times\left(\frac{9}{64}+\frac{71}{64}\right)=-20$ **or** $p=17$	A1
	E.g. $\mathrm{f}(4)=0\Rightarrow 4(4)^3-19(4)^2+4p-"20"=0\Rightarrow p=\ldots$	
	or $\frac{p}{4}=4\left(\frac{3}{8}+\frac{\sqrt{71}}{8}\mathrm{i}\right)+4\left(\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}\right)+\left(\frac{3}{8}+\frac{\sqrt{71}}{8}\mathrm{i}\right)\left(\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}\right)\rightarrow p=\ldots$	dM1
	$p=17,\ q=-20$	A1 (4)

Question Number	Scheme	Marks
M1 **A1** **dM1** **A1**	A correct attempt to use product of roots is $-\frac{q}{4}$ to find a value for q **or** pair sum is $\frac{p}{4}$ to find a value of p. Correct value for p or q Correct full method to find both p and q. Correct values for both	
Alt 2	Attempts at using the factor theorem are possible but unlikely to succeed. Score as follows: M1: Uses the factor theorem to generate two equations in the two unknowns (note they will need to use a complex root to achieve this and equate real and imaginary parts.). A1: Correct equations. dM1: Solves their two equations to find values for p and q. A1: Correct values Send to review if unsure.	
Alt 3 **(b)**	$4x^2-3x+p-12$ $x-4\overline{\big)4x^3-19x^2+px+q}$ $4x^3-16x^2$ $-3x^2+px+q$ $-3x^2+12x$ $(p-12)x+q$ $(p-12)x-4(p-12)$ $\Rightarrow q+4(p-12)=0 \;\;\&\;\; \frac{p-12}{4}=\left(\frac{3}{8}+\frac{\sqrt{71}}{8}\mathrm{i}\right)\left(\frac{3}{8}-\frac{\sqrt{71}}{8}\mathrm{i}\right)=\frac{5}{4}$ $P=17,\ q=-20$	M1 A1 dM1 A1 (4)
(b) **M1** **A1** **dM1** **A1**	Divides $x-4$ into the cubic to achieve a 3TQ quotient and a remainder Correct quotient and remainder Correct full method to find p or q Correct values	

Question Number	Scheme	Marks
3(a)	$k(k+5)-6=0$	M1
	$k^2+5k-6=0$	
	$\left((k-1)(k+6)=0 \Rightarrow\right) \quad k=1,\ -6$	A1 (2)
(b)	$\dfrac{1}{"k^2+5k-6"}\begin{pmatrix} k & 2 \\ 3 & k+5 \end{pmatrix}$	M1A1 (2)
		[4]

(a)	
M1	Attempts determinant and sets equal to zero (or equivalent method) to obtain an unsimplified quadratic equation
A1	Correct values for k (may solve the quadratic by any valid means)
(b)	
M1	Forms the matrix of signed minors (must have at least three correct elements) divided or multiplied by an attempt at the determinant
A1	Fully correct inverse

Question Number	Scheme	Marks
4		
(a)	$\alpha+\beta=-\frac{5}{2} \quad \alpha\beta=\frac{7}{2}$	B1
	$\alpha^3+\beta^3=(\alpha+\beta)^3-3\alpha\beta(\alpha+\beta)=\left(-\frac{5}{2}\right)^3-3\left(\frac{7}{2}\right)\left(-\frac{5}{2}\right)$	M1
	$=\frac{85}{8}$	A1 (3)
(b)	$\frac{\alpha^2}{\beta}+\frac{\beta^2}{\alpha}=\frac{\alpha^3+\beta^3}{\alpha\beta}=...$	M1
	$=\left(\frac{85}{8}\right)\times\left(\frac{2}{7}\right)=\frac{85}{28}$	A1
	$\frac{\alpha^2}{\beta}\times\frac{\beta^2}{\alpha}=\alpha\beta=\frac{7}{2}$	B1ft
	$x^2-"\frac{85}{28}"x+"\frac{7}{2}" \ (=0)$	M1
	$28x^2-85x+98=0$	A1 (5) **[8]**

	Note: if a candidate solves the equation and uses the roots to answer the question, then send to review.
(a)	
B1	Both correct. (Seen anywhere in the working)
M1	Uses their sum and product of roots in a correct expression for $\alpha^3+\beta^3$.
A1	Correct value. Must be exact. Accept 10.625
(b)	
M1	$\alpha^3+\beta^3$ and $\alpha\beta$ into $\frac{\alpha^3+\beta^3}{\alpha\beta}=...$ substitutes their values for (allow slips in substitution).
A1	Correct sum as a single fraction (may be seen or implied in their equation)
B1ft	Correct product or follow through their product
M1	Use $x^2-\text{sum of roots}\times x+\text{product of roots}$ with their values for sum and product. "= 0" may be missing.
A1	A correct final equation as shown or any integer multiple of this. " = 0" must be included.

Question Number	Scheme	Marks
5(a)	$\sum_{r=1}^{n}(r+1)(r+5)=\sum_{r=1}^{n}(r^2+6r+5)$	B1
	$=\sum_{r=1}^{n}r^2+6\sum_{r=1}^{n}r+5n$	
	$=\frac{n}{6}(n+1)(2n+1)+6\frac{n}{2}(n+1)+5n$	M1A1
	$=\frac{n}{6}(2n^2+3n+1+18n+18+30)$	dM1
	$=\frac{n}{6}(2n^2+21n+49)=\frac{n}{6}(n+7)(2n+7)$ *	A1 * (5)
(b)	$\sum_{r=n+1}^{2n}=\sum_{r=1}^{2n}-\sum_{r=1}^{n}=\frac{2n}{6}(2n+7)(4n+7)-\frac{n}{6}(n+7)(2n+7)$	M1
	$=\frac{n}{6}(2n+7)\{8n+14-(n+7)\}$	
	$=\frac{7n}{6}(2n+7)(n+1)$	A1 (2)
		[7]

(a)	
B1	Brackets multiplied out correctly. Summation signs not needed.
M1	Use at least two correct formulae from $\sum_{r=1}^{n}r$, $\sum_{r=1}^{n}r^2$ and $\sum_{r=1}^{n}1=n$.
A1	Fully correct expression.
dM1	Attempt to remove factor $\frac{n}{6}$ from an expression with common factor n present. (if "$5n$" is just 5 then this mark will not be scored). Must be seen before the given answer is quoted. No need to simplify the remaining quadratic factor.
A1*	Obtain the correct 3 term quadratic and factorise. This is a "show that" question, so the 3 TQ must be seen. No errors seen.
(b)	
M1	Use $\sum_{r=n+1}^{2n}=\sum_{r=1}^{2n}-\sum_{r=1}^{n}$
A1	Simplify to the correct answer.

Question Number	Scheme	Marks
6(a)	$\lambda = 4$	B1 (1)
(b)	$\arctan\frac{3}{"4"}$ or $\arctan\frac{-3}{"4"}$	M1
	(Second quadrant so arg z = 2.498...) = 2.5 (rad)	A1 (2)
(c)(i)	$\frac{z+3\mathrm{i}}{2-4\mathrm{i}} = \frac{-4+6\mathrm{i}}{2-4\mathrm{i}} \times \frac{2+4\mathrm{i}}{2+4\mathrm{i}}$ or $\frac{z+3\mathrm{i}}{2-4\mathrm{i}} = a+\mathrm{i}b \Rightarrow -4+6\mathrm{i} = (a+\mathrm{i}b)(2-4\mathrm{i})$	M1
	$= \frac{-8+12\mathrm{i}-16\mathrm{i}+24\mathrm{i}^2}{4+16} = -\frac{8}{5}-\frac{1}{5}\mathrm{i}$ Accept e.g. $\frac{-32-4\mathrm{i}}{20}$	dM1A1
	Or $2a+4b=-4, 2b-4a=6 \Rightarrow a=..,b=..$	
(ii)	$z^2 = (-4+3\mathrm{i})^2 = 16-24\mathrm{i}+9\mathrm{i}^2 = 16-24\mathrm{i}-9$	M1
	$= 7-24\mathrm{i}$	A1ft (5)
(d)		B1 B1ft B1ft (3) **[11]**

(a) **B1**	Correct answer. No working needed.
(b) **M1**	For $\arctan\left(\pm\frac{3}{"4"}\right)$ with their "4". Can be awarded from $\tan\theta = \pm\frac{3}{"4"} \Rightarrow \theta = ...$ or by implication if correct value for either arctan or correct final answer (rounded or not rounded, may be degrees) is seen.
A1	Cao 2.5
(c)(i) **M1**	Multiplies numerator and denominator by complex conjugate of denominator. Award if denominator of 4+16 or 20 is seen instead of product. May still have z at this stage, or even allow with $\lambda + 3\mathrm{i}$ as numerator. Alternatively, sets equal to $a + \mathrm{i}b$ and cross multiplies.
dM1	Using their λ or 4 substitutes **correctly** for z, fully expands the numerator and uses $\mathrm{i}^2 = -1$ Alt, uses $\mathrm{i}^2 = -1$, equates real and imaginary terms and solves their equations for a and b
A1	Correct answer only, as shown or single fraction accepting equivalent fractions or with exact decimals (−1.6 − 0.2i).
(ii) M1	Squaring an expression of form $k+3\mathrm{i}$ (with a real value for k) to get 3 terms (may be implied) and uses $\mathrm{i}^2 = -1$
A1ft	Correct answer, follow through their $\lambda > 0$ (ie for $"\lambda^2 - 9" - "6\lambda"\mathrm{i}$ must be negative i term)
(d)	NB: Penalise once only (in the first mark due) for mislabelling or failing to label points as long as they look to be placed correctly. Award if lines/arrows not included. Points may be labelled by letter, name or their Cartesian coordinates (which may be given on the axes).
B1	Plots z in second quadrant and z^* as mirror image in the Real axis. Both must be labelled.
B1ft	Plot and label C for their solution to (c)(ii) It must be the correct side of B (for their

B1ft	answers) and a correct relative scale (so noticeably closer to O than their B if correct values). Plot and label their D (- 24 need not be to scale, but should be further from O than their B).

Question Number	Scheme	Marks
7(a)	$\begin{vmatrix} 4 & -5 \\ -3 & 2 \end{vmatrix} = 8-15=-7 \Rightarrow \text{Area } T' = "\pm 7" \times 23 = ...$	M1
	$\text{Area } T' = 161$	A1 (2)
(b)	$\begin{pmatrix} 4 & -5 \\ -3 & 2 \end{pmatrix}\begin{pmatrix} 3p+2 \\ 2p-1 \end{pmatrix} = \begin{pmatrix} 17 \\ -18 \end{pmatrix}$ or $\begin{pmatrix} 3p+2 \\ 2p-1 \end{pmatrix} = \frac{1}{8-15}\begin{pmatrix} 2 & 5 \\ 3 & 4 \end{pmatrix}\begin{pmatrix} 17 \\ -18 \end{pmatrix}$	
	$4(3p+2)-5(2p-1)=17$ or $-3(3p+2)+2(2p-1)=-18$ or $3p+2=-\frac{1}{7}(34-90)$ or $2p-1=-\frac{1}{7}(51-72)$ $(\text{e.g. } 2p+13=17 \Rightarrow .....)$	M1
	$p=2$	A1 (2)
(c)	Rotation; through 90° clockwise (or 270° anticlockwise) about origin	B1;B1 (2)
(d)	**CA** = **B** $\mathbf{A}^{-1} = -\frac{1}{7}\begin{pmatrix} 2 & 5 \\ 3 & 4 \end{pmatrix}$ or $\begin{pmatrix} a & b \\ c & d \end{pmatrix}\begin{pmatrix} 4 & -5 \\ -3 & 2 \end{pmatrix} = \begin{pmatrix} 4a-3b & -5a+2b \\ 4c-3d & -5c+2d \end{pmatrix}$	B1
	$\mathbf{C} = -\frac{1}{7}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\begin{pmatrix} 2 & 5 \\ 3 & 4 \end{pmatrix} = ...$ or $\begin{cases} 4a-3b=0 & -5a+2b=1 \\ 4c-3d=-1 & -5c+2d=0 \end{cases} \Rightarrow ...$	M1
	$\mathbf{C} = -\frac{1}{7}\begin{pmatrix} 3 & 4 \\ -2 & -5 \end{pmatrix}$ or $\begin{pmatrix} -\frac{3}{7} & -\frac{4}{7} \\ \frac{2}{7} & \frac{5}{7} \end{pmatrix}$ oe	A1 (3)
		[9]

(a)	
M1	Attempts to find the determinant of **M** and use as a scale factor. Accept if a slip in calculation is made and accept if negative is used for this mark. Dividing by the determinant is M0.
A1	Correct answer only. No working needed (correct answer implies the method).
(b)	
M1	Form a matrix equation using either **A** or an attempt at $\mathbf{A}^{-1}$, obtain a linear equation and solve for *p*.
A1	Correct value for *p*, obtained from a correct equation. (No need to check in other equation.)
(c)	
B1	Rotation, rotates, rotate or rotating (oe) Accept "turn"
B1	Correct angle (degrees or radians) with direction specified and about origin or (0, 0)
(d)	
B1	Correct matrix for $\mathbf{A}^{-1}$ May have been found in (b) but must be used in (d). Alternatively, correct **CA** with unknowns for entries of **C**.
M1	Multiply **B** by $\mathbf{A}^{-1}$ **on the right.** Alternatively, sets **CA** equal to **B** and solves equations.
A1	Correct matrix **C**(isw after a correct answer).

Question Number	Scheme	Marks
8(a)	$200t^3 = 25$ or $\left(\frac{25}{x}\right)^2 = 40x$ or $y^2 = 40\left(\frac{25}{y}\right)$	M1
	$t = \frac{1}{2}$ or $x = \frac{5}{2}$ or $y = 10$	A1
	$\left(\frac{5}{2}, 10\right)$	A1 (3)
(b)	$y^2 = 40x \Rightarrow x = \frac{y^2}{40} \Rightarrow \frac{dx}{dy} = \frac{2y}{40}$ or $2y\frac{dy}{dx} = 40 \Rightarrow \frac{dy}{dx} = \frac{40}{2y}$ or $\frac{dy}{dx} = \frac{\sqrt{10}}{\sqrt{x}}$	B1
	or by parametric differentiation: $\frac{dy}{dx} = \frac{1}{t}$	
	at $(10, 20)$: $\frac{dy}{dx} = 1$ or $\frac{dx}{dy} = 1$	M1
	Grad normal $= -1$	A1
	$y - 20 = -(x - 10)$	M1
	$x + y - 30 = 0$	A1 (5)
(c)	$xy = 25 \quad x + y - 30 = 0$	
	$x + \frac{25}{x} - 30 = 0$ or $\frac{25}{y} + y - 30 = 0$ or $\frac{5}{t} + 5t - 30 = 0$	M1
	$x^2 - 30x + 25 = 0$ or $y^2 - 30y + 25 = 0$ or $5t^2 - 30t + 5 = 0$	dM1
	$x = 15 \pm \sqrt{200}$ or $y = 15 \pm \sqrt{200}$ or $t = 3 \pm 2\sqrt{2}$ (or exact equivalents)	A1
	eg $x = 15 + \sqrt{200} \Rightarrow y = 15 - \sqrt{200}$	ddM1
	$\left(15 + 10\sqrt{2}, 15 - 10\sqrt{2}\right) \quad \left(15 - 10\sqrt{2}, 15 + 10\sqrt{2}\right)$	A1A1 (6)
		[14]

(a)	
M1	Attempt an equation in a single variable.
A1	Correct value for x, y or t
A1	Correct values for x and y. Need not be in coordinate brackets. No other points seen.
(b)	
B1	Any correct expression involving the derivative, $\frac{dy}{dx}$ or $\frac{dx}{dy}$, for P
M1	Attempt to obtain value of their derivative at (10, 20). May be from an incorrect curve.
A1	Correct gradient of normal.
M1	Equation of normal by any complete method. Must involve a sign change of their derivative and have numerical gradient. Cam be scored from an incorrect starting equation/point.
A1	Correct equation in form demanded (though terms may be in different order).
(c)	
M1	Use the equation of H and their equation of the normal from (b) to obtain an equation in a single variable
dM1	Obtains a 3TQ and attempts to solve by any valid means.
A1	Correct values for x or y or t. Must be exact but need not be fully simplified (but discriminant must be evaluated).
ddM1	Use at least one of their values for x or y to obtain a value for the other coordinate or t to find at least one set of coordinates. (Can be scored with inexact values.)
A1	Either pair of coordinates correct. Allow if unsimplified.
A1	Second pair of coordinates correct and no extra solutions and both pairs in simplest form (as shown in scheme). Need not be coordinates as long as correctly paired.
	Award A1A0 if $x = 15 \pm 10\sqrt{2}, y = 15 \pm 10\sqrt{2}$ is given

Question Number	Scheme	Marks
9(i)	$n=1 \quad u_1 = 3\times\frac{2}{3}-1=1$ (so true for n = 1 (†))	B1
	Assume true for $n=k$ ie $u_k = 3\left(\frac{2}{3}\right)^k - 1$ (†)	
	$u_{k+1} = \frac{1}{3}(2u_k - 1) = \frac{1}{3}\left(2\left(3\left(\frac{2}{3}\right)^k - 1\right) - 1\right) = \frac{1}{3}\left(6\left(\frac{2}{3}\right)^k - 2 - 1\right)$	M1A1
	$= \frac{1}{3}\left(2\times 3\left(\frac{2}{3}\right)^{k+1} \times \left(\frac{3}{2}\right) - 2 - 1\right)$	dM1
	$= \frac{1}{3}\times 2\times 3\left(\frac{2}{3}\right)^{k+1} \times \left(\frac{3}{2}\right) + \frac{1}{3}(-2-1)$	
	$= 3\left(\frac{2}{3}\right)^{k+1} - 1$	A1
	$\therefore$ if true for $n=k$, also true for $n=k+1$ (†)	
	$\left(\text{True for } n=1\right)$ so $u_n = 3\left(\frac{2}{3}\right)^n - 1$ is true for all $n \in \mathbb{Z}^+$	A1cso (6)
(ii)	$\mathrm{f}(1) = 2^3 + 3^3 = 8 + 27 = 35$ (Multiple of 7) (so true for n = 1 (†))	B1
	Assume $\mathrm{f}(k)$ is a multiple of 7 $\mathrm{f}(k) = 2^{k+2} + 3^{2k+1}$ is a multiple of 7 (†)	
	$\mathrm{f}(k+1) - M\mathrm{f}(k) = 2^{k+3} + 3^{2k+3} - M\left(2^{k+2} + 3^{2k+1}\right)$	M1
	$= 2^{k+2}(2-M) + 3^{2k+1}\left(3^2 - M\right)$	A1
	$= (2-M)\left(2^{k+2} + 3^{2k+1}\right) + 3^{2k+1}\times 7$ or $(9-M)\left(2^{k+2} + 3^{2k+1}\right) - 7\times 2^{k+2}$ oe	dM1
	$\therefore \mathrm{f}(k+1) = 2\mathrm{f}(k) + 7\times 3^{2k+1}$ oe e.g. $9\mathrm{f}(k) - 7\times 2^{k+2}$	A1
	Or e.g. $7\times 3^{2k+1}$ is a multiple of 7, so if f(k) is a multiple of 7 <u>then f(k+1) is also a multiple of 7</u>	
	If the result is true for $n = k$ it is also true for $n = k + 1$ (†)	
	As the result has been shown to be true for n = 1, it is true for all $n \in \mathbb{Z}^+$	A1 cso (6) **[12]**

9(i)	
B1	Check that the formula gives 1 when $n = 1$ Working must be shown. (Need not state true for $n = 1$ for this mark – but see final A)
M1	(Assume true for $n = k$ and) attempts to substitute the formula for u_k into $u_{k+1} = \frac{1}{3}(2u_k - 1)$ or equivalent with suffixes increased. Allow slips.
A1	Correct substitution.
dM1	Obtain an expression with $\left(\frac{2}{3}\right)^{k+1}$ and no other k. Alternatively, expands u_{k+1} to a matching expression (ie work from both directions).
A1	Correct expression when $n = k + 1$ At least one intermediate stage of working must be shown and no errors (though notational slips may be condoned). If working from both directions, it is for correct work to reach matching expressions.
A1cso	Correct concluding statements following correct solution which has included each of the points (†) at some stage during the working. Depends on all except the first B mark (e.g. if they think they have checked $n = 1$ but have really checked $n = 2$). Note: Allow the M's and first two A's for students who go from k+1 to k+2 but treat it as k to $k + 1$.
(ii)	
B1	Checks the case $n = 1$. Minimum statement of f(1) = 35
M1	Attempts an expression for f(k + 1) – Mf(k) with any value of M. Need not be simplified. Most likely with $M = 1$ but may be seen with other values of M. With $M = 0$, $\mathrm{f}(k+1) = 2^{k+3} + 3^{2k+3}$ is all that is required.
A1	A correct expression with terms 2^{k+2} and 3^{2k+1} clearly identified.
dM1	Attempts to extract/identify f(k) within a correct expression to give terms divisible by 7. With $M = 0$ look for $\mathrm{f}(k+1) = 2\times\left(2^{k+2} + 3^{3k+1}\right) + 7\times 3^{2k+1}$ or $9\times\left(2^{k+2} + 3^{3k+1}\right) - 7\times 2^{k+2}$ oe and similar for other value of M.
A1	One of the correct expressions for f(k+1) shown (or with powers of 2 and 3) or full reason why f(k+1) is divisible by 7, following a suitable expression.
A1cso	Correct concluding statements following correct solution which has included each of the points (†) at some stage during the working. Depends on all previous marks.

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom