Version 1.4



General Certificate of Education (A-level)
January 2013

Chemistry CHEM5

(Specification 2420)

Unit 5: Energetics, Redox and Inorganic Chemistry

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2013 AQA and its licensors. All rights reserved.

Copyright
AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334).
Registered address: AQA, Devas Street, Manchester M15 6EX.

Question	Marking Guidance	Mark	Comments
1(a)	(Enthalpy change to) break the bond in 1 mol of chlorine (molecules)	1	Allow (enthalpy change to) convert 1 mol of chlorine molecules into atoms Do not allow energy or heat instead of enthalpy, allow heat energy
	To form (2 mol of) gaseous chlorine atoms / free radicals	1	Can score 2 marks for 'Enthalpy change for the reaction': $Cl_2(g) \rightarrow 2Cl(g)$ Equation alone gains M2 only Can only score M2 if 1 mol of chorine molecules used in M1 (otherwise it would be confused with atomisation enthalpy) Any mention of ions, CE = 0
1(b)	(For atomisation) only 1 mol of chlorine atoms, not 2 mol (as in bond enthalpy) is formed / equation showing ½ mol Chlorine giving 1 mol of atoms	1	Allow breaking of one bond gives two atoms Allow the idea that atomisation involves formation of 1 mol of atoms not 2 mol Allow the idea that atomisation of chlorine involves half the amount of molecules of chlorine as does dissociation Any mention of ions, CE = 0
1(c)(i)	$\frac{1}{2}F_2(g) + \frac{1}{2}Cl_2(g) \rightarrow ClF(g)$	1	

1(c)(ii)	ΔH = ½E(F–F) + ½ E(Cl-Cl) – E(Cl-F) E(Cl-F) = ½E(F–F) + ½E(Cl-Cl) - ΔH = 79 + 121 – (–56) = 256 (kJ mol^{-1})	1 1	Allow correct cycle -256 scores zero Ignore units even if wrong
1(c)(iii)	½Cl_2 + 3/2F_2 → ClF_3 ΔH = ½ E(Cl-Cl) + 3/2 E(F–F) - 3E(Cl-F) = 121 + 237 – 768 / (or 3 x value from (c)(ii)) = –410 (kJ mol^{-1})	1 1 1	If equation is doubled CE=0 unless correcr answer gained by /2 at end This would score M1 This also scores M1 (note = 358 – 768) If given value of 223 used ans = –311 Allow 1/3 for +410 and +311
1(c)(iv)	(Bond enthalpy of) Cl-F bond in ClF is different from that in ClF_3	1	Allow Cl-F bond (enthalpy) is different in different compounds (QoL)
1(d)	NaCl is ionic / not covalent	1	

Question	Marking Guidance	Mark	Comments
2(a)	$MgCl_2(s) \rightarrow Mg^{2+}(g) + 2Cl^-(g)$	1	
2(b)	The magnesium <u>ion</u> is smaller / has a smaller radius / greater charge density (than the calcium ion) Attraction between ions / to the chloride ion stronger	1 1	If not ionic or if molecules / IMF / metallic / covalent / bond pair / electronegativity mentioned, CE = 0 Allow ionic bonds stronger Do not allow any reference to polarisation or covalent character Mark independently
2(c)	The oxide ion has a greater charge / charge density than the chloride ion So it attracts the magnesium ion more strongly	1 1	If not ionic or if molecules / IMF / metallic / covalent / bond pair mentioned, CE = 0 Allow oxide ion smaller than chloride ion Allow ionic bonds stronger Mark independently
2(d)	$\Delta H_{solution} = \Delta H_L + \Sigma\Delta H_{hyd}$ Mg^{2+} ions + $\Sigma\Delta H_{hyd}$ Cl^- ions $-155 = 2493 + \Delta H_{hyd}$ Mg^{2+} ions $- 2\times364$ ΔH_{hyd} Mg^{2+} ions $= -155 - 2493 + 728$ $= -1920$ (kJ mol^{-1})	1 1 1	Allow correct cycle Ignore units Allow max 1 for +1920 Answer of + or -1610, CE = 0 Answer of -2284, CE = 0

2(e)	Water is polar / O on water has a delta negative charge Mg^{2+} ion / +ve ion / + charge attracts (negative) O on a water molecule	1 1	Allow <u>O</u> (not water) has lone pairs (can score on diagram) Allow Mg^{2+} attracts lone pair(s) M2 must be stated in words (QoL) Ignore mention of co-ordinate bonds CE = 0 if O^{2-} or water ionic or H bonding

2(f)	Magnesium oxide reacts with water / forms $Mg(OH)_2$	1	Allow MgO does not dissolve in water / sparingly soluble / insoluble

Question	Marking Guidance	Mark	Comments
3(a)	$\Delta G = \Delta H - T\Delta S$	1	Or expression $\Delta H - T\Delta S$ must be evaluated
	If ΔG / expression <=0 reaction is feasible	1	Or any explanation that this expression <=0 Do not allow just $\Delta G = 0$
3(b)	The molecules become more disordered / random when water changes from a liquid to a gas / evaporates	1	For M1 must refer to change in state AND increase in disorder
	Therefore the entropy change is positive / Entropy increases	1	Only score M2 if M1 awarded
	$T\Delta S > \Delta H$	1	Allow M3 for *T* is large / high (provided M2 is scored)
	$\Delta G < 0$	1	Mark M3, M4 independently
3(c)(i)	Condition is $T = \Delta H/\Delta S$	1	
	ΔS = 189 –205/2 – 131 = –44.5;	1	
	ΔH = –242 therefore T = (–242 × 1000)/–44.5)	1	
	= 5438 K (allow 5400 – 5500 K)	1	Units essential (so 5438 alone scores 3 out of 4) 2719 K allow score of 2 5.4 (K) scores 2 for M1 and M2 only 1646 (K) scores 1 for M1 only
3(c)(ii)	It would decompose into <u>hydrogen and oxygen</u> / its elements	1	Can score this mark if mentioned in M2
	Because ΔG for this reaction would be <= 0	1	Allow the reverse reaction / decomposition is feasible Only score M2 if M1 awarded

3(d)	$\Delta H = T\Delta S$	1	Allow correct substituted values instead of symbols
	ΔS = 70-189 = -119 $J K^{-1} mol^{-1}$	1	
	ΔH = (-119 × 373)/1000 = -44.4 kJ (mol^{-1}) (allow -44 to -45)	1	Allow -44000 to -45000 J (mol^{-1}) Answer must have correct units of kJ or J

Question	Marking Guidance	Mark	Comments
4(a)	Na_2O is an ionic <u>lattice</u> / giant ionic / ionic crystal	1	CE= 0 if molecules, atoms, metallic mentioned Mention of electronegativity max 1 out of 2
	With strong forces of attraction between ions	1	Allow strong ionic bonds/lots of energy to separate ions
4(b)	SO_3 is a larger molecule than SO_2	1	Allow greater M_r / surface area
	So <u>van der Waals'</u> forces <u>between molecules</u> are stronger	1	Any mention of ions, CE= 0
4(c)	Ionic	1	Do not allow ionic with covalent character
	Contains <u>O^{2-}</u> ions / oxide ions	1	Equations of the form $O^{2-} + H^+ \rightarrow OH^-$ / $O^{2-} + 2H^+ \rightarrow H_2O$ / $O^{2-} + H_2O \rightarrow 2OH^-$ score M2 and M3
	These / O^{2-} ions (accept protons to) form OH^- / hydroxide / water (must score M2 to gain M3)	1	
4(d)(i)	$SO_2 + H_2O \rightarrow H^+ + HSO_3^-$	1	Allow $2H^+ + SO_3^{2-}$ but no ions, no mark Only score (d)(ii) if (d)(i) correct
4(d)(ii)	Reaction is an equilibrium / reversible reaction displaced mainly to the left / partially ionised / dissociated	1	Allow reaction does not go to completion
4(e)	SiO_2 reacts with bases / NaOH / CaO / $CaCO_3$	1	Ignore incorrect formulae for silicate



Question	Marking Guidance	Mark	Comments
5(a)	Yellow (solution)	1	
	Orange <u>solution</u>	1	
	$2CrO_4^{2-} + 2H^+ \rightarrow Cr_2O_7^{2-} + H_2O$	1	Allow equation with H_2SO_4
5(b)	Yellow / purple (solution)	1	Allow orange / brown (solution)
	Brown precipitate / solid	1	
	$[Fe(H_2O)_6]^{3+} + 3OH^- \rightarrow Fe(H_2O)_3(OH)_3 + 3H_2O$	1	
5(c)	Blue (solution)	1	Allow pale blue
	Dark / deep blue <u>solution</u>	1	Ignore any reference to blue ppt
	$[Cu(H_2O)_6]^{2+} + 4NH_3 \rightarrow [Cu(H_2O)_2(NH_3)_4]^{2+} + 4H_2O$	1	Can be in two equations
5(d)	Colourless (solution)	1	
	White precipitate / solid	1	Do not allow grey
	Bubbles / effervescence / gas evolved / given off	1	Do not allow just CO_2
	$2[Al(H_2O)_6]^{3+} + 3CO_3^{2-} \rightarrow 2Al(H_2O)_3(OH)_3 + 3CO_2 + 3H_2O$	1	

Question	Marking Guidance	Mark	Comments
6(a)	Variable / many oxidation states	1	
6(b)	$V_2O_5 + SO_2 \rightarrow V_2O_4 + SO_3$ $V_2O_4 + ½O_2 \rightarrow V_2O_5$	1 1	Equations can be in either order Allow multiples
6(c)(i)	In a different phase / state <u>from reactants</u>	1	
6(c)(ii)	Impurities poison / deactivate the catalyst / block the active sites	1	Allow (adsorbs onto catalyst AND reduces surface area)
6(d)(i)	The catalyst is a reaction product	1	
6(d)(ii)	Mn^{2+}/ Mn^{3+} ion(s)	1	
6(d)(iii)	$4Mn^{2+} + MnO_4^- + 8H^+ \rightarrow 5Mn^{3+} + 4H_2O$ $2Mn^{3+} + C_2O_4^{2-} \rightarrow 2Mn^{2+} + 2CO_2$	1 1	Equations can be in either order

Question	Marking Guidance	Mark	Comments
7(a)	Diagram of an Fe^{3+}/Fe^{2+} electrode that includes the following parts labelled:		
	Solution containing Fe^{2+} and Fe^{3+} ions	1	
	Platinum electrode connected to one terminal of a voltmeter	1	Must be in the solution of iron ions (one type will suffice)
	Salt bridge	1	Do not allow incorrect material for salt bridge and salt bridge must be in the solution (ie it must be shown crossing a meniscus)
	298 K and 100 kPa / 1 bar	1	
	<u>all solutions</u> unit / 1 mol dm^{-3} concentration	1	Allow zero current / high resistanve voltmeter as alternative to M4 or M5 Ignore hydrogen electrode even if incorrect
7(b)	$Cu^{2+} + Fe \rightarrow Cu + Fe^{2+}$	1	Ignore state symbols
	$Fe\|Fe^{2+}\|\|Cu^{2+}\|Cu$ correct order	1	Allow $Cu\|Cu^{2+}\|\|Fe^{2+}\|Fe$
	Phase boundaries and salt bridge correct, no Pt	1	Allow single/double dashed line for salt bridge Penalise phase boundary at either electrode end Can only score M3 if M2 correct
	Copper electrode	1	Allow any reference to copper

7(c)	$E^{\ominus}$ Au^{+}(/Au) > $E^{\ominus}$ O_2(/H_2O)	1	Allow *E* cell/e.m.f. = 0.45 V Allow 1.68 > 1.23
	So Au^{+} ions will oxidise water / water reduces Au^{+}	1	QoL
	$2Au^{+} + H_2O \rightarrow 2Au + 1/2O_2 + 2H^{+}$	1	Allow multiples

7(d)	$E^{\ominus}$ Ag^{+}(/Ag) > $E^{\ominus}$ Fe^{2+}(/Fe)	1	Allow *E* cell/e.m.f. = 1.24 Allow 0.80 > -0.44
	And $E^{\ominus}$ Ag^{+}(/Ag) > $E^{\ominus}$ Fe^{3+}(/Fe^{2+})	1	Allow *E* cell/e.m.f. = 0.03 Allow 0.80 > 0.77
	So silver ions will oxidise iron (to iron(II) ions) and then oxidise Fe(II) ions (further to Fe(III) ions producing silver metal)	1	Allow Ag^{+} ions will oxidise iron to iron(III)

Question	Marking Guidance	Mark	Comments
8(a)	A ligand is an electron pair / lone pair donor	1	Allow uses lone / electron pair to form a co-ordinate bond
	A bidentate ligand donates two electron pairs (to a transition metal ion) from different atoms / two atoms (on the same molecule / ion)	1	QoL
8(b)	$CoCl_4^{2-}$ diagram 	1	Four chlorines attached to Co with net 2- charge correct
	Tetrahedral shape	1	Charge can be placed anywhere, eg on separate formula Penalise excess charges
	109^o 28’	1	Allow 109^o to 109.5^o
	$[Co(NH_3)_6]^{2+}$ diagram 	1	Six ammonia / NH_3 molecules attached to Co with 2+ charge correct
	Octahedral shape	1	
	90^o	1	Allow 180^o if shown clearly on diagram
			CE= 0 if wrong complex but mark on if only charge is incorrect
8(c)	In different complexes the d orbitals / d electrons (of the cobalt) will have different energies / d orbital splitting will be different	1	
	Light / energy is absorbed causing an electron to be excited	1	
	Different frequency / wavelength / colour of light will be absorbed / transmitted / reflected	1	

8(d)	1 mol of H_2O_2 oxidises 2 mol of Co^{2+}	1	Or $H_2O_2 + 2Co^{2+} \rightarrow 2OH^- + 2Co^{3+}$
	M_r $CoSO_4.7H_2O$ = 281	1	If M_r wrong, max 3 for M1, M4, M5
	Moles Co^{2+} = 9.87/281 = 0.03512	1	
	Moles H_2O_2 = 0.03512/2 = 0.01756	1	M4 is method mark for (M3)/2 (also scores M1)
	Volume H_2O_2 = (moles × 1000)/concentration = 0.01756 × 1000)/5.00		
	= 3.51 cm^3 / (3.51 x 10^{-3} dm^3)	1	Units essential for answer
			M5 is method mark for (M4) x 1000/5 Allow 3.4 to 3.6 cm^3
			If no 2:1 ratio or ratio incorrect Max 3 for M2, M3 & M5
			Note : Answer of 7 cm^3 scores 3 for M2, M3, M5 (and any other wrong ratio max 3)
			Answer of 16.8 cm^3 scores 3 for M1, M4, M5 (and any other wrong M_r max 3)
			Answer of 33.5 cm^3 scores 1 for M5 only (so wrong M_r AND wrong ratio max 1)