Version 1.0



General Certificate of Education (A-level)
June 2013

Chemistry CHEM5

(Specification 2420)

Unit 5: Energetics, Redox and Inorganic Chemistry

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2013 AQA and its licensors. All rights reserved.

Copyright
AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334).
Registered address: AQA, Devas Street, Manchester M15 6EX.

Question	Marking Guidance	Mark	Comments
1(a)	Enthalpy change (to separate)1 mol of an (ionic) substance into its ions	1	If ionisation or hydration / solution, CE = 0 If atoms / molecules / elements mentioned, CE = 0 Allow heat energy change but not energy change alone. If forms 1 mol ions, lose M1
	Forms ions in the gaseous state	1	If lattice formation not dissociation, allow M2 only. Ignore conditions. Allow enthalpy change for $MX(s) \rightarrow M^+(g) + X^-(g)$ (or similar) for M1 and M2
1(b)	Any **one** of: • Ions are point charges • Ions are perfect spheres • Only electrostatic attraction / bonds (between ions) • No covalent interaction / character • Only ionic bonding / no polarisation of ions	1 max	If atoms / molecules mentioned, CE = 0
1(c)	(Ionic) radius / distance between ions / size	1	Allow in any order. Do not allow charge / mass or mass / charge.
	(Ionic) charge / charge density	1	Do not allow 'atomic radius'.

1(d)	$\Delta H_L = \Delta H_a$(chlorine) + ΔH_a(Ag) + I.E(Ag) +EA(Cl) - $\Delta H_f^{\ominus}$ = 121 + 289 + 732 -364 + 127 = (+) 905 (kJ mol^{-1})	1 1 1	Or cycle If $AgCl_2$, CE=0/3 Allow 1 for -905 Allow 1 for (+)844.5 (use of 121/2) Ignore units even if incorrect.
1(e)	**M1** Greater **M2** (Born-Haber cycle method allows for additional) covalent interaction ***OR*** **M1** Equal **M2** AgCl is perfectly ionic / no covalent character	1 1	Do not penalise $AgCl_2$ Allow AgCl has covalent character. Only score M2 if M1 is correct.

Question	Marking Guidance	Mark	Comments
2(a)	Chloride (ions) are smaller (than bromide ions)	1	Must state or imply ions. Allow chloride has greater charge density (than bromide). Penalise chlorine ions once only (max 2/3).
	So the force of attraction between chloride ions and water is stronger	1	This can be implied from M1 and M3 but do not allow intermolecular forces.
	Chloride ions attract the δ+ on H of water / electron deficient H on water	1	Allow attraction between ions and polar / dipole water. Penalise H^+ (ions) and mention of hydrogen bonding for **M3** Ignore any reference to electronegativity. Note: If water not mentioned can score M1 only.
2(b)	$\Delta H_{solution} = \Delta H_L + \Delta H_{hyd}$ K^+ ions + ΔH_{hyd} Br^- ions / = 670 – 322 – 335	1	Allow $\Delta H_{solution} = \Delta H_L + \Sigma\Delta H_{hyd}$
	= (+)13 (kJ mol^{-1})	1	Ignore units even if incorrect. +13 scores M1 and M2 -13 scores 0 -16 scores M2 only (transcription error).

2(c)(i)	The entropy change is positive / entropy increases	1	ΔS is negative loses M1 and M3
	Because 1 mol (solid) $\rightarrow$ 2 mol (aqueous ions) / no of particles increases	1	Allow the aqueous ions are more disordered (than the solid). Mention of atoms / molecules loses M2
	Therefore $\underline{T\Delta S > \Delta H}$	1	
2(c)(ii)	Amount of KCl = $5/M_r$ = 5/74.6 = $\underline{0.067(0)\ \text{mol}}$	1	If moles of KCl not worked out can score M3, M4 only (answer to M4 likely to be 205.7 K)
	Heat absorbed = 17.2 × 0.0670 = 1.153 kJ	1	Process mark for M1 × 17.2
	Heat absorbed = mass × sp ht × ΔT $(1.153 \times 1000) = 20 \times 4.18 \times \Delta T$	1	If calculation uses 25 g not 20, lose M3 only (M4 = 11.04, M5 = 287)
	ΔT = 1.153 × 1000 / (20 × 4.18) = 13.8 K	1	If 1000 not used, can only score M1, M2, M3 M4 is for a correct ΔT Note that 311.8 K scores 4 (M1, M2, M3, M4).
	T = 298 – 13.8 = 284(.2) K	1	If final temperature is negative, M5 = 0 Allow no units for final temp, penalise wrong units.

Question	Marking Guidance	Mark	Comments
3(a)(i)	(At 0 K) particles are stationary / not moving / not vibrating	1	Allow have zero energy. Ignore atoms / ions.
	No disorder / perfect order / maximum order	1	Mark independently.
3(a)(ii)	As *T* increases, particles start to move / vibrate	1	Ignore atoms / ions. Allow have more energy. If change in state, CE = 0
	Disorder / randomness increases / order decreases	1	
3(a)(iii)	Mark on temperature axis vertically below second 'step'	1	Must be marked as a line, an 'x' , T_b or 'boiling point' on the temperature axis.
3(a)(iv)	L_2 corresponds to boiling / evaporating / condensing / $l \rightarrow g$ / $g \rightarrow l$ And L_1 corresponds to melting / freezing / $s \rightarrow l$ / $l \rightarrow s$	1	There must be a clear link between L_1, L_2 and the change in state.
	Bigger change in disorder for L_2 / boiling compared with L_1 / melting	1	M2 answer must be in terms of changes in state and not absolute states eg must refer to change from liquid to gas not just gas. Ignore reference to atoms even if incorrect.

3(b)(i)	$\Delta G = \Delta H - T\Delta S$ $\Delta H = c$ and $(-)\Delta S = m$ / ΔH and ΔS are constants (approx)	1 1	 Allow ΔH is the intercept, and $(-)\Delta S$ is the slope / gradient. Can only score M2 if M1 is correct.
3(b)(ii)	Because the entropy change / ΔS is positive / $T\Delta S$ gets bigger	1	Allow $-T\Delta S$ gets more negative.
3(b)(iii)	Not feasible / unfeasible / not spontaneous	1	
3(c)(i)	+ 44.5 J K^{-1} mol^{-1}	1	Allow answer without units but if units given they must be correct (including mol^{-1})
3(c)(ii)	At 5440 $\Delta H = T\Delta S$ = 5440 × 44.5 = 242 080 (***OR*** using given value = 5440 × 98 = 533 120) ΔH = 242 kJ mol^{-1} (***OR*** using given value ΔH = 533 kJ mol^{-1})	1 1 1	 Mark is for answer to (c)(i) × 5440 Mark is for correct answer to M2 with correct units (J mol^{-1} or kJ mol^{-1}) linked to answer. If answer consequentially correct based on (c)(i) except for incorrect sign (eg -242), max 1/3 provided units are correct.

Question	Marking Guidance	Mark	Comments
4(a)	MgO is ionic	1	If not ionic, CE = 0
	Melt it	1	If solution mentioned, cannot score M2 or M3
	(Molten oxide) conducts electricity	1	Allow acts as an electrolyte. Cannot score M3 unless M2 is correct.
4(b)	Macromolecular	1	CE = 0 if ionic, metallic or molecular. Allow giant molecule.
	Covalent bonding	1	Giant covalent scores M1 and M2
	Water cannot (supply enough energy to) break the covalent bonds / lattice	1	Hydration enthalpy < bond enthalpy.
4(c)	(Phosphorus pentoxide's melting point is) lower	1	If M1 is incorrect, can only score M2
	Molecular with covalent bonding	1	M2 can be awarded if molecular mentioned in M3
	Weak / easily broken / not much energy to break intermolecular forces ***OR*** weak vdW / dipole-dipole forces of attraction between molecules	1	Intermolecular / IMF means same as between molecules.

4(d)	Reagent (water or acid) Equation eg $MgO + 2HCl \rightarrow MgCl_2 + H_2O$	1 1	Can be awarded in the equation. $MgO + H_2O \rightarrow Mg(OH)_2$ Equations can be ionic but must show all of the reagent eg $H^+ + Cl^-$ Simplified ionic equation without full reagent can score M2 only. Allow $6MgO + P_4O_{10} \rightarrow 2Mg_3(PO_4)_2$
4(e)	$P_4O_{10} + 12NaOH \rightarrow 4Na_3PO_4 + 6H_2O$	1	Allow P_2O_5 and acid salts. Must be NaOH not just hydroxide ions.

Question	Marking Guidance	Mark	Comments
5(a)	It has mobile ions / ions can move through it / free ions	1	Do not allow movement of electrons. Allow specific ions provided they are moving but do not react.
5(b)	Chloride ions react with copper ions / Cu^{2+} ***OR*** $[CuCl_4]^{2-}$ formed	1	If incorrect chemistry, mark = 0
5(c)	The Cu^{2+} ions / $CuSO_4$ in the left-hand electrode more concentrated So the reaction of Cu^{2+} with $2e^-$ will occur (in preference at) left-hand electrode / $Cu \rightarrow Cu^{2+}$ + electrons at right-hand electrode	1 1	Allow converse. Allow left-hand electrode positive / right-hand electrode negative. Also reduction at left-hand electrode / oxidation at right-hand electrode. Also left-hand electrode has oxidising agent / right-hand electrode has reducing agent. Allow *E* left-hand side > *E* right-hand side
5(d)	(Eventually) the copper ions / $CuSO_4$ in each electrode will be at the same concentration	1	
5(e)(i)	-3.05 (V)	1	Must have minus sign. -3.05 only.

5(e)(ii)	$LiMnO_2 \rightarrow Li + MnO_2$ correct equation Correct direction	1 1	Allow 1 for reverse equation. Allow multiples. If Li+ not cancelled but otherwise correct, max = 1 If electrons not cancelled, CE = 0 $LiMnO_2 \rightarrow Li + MnO_2$ scores 2 $Li^+ + LiMnO_2 \rightarrow Li^+ + Li + MnO_2$ scores 1 $Li + MnO_2 \rightarrow LiMnO_2$ scores 1

5(e)(iii)	Electricity for recharging the cell may come from power stations <u>burning</u> (fossil) fuel	1	Allow any reference to <u>burning</u> (of carbon-containing) fuels. Note combustion = burning.

Question	Marking Guidance	Mark	Comments
6(a)	$\Delta E = h\nu$	1	Allow = h*f*
	$\nu = \Delta E / h = 2.84 \times 10^{-19} / 6.63 \times 10^{-34} = 4.28 \times 10^{14}$ s^{-1} / Hz	1	Allow 4.3×10^{14} s^{-1} / Hz Answer must be in the range: $4.28 - 4.30 \times 10^{14}$
6(b)	(One colour of) light is absorbed (to excite the electron)	1	If light emitted, CE = 0
	The remaining colour / frequency / wavelength / energy is transmitted (through the solution)	1	Allow light reflected is the colour that we see.
6(c)	Bigger	1	
	Blue light would be absorbed ***OR*** light that has greater energy than red light would be absorbed ***OR*** higher frequency (of light absorbed / blue light) leads to higher ΔE	1	Can only score M2 if M1 is correct.

6(d)	Any **three** from: • (Identity of the) metal • Charge (on the metal) / oxidation state / charge on complex • (Identity of the) ligands • Co-ordination number / number of ligands • Shape	3 max	

Question	Marking Guidance	Mark	Comments
7(a)	Iron(II): green (solution) gives a green precipitate	1	Apply list principle throughout if extra colours and/or extra observations given. Ignore state symbols in equations. Not blue-green ppt.
	$\underline{[Fe(H_2O)_6]^{2+}} + CO_3^{2-} \rightarrow FeCO_3 + 6H_2O$	1	Must start from $[Fe(H_2O)_6]^{2+}$ Allow equations with Na_2CO_3
	Iron(III):: yellow / purple / brown / lilac / violet (solution) gives a brown / rusty precipitate	1	
	Effervescence / gas / bubbles	1	Allow CO_2 evolved but not just CO_2
	$\underline{2[Fe(H_2O)_6]^{3+}} + 3CO_3^{2-} \rightarrow 2[Fe(H_2O)_3(OH)_3] + 3CO_2 + 3H_2O$	1	
7(b)	Copper(II): blue (solution) gives a green / yellow solution ***OR*** blue solution (turns) to green / yellow / olive green	1	Apply list principle throughout if extra colours and/or extra observations given. Ignore state symbols in equations.
	$\underline{[Cu(H_2O)_6]^{2+}} + 4Cl^- \rightarrow [CuCl_4]^{2-} + 6H_2O$	1	Allow equations with HCl
	Cobalt(II): pink (solution) gives a blue solution ***OR*** pink solution turns blue	1	
	$\underline{[Co(H_2O)_6]^{2+}} + 4Cl^- \rightarrow [CoCl_4]^{2-} + 6H_2O$	1	

7(c)	Iron(II): green (solution) gives a green precipitate	1	Apply list principle throughout if extra colours and/or extra observations given. Ignore state symbols in equations.
	$\underline{[Fe(H_2O)_6]^{2+}} + 2OH^- \rightarrow Fe(H_2O)_4(OH)_2 + 2H_2O$	1	Allow equations with NaOH
	Chromium(III): green / ruby / purple / violet / red-violet (solution) gives a green solution ***OR*** green / ruby / purple / violet / red-violet solution turns green	1	Ignore green ppt.
	$\underline{[Cr(H_2O)_6]^{3+}} + 6OH^- \rightarrow [Cr(OH)_6]^{3-} + 6H_2O$	1	Allow also with 4 or 5 OH balanced with 2 or 1 waters. Also allow two correct equations showing $Cr(H_2O)_3(OH)_3$ as intermediate.

7(d)	Al: colourless (solution) gives a white ppt	1	Apply list principle throughout if extra colours and/or extra observations given. Ignore state symbols in equations.
	$\underline{[Al(H_2O)_6]^{3+}} + 3NH_3 \rightarrow Al(H_2O)_3(OH)_3 + 3NH_4^+$	1	Allow $+ 3OH^- \rightarrow 3H_2O$ if $NH_3 + H_2O \rightarrow NH_4^+ + OH^-$ also
	Ag: colourless (solution) remains a colourless solution / no visible change	1	Ignore brown ppt.
	$\underline{[Ag(H_2O)_2]^+} + 2NH_3 \rightarrow [Ag(NH_3)_2]^+ + 2H_2O$	1	Allow 2 / 3 equations involving Ag_2O or $Ag(OH)_2$

Question	Marking Guidance	Mark	Comments
8(a)	Cobalt has variable oxidation states	1	Allow exists as Co(II) and Co(III)
	(It can act as an intermediate that) lowers the activation energy	1	Allow (alternative route with) lower E_a
	$CH_3CHO + 2Co^{3+} + H_2O \rightarrow CH_3COOH + 2Co^{2+} + 2H^+$	1	Allow multiples; allow molecular formulae Allow equations with H_3O^+
	$\frac{1}{2}O_2 + 2Co^{2+} + 2H^+ \rightarrow 2Co^{3+} + H_2O$	1	
8(b)(i)	$[Co(H_2O)_6]^{2+} + 3H_2NCH_2CH_2NH_2 \rightarrow [Co(H_2NCH_2CH_2NH_2)_3]^{2+} + 6H_2O$	1	Do not allow en in equation, allow $C_2H_8N_2$
	The number of particles increases / changes from 4 to 7	1	Can score M2 and M3 even if equation incorrect or missing provided number of particles increases.
	So the entropy change is positive / disorder increases / entropy increases	1	
8(b)(ii)	Minimum for **M1** is 3 bidentate ligands bonded to Co Ligands need not have any atoms shown but diagram must show 6 bonds from ligands to Co, 2 from each ligand	1	Ignore all charges for M1 and M3 but penalise charges on any ligand in M2
	Minimum for **M2** is one ligand identified as H_2N-----NH_2	1	Allow linkage as -C-C- or just a line.
	Minimum for **M3** is one bidentate ligand showing two arrows from separate nitrogens to cobalt	1	

8(c)	Moles of cobalt = (50 × 0.203)/1000 = <u>0.01015</u> mol	1	Allow 0.0101 to 0.0102
	Moles of AgCl = 4.22/143.4 = 0.0294	1	Allow 0.029 If not AgCl (eg $AgCl_2$ or $AgNO_3$), lose this mark and can only score **M1**, **M4** and **M5**
	Ratio = Cl^- to Co = 2.9 : 1	1	Do not allow 3 : 1 if this is the only answer but if 2.9:1 seen somewhere in answer credit this as **M3**
	$[Co(NH_3)_6]Cl_3$ (square brackets not essential)	1	
	Difference due to incomplete oxidation in the preparation	1	Allow incomplete reaction. Allow formation $[Co(NH_3)_5Cl]Cl_2$ etc. Some chloride ions act as ligands / replace NH_3 in complex. Do not allow 'impure sample' or reference to practical deficiencies.