OCR
RECOGNISING ACHIEVEMENT

GCE

Chemistry A

Advanced GCE **F325**

Equilibria, Energetics and Elements

Mark Scheme for June 2010

Oxford Cambridge and RSA Examinations

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2010

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

Question			Expected Answers	Marks	Additional Guidance
1	a		**F** **B** **G** **E** **D** **FIVE** correct ✓✓✓ **FOUR** correct ✓✓ **THREE** correct ✓	3	**ALLOW** **1450** **736** **G** **76** **–642**
	b		Correct calculation –642 – (+76 + (2 × 150) + 736 + 1450 + (2 × –349)) ✓ –642 – 1864 = – 2506 ✓ (kJ mol^{-1})	2	**ALLOW** for 1 mark: –2705 (2 × 150 and 2 × 349 not used for Cl) –2356 (2 × 150 not used for Cl) –2855 (2 × 349 not used for Cl) +2506 (wrong sign) **DO NOT ALLOW** any other answers
	c		Magnesium ion **OR** Mg^{2+} has greater charge (than sodium ion **OR** Na^{+}) **OR** Mg^{2+} has greater charge density ✓ Magnesium ion **OR** Mg^{2+} is smaller ✓ Mg^{2+} has a stronger attraction (than Na^{+}) to Cl^{-} ion **OR** Greater attraction between oppositely charged ions ✓	3	***ANNOTATIONS MUST BE USED*** **ALLOW** magnesium/Mg is 2+ but sodium/Na is 1+ **DO NOT ALLOW** Mg **atom** is 2+ but Na **atom** is 1+ **ALLOW** 'charge density' here **only** **ALLOW** Mg **OR** magnesium is smaller **DO NOT ALLOW** Mg^{2+} has a smaller **atomic** radius **ALLOW** anion **OR** negative ion for Cl^{-} **DO NOT ALLOW** chlorine ions **DO NOT ALLOW** Mg has greater attraction **ALLOW** 'attracts with more force' for greater attraction but **DO NOT ALLOW** 'greater force (could be repulsion) **ALLOW** reverse argument throughout in terms of Na^{+}
			Total	**8**	

Question			Expected Answers	Marks	Additional Guidance
2	a		$BrO_3^- + 5Br^- + 6H^+ \longrightarrow 3Br_2 + 3H_2O$ ✓	1	**ALLOW** multiples
	b			1	***ANNOTATIONS MUST BE USED***
			graph:		**Both** explanation and 1st order required for mark
			Straight/diagonal line through origin **OR** 0,0 **AND** 1st order with respect to BrO_3^- ✓		**DO NOT ALLOW** diagonal line **OR** straight line **OR** constant gradient on its own (no mention of origin **OR** 0,0) **ALLOW** 'As BrO_3^- doubles, rate doubles' **AND** 1st order **ALLOW** rate is proportional to concentration **AND** 1st order
			initial rates data:	4	
			When $[Br^-]$ is doubled, rate × 2 ✓ 1st order with respect to Br^- ✓		Mark order and explanation independently Mark order first, then explanation
			When $[H^+]$ × 2, rate × 4 (2^2) ✓ 2nd order with respect to H^+ ✓		
			Rate equation rate = k $[BrO_3^-]$ $[Br^-]$ $[H^+]^2$ ✓	1	**ALLOW** ECF from candidate's orders above

Question	Expected Answers	Marks	Additional Guidance
	Calculation of rate constant (3 marks) $k = \frac{\text{rate}}{[BrO_3^-][Br^-][H^+]^2}$ **OR** $\frac{1.19 \times 10^{-5}}{(5.0 \times 10^{-2})(1.5 \times 10^{-1})(3.1 \times 10^{-1})^2}$ ✓ $= 1.7 \times 10^{-2}$ **OR** 1.65×10^{-2} ✓ $dm^9\ mol^{-3}\ s^{-1}$ ✓	3	***ANNOTATIONS MUST BE USED*** **Calculation can be from any of the experimental runs – they all give the same value of *k*** **ALLOW** $mol^{-3}\ dm^9\ s^{-1}$ **ALLOW** 1.6510579×10^{-2} and correct rounding to 1.7×10^{-2} **Correct numerical answer subsumes previous marking point** **DO NOT ALLOW** fraction: $\frac{238}{14415}$ --- **ALLOW ECF from incorrect rate equation.** Examples are given below for 1st line of initial rates data. IF other rows have been used, then calculate the rate constant from data chosen. --- **Example 1:** 1st order with respect to H^+ *rate* = $k\ [BrO_3^-]\ [Br^-]\ [H^+]$ $k = \frac{\text{rate}}{[BrO_3^-][Br^-][H^+]}$ **OR** $\frac{1.19 \times 10^{-5}}{(5.0 \times 10^{-2})(1.5 \times 10^{-1})(3.1 \times 10^{-1})}$ ✓ $= 5.1 \times 10^{-3}$ **OR** 5.12×10^{-3} ✓ $dm^6\ mol^{-2}\ s^{-1}$ ✓ **ALLOW** $5.11827957 \times 10^{-3}$ and correct rounding to 5.1×10^{-3} --- **Example 2:** Zero order with respect to BrO_3^- *rate* = $k\ [Br^-]\ [H^+]^2$ $k = \frac{\text{rate}}{[Br^-][H^+]^2}$ **OR** $\frac{1.19 \times 10^{-5}}{(1.5 \times 10^{-1})(3.1 \times 10^{-1})^2}$ ✓ $= 8.3 \times 10^{-4}$ **OR** 8.26×10^{-4} ✓ $dm^6\ mol^{-2}\ s^{-1}$ ✓ **ALLOW** $8.255289629 \times 10^{-4}$ and correct rounding to 8.3×10^{-4}
	Total	**10**	

Question			Expected Answers	Marks	Additional Guidance
3	a		measured pH > 1 **OR** $[H^+] < 0.1$ (mol dm^{-3}) ✓	4	**ALLOW** C_2H_5 throughout question **ALLOW** $[H^+] < [CH_3CH_2COOH]$ **OR** $[H^+] < [HA]$ **ALLOW** measured pH is higher than expected **ALLOW** measured pH is not as acidic as expected **ALLOW** a quoted pH value or range > 1 and < 7 **OR** between 1 and 7
			$[H^+] = 10^{-pH}$ ✓		**ALLOW** $[H^+]$ = antilog –pH **OR** $[H^+]$ = inverse log –pH
			$K_a = \frac{[H^+][CH_3CH_2COO^-]}{[CH_3CH_2COOH]}$ **OR** $\frac{[H^+]^2}{[CH_3CH_2COOH]}$ ✓		**ALLOW** $\frac{[H^+][A^-]}{[HA]}$ **OR** $\frac{[H^+]^2}{[HA]}$
			Calculate K_a from $\frac{[H^+]^2}{0.100}$ ✓		**IF** K_a is **NOT** given and $K_a = \frac{[H^+]^2}{0.100}$ is shown, award mark for K_a also (i.e. $K_a = \frac{[H^+]^2}{0.100}$ is automatically awarded the last 2 marks)
	b		**Marks are for correctly calculated values.** **Working shows how values have been derived.** $[H^+] = 10^{-13.46} = 3.47 \times 10^{-14}$ (mol dm^{-3}) ✓	2	**ALLOW** $3.467368505 \times 10^{-14}$ and correct rounding to 3.5×10^{-14} **ALLOW** 0.28840315 and correct rounding to 0.29, i.e. **ALLOW** 0.288
			$[OH^-] = \frac{1.0 \times 10^{-14}}{3.47 \times 10^{-14}} = 0.29$ (mol dm^{-3}) ✓		**ALLOW** alternative approach using pOH: pOH = 14 – 13.46 = 0.54 ✓ $[OH^-] = 10^{-0.54} = 0.29$ (mol dm^{-3}) ✓ Correct answer gets **BOTH** marks

Question	Expected Answers	Marks	Additional Guidance
c		7	***ANNOTATIONS MUST BE USED***
	Propanoic acid reacts with sodium hydroxide forming propanoate ions/sodium propanoate **OR** $CH_3CH_2COOH + NaOH \rightarrow CH_3CH_2COONa + H_2O$ ✓		**ALLOW** C_2H_5 throughout question **ALLOW** Adding NaOH forms propanoate ions/sodium propanoate (imples that the NaOH is added to the propanoic acid)
	Some propanoic acid remains **OR** propanoic acid **AND** propanoate (ions) / sodium propanoate present ✓		**ALLOW**: weak acid **AND** its conjugate base/salt present Throughout, do not penalise comments that imply that pH is constant in presence of buffer
	equilibrium: $CH_3CH_2COOH \rightleftharpoons H^+ + CH_3CH_2COO^-$ ✓		**DO NOT ALLOW** HA and A^- in this equilibrium expression For description of action of buffer below, **ALLOW** HA for CH_3CH_2COOH; **ALLOW** A^- for $CH_3CH_2COO^-$ Equilibrium responses must refer back to a written equilibrium. **IF** no equilibrium shown, use the equilibrium as written in expected answers (which is also written on page 6 of the paper)
	Added alkali CH_3CH_2COOH reacts with added alkali **OR** $CH_3CH_2COOH + OH^- \rightarrow$ **OR** added alkali reacts with H^+ **OR** $H^+ + OH^- \rightarrow$ ✓		**ALLOW** weak acid reacts with added alkali
	$\rightarrow CH_3CH_2COO^-$ **OR** Equilibrium $\rightarrow$ right ✓		
	Added acid $CH_3CH_2COO^-$ reacts with added acid **OR** $[H^+]$ increases ✓		**ALLOW** conjugate base reacts with added acid **DO NOT ALLOW** salt reacts with added acid
	$\rightarrow CH_3CH_2COOH$ **OR** Equilibrium $\rightarrow$ left ✓		
		5	

Question			Expected Answers	Marks	Additional Guidance
	d		$HNO_3 + CH_3CH_2COOH \rightleftharpoons CH_3CH_2COOH_2^+ + NO_3^-$ ✓ acid 1 base 2 acid 2 base 1 ✓	2	State symbols **NOT** required **ALLOW** 1 **AND** 2 labels the other way around. **ALLOW** 'just acid' and 'base' labels throughout if linked by lines so that it is clear what the acid–base pairs are. **IF** proton transfer is wrong way around then **ALLOW** 2nd mark for idea of acid–base pairs, i.e. $HNO_3 + CH_3CH_2COOH \rightleftharpoons CH_3CH_2COO^- + H_2NO_3^+$ × base 2 acid 1 base 1 acid 2 ✓
	e	**i**	$2CH_3CH_2COOH + Mg \rightarrow (CH_3CH_2COO)_2Mg + H_2$ ✓	1	**IGNORE** state symbols **ALLOW** ionic equation: $2H^+ + Mg \rightarrow Mg^{2+} + H_2$ **IGNORE** any random charges in formula of $(CH_3CH_2COO)_2Mg$ as long as the charges are **correct (charges are treated as working)** i.e. $(CH_3COO^-)_2Mg$ **OR** $(CH_3COO)_2^-Mg$ should **not** be penalised However, Mg^{2+} instead of Mg on the left side of equation is obviously wrong
		ii	$2H^+ + CO_3^{2-} \longrightarrow H_2O + CO_2$ **OR** $2H^+ + CO_3^{2-} \longrightarrow H_2CO_3$ **OR** $H^+ + CO_3^{2-} \longrightarrow HCO_3^-$ ✓	1	State symbols **NOT** required
			Total	**17**	

Question			Expected Answers	Marks	Additional Guidance
4	a	i	Complete circuit (with voltmeter) and salt bridge linking two half-cells ✓ Pt electrode in solution of Fe^{2+}/Fe^{3+} ✓ Ag in solution of Ag^{+} ✓	3	**DO NOT ALLOW** 'solution of a silver halide', e.g. AgCl (as these are insoluble) but **DO ALLOW** any solution of any other silver salt (whether insoluble or not) **IF** candidate has used incorrect redox systems, then mark ECF as follows: **(i) each** incorrect system will cost the candidate **one** mark **(ii) ECF** if species have been quoted (see Additional Guidance below) **(iii) ECF** for equation **(iv) ECF** for cell potential ***YOU MAY NEED TO WORK OUT THESE ECF RESPONSES YOURSELF DEPENDING ON THE INCORRECT REDOX SYSTEMS CHOSEN***
		ii	electrons **AND** ions ✓	1	For electrons, **ALLOW** e^{-} For 'ions', **ALLOW** formula of an ion in one of the half-cells or salt bridge, e.g. Ag^{+}, Fe^{2+}, Fe^{3+} **ALLOW ECF** as in **(i)**
		iii	$Ag + Fe^{3+} \longrightarrow Ag^{+} + Fe^{2+}$ ✓	1	**ALLOW ECF** as in **(i)** **ALLOW** equilibrium sign
		iv	0.43 V ✓	1	**ALLOW ECF** as in **(i)**
	b	i	Cl_2 **OR** O_2 **AND** H^{+} ✓	1	**ALLOW** chlor**ine** **ALLOW** O_2 **AND** $4H^{+}$ **ALLOW** O_2 **AND** acid **DO NOT ALLOW** O_2 alone **DO NOT ALLOW** equation or equilibrium
		ii	I^{-} ✓	1	**ALLOW** $2I^{-}$ **OR** iod**ide** **DO NOT ALLOW** equation or equilibrium

Question			Expected Answers	Marks	Additional Guidance
	c		A fuel cell converts energy from reaction of a fuel with oxygen into a voltage/electrical energy ✓ $2H_2 + O_2 \rightarrow 2H_2O$ ✓ Two from: • under pressure **OR** at low temperature **OR** as a liquid • adsorbed on solid • absorbed within solid ✓✓ Energy is needed to make the hydrogen **OR** energy is needed to make fuel cell ✓	5	***ANNOTATIONS MUST BE USED*** **ALLOW** combustion for reaction of fuel with oxygen/reactants **ALLOW** a fuel cell requires constant supply of fuel **OR** operates continuously as long as a fuel (and oxygen) are added **ALLOW** multiples, e.g. $H_2 + ½O_2 \rightarrow H_2O$ **IGNORE** state symbols **ALLOW** 'material' **OR** metal for solid **ALLOW** as a metal hydride
			Total	**13**	

Question			Expected Answers	Marks	Additional Guidance
5	a	i	$(K_c =) \frac{[NH_3]^2}{[N_2]\,[H_2]^3}$ ✓	1	Must be square brackets
		ii	$dm^6\ mol^{-2}$ ✓	1	**ALLOW** $mol^{-2}\ dm^6$ **ALLOW ECF** from incorrect K_c expression
	b		**Unless otherwise stated, marks are for correctly calculated values. Working shows how values have been derived.** $[N_2] = \frac{7.2}{6.0}$ **OR** 1.2 ($mol\ dm^{-3}$) **AND** $[H_2] = \frac{12}{6.0}$ **OR** 2.0 ($mol\ dm^{-3}$) ✓ $[NH_3] = \sqrt{(K_c \times [N_2] \times [H_2]^3)}$ **OR** $\sqrt{(8.00 \times 10^{-2} \times 1.2 \times 2.0^3)}$ ✓ = 0.876 **OR** 0.88 ($mol\ dm^{-3}$) ✓ amount NH_3 = 0.876 × 6 = 5.26 **OR** 5.3 (mol) ✓	4	***ANNOTATIONS MUST BE USED*** For **all** parts, **ALLOW** numerical answers from 2 significant figures up to the calculator value 1st mark is for realising that concentrations need to be calculated. **Correct numerical answer with no working would score all previous calculation marks** **ALLOW** calculator value: 0.876356092 down to 0.88, correctly rounded **ALLOW** calculator value down to 5.3, correctly rounded




Question	Expected Answers	Marks	Additional Guidance
b	**EXAMPLES OF INCORRECT RESPONSES IN (b) THAT MAY BE WORTHY OF CREDIT**		---------- **ALLOW ECF from incorrect concentrations (3 marks)** For example, If concentrations **not** calculated at start, then $[NH_3] = \sqrt{(8.00 \times 10^{-2} \times 7.2 \times 12.0^3)}$ ✓ = 31.5 mol dm^{-3} ✓ Equilibrium amount of $NH_3 = 31.5 \times 6 = 189.6$ (mol) ✓ ---------- **IF** candidate has K_c expression upside down, then all **4 marks** are available in **(b)** by **ECF** Correct $[N_2]$ AND $[H_2]$ ✓ $[NH_3] = \sqrt{\frac{[N_2][H_2]^3}{K_c}} = = \sqrt{\frac{1.2 \times 2^3}{8.00 \times 10^{-2}}}$ ✓ = 11.0 mol dm^{-3} ✓ Equilibrium amount of $NH_3 = 11.0 \times 6 = 66.0$ (mol) ✓ ---------- **IF** candidate has used K_c value of 8.00×10^{-2} **AND** values for N_2 **AND** H_2 with powers wrong, mark by **ECF** from calculated as below (**3 max in (b)**) Correct $[N_2]$ AND $[H_2]$ ✓ $[NH_3]$ expression × **ECF**: Calculated $[NH_3]$ ✓ **ECF**: Equilibrium amount of NH_3 ✓

Question			Expected Answers	Marks	Additional Guidance
	c	i	Equilibrium shifts to right **OR** Equilibrium towards ammonia ✓ Right hand side has fewer number of (gaseous) moles ✓	2	**ALLOW** 'moves right' **OR** 'goes right' **OR** 'favours right' **OR** 'goes forwards' **ALLOW** 'ammonia side' has fewer moles **ALLOW** 'there are more (gaseous) moles on left'
		ii	K_c does not change ✓ Increased pressure increases concentration terms on bottom of K_c expression more than the top **OR** system is now no longer in equilibrium ✓ top of K_c expression increases and bottom decreases until K_c is reached ✓	3	***ANNOTATIONS MUST BE USED*** Any response in terms of K_c changing scores **ZERO** for Part **(ii)** **ALLOW** K_c is temperature dependent only **OR** K_c does not change with pressure **ALLOW** $\frac{[NH_3]^2}{[N_2]\ [H_2]^3}$ no longer equal to K_c
	d	i	$CH_4 + H_2O \longrightarrow 3H_2 + CO$ ✓	1	State symbols **NOT** required **ALLOW**: $CH_4 + H_2O \longrightarrow CH_3OH + H_2$ $CH_4 + 2H_2O \longrightarrow 4H_2 + CO_2$ $CH_4 + H_2O \longrightarrow 2H_2 + HCHO$ $CH_4 + 2H_2O \longrightarrow 3H_2 + HCOOH$
		ii	Electrolysis of water **OR** $H_2O \longrightarrow H_2 + ½O_2$ ✓	1	**ALLOW** electrolysis of brine **DO NOT ALLOW** reforming **DO NOT ALLOW** cracking **DO NOT ALLOW** reaction of metal with acid

Question			Expected Answers	Marks	Additional Guidance
	e	i	**Unless otherwise stated, marks are for correctly calculated values.** **Working shows how values have been derived.** $\Delta S = \Sigma S$(products) – ΣS(reactants) / = (2 × 192) – (191 + 3 × 131) ✓ = –200 (J K^{-1} mol^{-1}) **OR** –0.200 (kJ K^{-1} mol^{-1}) ✓ Use of 298 K (could be within ΔG expression below) ✓ $\Delta G = \Delta H - T\Delta S$ **OR** ΔG = –92 – (298 × –0.200) **OR** ΔG = –92000 – (298 × –200) ✓ = –32.4 kJ mol^{-1} **OR** –32400 J mol^{-1} ✓ **(Units must be shown)** For feasibility, $\Delta G < 0$ **OR** ΔG is negative ✓	5 1	***ANNOTATIONS MUST BE USED*** See **Appendix 1** for extra guidance for marking **5e(i)** and **5e(ii)** **NO UNITS** required at this stage **IGNORE** units **ALLOW** –32.4 kJ **OR** –32400 J **(Units must be shown)** Award all 5 marks **above** for correct answer with no working **IF** 25 ºC has been used instead of 298 K, correctly calculated ΔG values are = –87 kJ mol^{-1} **OR** –87000 J mol^{-1} **4 marks** are still available up to this point and maximum possible from **(e)(i)** is 5 marks
		ii	As the temperature increases, $T\Delta S$ becomes more negative **OR** $T\Delta S$ becomes more negative than ΔH **OR** $T\Delta S$ becomes more significant ✓ Eventually $\Delta H - T\Delta S$ becomes positive ✓	2	**ALLOW** $T\Delta S > \Delta H$ (i.e. assume no sign at this stage) **ALLOW** 'entropy term' as alternative for $T\Delta S$ **ALLOW –** $T\Delta S$ becomes more positive **ALLOW –** $T\Delta S$ decreases **ALLOW** ΔG becomes positive **OR** $\Delta G > 0$

Question			Expected Answers	Marks	Additional Guidance
		iii	Activation energy is too high **OR** reaction too slow ✓	1	**ALLOW** increases the rate **OR** more molecules exceed activation energy **OR** more successful collisions **ALLOW** rate constant increases **IGNORE** comments on yield
			Total	**22**	

Question			Expected Answers	Marks	Additional Guidance
6	a	i	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^1$ ✓	1	**ALLOW** $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1 3d^5$ (i.e. 4s before 3d) **ALLOW** $[Ar]4s^1 3d^5$ **OR** $[Ar]3d^5 4s^1$
		ii	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^3$ ✓	1	**ALLOW** $[Ar]3d^3$ **ALLOW** $1s^2 2s^2 2p^6 3s^2 3p^6 3d^3 4s^0$ **OR** $[Ar]3d^3 4s^0$
	b		$Zn \longrightarrow Zn^{2+} + 2e^-$ ✓ $Cr_2O_7^{2-} + 14H^+ + 8e^- \longrightarrow 2Cr^{2+} + 7H_2O$ ✓ $4Zn + Cr_2O_7^{2-} + 14H^+ \longrightarrow 4Zn^{2+} + 2Cr^{2+} + 7H_2O$ ✓	3	**ALLOW** multiples **WATCH** for balancing of the equations printed on paper **IF** printed equations and answer lines have different balancing numbers **OR** electrons, **IGNORE** numbers on printed equations (i.e. treat these as working) and mark responses on answer lines **only** **NO ECF** for overall equation i.e. the expected answer is the **ONLY** acceptable answer
	c	i	Ligand substitution ✓	1	**ALLOW** ligand exchange
		ii	$[Cr(H_2O)_6]^{3+} + 6NH_3 \longrightarrow [Cr(NH_3)_6]^{3+} + 6H_2O$ ✓ ✓	2	1 mark is awarded for each side of equation **ALLOW** equilibrium sign **ALLOW** 1 mark for 2+ shown instead of 3+ on both sides of equation **ALLOW** 1 mark for substitution of 4 NH_3: $[Cr(H_2O)_6]^{3+} + 4NH_3 \longrightarrow [Cr(NH_3)_4(H_2O)_2]^{3+} + 4H_2O$
	d	i	Donates an electron pair to a metal ion **OR** forms a coordinate bond to a metal ion ✓	1	**ALLOW** donates an electron pair to a metal **ALLOW** dative (covalent) bond for coordinate bond
		ii	Donates **two** electron pairs **OR** forms **two** coordinate bonds ✓ Lone pairs on two O atoms ✓	2	First mark is for the idea of two coordinate bonds **ALLOW** lone pair on O and N **DO NOT ALLOW** lone pairs on COO^- (could involve C) Second mark is for the atoms that donate the electron pairs Look for the atoms with lone pairs also on response to **(d)(iii)** and credit here if not described in **(d)(ii)**

Question	Expected Answers	Marks	Additional Guidance
iii	Forms two optical isomers **OR** two enantiomers **OR** two non-superimposable mirror images ✓  ✓✓ For each structure	3	**IGNORE** any charges shown **ALLOW** any attempt to show bidentate ligand. Bottom line is the diagram on the left. 1 mark for 3D diagram with ligands attached for ONE stereoisomer. Must contain 2 out wedges, 2 in wedges and 2 lines in plane of paper:  2nd mark for reflected diagram of SECOND stereoisomer. The diagram below would score the 2nd mark but not the first 

Question			Expected Answers	Marks	Additional Guidance
	e		N : H : Cr : O 11.1/14 : 3.17/1 : 41.27/52 : 44.45/16 **OR** 0.793 : 3.17 : 0.794 : 2.78 ✓	8	***ANNOTATIONS MUST BE USED***
			A: $N_2H_8Cr_2O_7$ ✓		**ALLOW A**: $(NH_4)_2Cr_2O_7$
			Ions: NH_4^+ ✓ $Cr_2O_7^{2-}$ ✓		**IF** candidate has obtained NH_4CrO_4 for A, **ALLOW** NH_4^+ **DO NOT ALLOW** CrO_4^-
			B: Cr_2O_3 ✓		
			Correctly calculates molar mass of **C** = 1.17 × 24.0 = 28.08 (g mol^{-1}) ✓		**ALLOW**: (relative) molecular mass **ALLOW**: 28 **ALLOW**: '**C** is 28'
			C: N_2 ✓		
			Equation: $(NH_4)_2Cr_2O_7 \longrightarrow Cr_2O_3 + 4H_2O + N_2$ ✓		**ALLOW** $N_2H_8Cr_2O_7$ in equation.
			Total	**22**	

Question			Expected Answers	Marks	Additional Guidance
7	a	i	$H_2O_2 \longrightarrow O_2 + 2H^+ + 2e^-$ ✓✓	2	All other multiples score 1 mark e.g. ½ $H_2O_2 \longrightarrow$ ½ $O_2 + H^+ + e^-$ $5H_2O_2 \longrightarrow 5O_2 + 10H^+ + 10e^-$
	b		**Marks are for correctly calculated values.** **Working shows how values have been derived.**		***ANNOTATIONS MUST BE USED***
			$n(KMnO_4) = \frac{0.0200 \times 23.45}{1000} = 4.69 \times 10^{-4}$ (mol) ✓		**DO NOT ALLOW** 4.7×10^{-4}
			$n(H_2O_2) = 5/2 \times 4.69 \times 10^{-4} = 1.1725 \times 10^{-3}$ (mol) ✓		**ALLOW** 1.173×10^{-3} OR 1.17×10^{-3} (i.e. 3 significant figures upwards) **ALLOW** by **ECF**: 5/2 × ans above
			$n(H_2O_2)$ in 250 cm^3 solution $= 10 \times 1.1725 \times 10^{-3} = 1.1725 \times 10^{-2}$ (mol) ✓		**ALLOW** by **ECF** 10 × ans above **ALLOW** concentration H_2O_2 = 0.0469 mol dm^{-3}
			concentration in g dm^{-3} of original H_2O_2 $= 40 \times 1.1725 \times 10^{-2} \times 34 = 15.9$ (g dm^{-3}) ✓	4	**ALLOW** by **ECF** $40 \times n(H_2O_2) \times 34$ **ALLOW** 0.0469 x 10 x 34 = 15.9 g dm^{-3} ✓ **ALLOW** two significant figures, 16 (g dm^{-3}) up to calculator value of 15.946 g dm^{-3}
			$n(O_2) = 5/2 \times 4.69 \times 10^{-4} = 1.1725 \times 10^{-3}$ (mol) ✓		**ALLOW** 0.028 dm^3 **OR** 0.02814 dm^3 **ALLOW** 28 cm^3 **OR** 28.14 cm^3 Value **AND** units required **DO NOT ALLOW** 0.03 dm^3
			volume $O_2 = 24.0 \times 1.1725 \times 10^{-3} = 0.0281$ dm^3 ✓	2	**ALLOW** by **ECF**: 24.0 × calculated moles of O_2 (2 significant figures up to calculator value)
			Total	**8**	

Appendix 1

Extra guidance for marking atypical responses to **5e(i)** and **5e(ii)**

Question			Expected Answer	Mark	Additional Guidance
5	e	i	**TOTAL ENTROPY APPROACH:** **ALL MARKS AVAILABLE** **Unless otherwise stated, marks are for correctly calculated values.** **Working shows how values have been derived.**		***ANNOTATIONS MUST BE USED***
			$\Delta S = \Sigma S$(products) – ΣS(reactants) / = (2 × 192) – (191 + 3 × 131) ✓ = –200 (J K^{-1} mol^{-1}) **OR** –0.200 (kJ K^{-1} mol^{-1}) ✓		**NO UNITS** required at this stage **IGNORE** units
			Use of 298 K (could be within expression below) ✓		
			$\Delta S_{total} = \Delta S_{system} + \Delta S_{surroundings}$ $\Delta S_{surroundings} = -\frac{\Delta H}{T}$ **OR** $\Delta S_{total} = \Delta S_{system} - \frac{\Delta H}{T}$ **OR** $\Delta S_{total} = -0.200 - \frac{-92}{298}$ **OR** $\Delta S_{total} = -200 - \frac{-92000}{298}$ ✓ = 0.109 kJ (K^{-1} mol^{-1}) **OR** 109 J (K^{-1} mol^{-1}) ✓	5	**ALLOW** 0.109 kJ **OR** 109 J **IF** 25ºC has been used instead of 298 K, correctly calculated ΔS_{total} values are = 3.48 kJ K^{-1} mol^{-1} **OR** 3,480 J K^{-1} mol^{-1}
			Feasible when ΔS_{total} >0 ✓	1	

Question			Expected Answer	Mark	Additional Guidance
5	e	i	**MAX/MIN TEMPERATURE APPROACH:** **5 MARKS MAX AVAILABLE** **Unless otherwise stated, marks are for correctly calculated values.** **Working shows how values have been derived.**		***ANNOTATIONS MUST BE USED*** **This candidate has not answered the question but many marks are still available.**
			$\Delta S = \Sigma S$(products) – ΣS(reactants) / = (2 × 192) – (191 + 3 × 131) ✓ = –200 (J K^{-1} mol^{-1}) **OR** –0.200 (kJ K^{-1} mol^{-1}) ✓		**NO UNITS** required at this stage **IGNORE** units
			Use of 298 K (could be within ΔG expression below) ✓		
			$\Delta G = \Delta H - T\Delta S$ **OR** When $\Delta G = 0$, $0 = \Delta H - T\Delta S$; **OR** $T = \frac{\Delta H}{\Delta S} = \frac{-92}{-0.200}$ **OR** $T = \frac{\Delta H}{\Delta S} = \frac{-92000}{-200}$ ✓ = 460 K ✓ = 187 ºC (use of 298) ✓		
			The condition $\Delta G = 0$ because temperature at which $\Delta G = 0$ is the maximum temperature for feasibility **AND** justification for the being the maximum ✓		By this approach, the calculated temperature is the switchover between feasibility and non-feasibility but it cannot be assumed that this is the maximum temperature

Question			Expected Answer	Mark	Additional Guidance
5	e	ii	As the temperature increases, $\Delta H/T$ becomes **less** negative **OR** $\Delta H/T$ becomes **more** negative than ΔS(system) **OR** $\Delta H/T$ becomes **less** significant **OR** ΔS(surroundings) becomes **less** significant **OR** ΔS(system) > $\Delta H/T$ **OR** ΔS(system) > ΔS(surroundings) ✓ Eventually ΔS(total) becomes **negative** ✓	2	**ALLOW** $\Delta H/T > \Delta S_{system}$ (i.e. assume no sign at this stage) **ALLOW –**$\Delta H/T$ becomes more positive **ALLOW –**$\Delta H/T$ increases

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)
Telephone: 01223 553998
Facsimile: 01223 552627
Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2010