OCR
RECOGNISING ACHIEVEMENT

Thursday 19 January 2012 – Afternoon

AS GCE CHEMISTRY A

F322 Chains, Energy and Resources

F3144401122

Candidates answer on the Question Paper.

OCR supplied materials:
- *Data Sheet for Chemistry A* (inserted)

Other materials required:
- Scientific calculator

Duration: 1 hour 45 minutes

F322

Candidate forename		Candidate surname	

Centre number						Candidate number				

INSTRUCTIONS TO CANDIDATES

- The Insert will be found in the centre of this document.
- Write your name, centre number and candidate number in the boxes above. Please write clearly and in capital letters.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Write your answer to each question in the space provided. If additional space is required, you should use the lined pages at the end of this booklet. The question number(s) must be clearly shown.
- Do **not** write in the bar codes.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets **[ ]** at the end of each question or part question.



Where you see this icon you will be awarded marks for the quality of written communication in your answer.

This means for example you should:
- ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear;
- organise information clearly and coherently, using specialist vocabulary when appropriate.


- You may use a scientific calculator.
- A copy of the *Data Sheet for Chemistry A* is provided as an insert with this question paper.
- You are advised to show all the steps in any calculations.
- The total number of marks for this paper is **100**.
- This document consists of **24** pages. Any blank pages are indicated.

© OCR 2012 [H/500/7834]
OCR is an exempt Charity

Answer **all** the questions.

1 Crude oil is a source of many hydrocarbons.

The skeletal formulae of some of these hydrocarbons are shown below.



(a) Explain why compound **A** is both *saturated* and a *hydrocarbon*.

.....

.....

.....

..... **[2]**

(b) What is the empirical formula for compound **A**?

..... **[1]**

(c) Give the letters, **A**, **B**, **C**, **D**, **E**, **F**, **G**, **H** or **I**, of two hydrocarbons that are structural isomers of each other.

..... and ..... **[1]**

© OCR 2012

(d) The petroleum industry processes straight chain alkanes into cyclic hydrocarbons such as **A**, **B** and **E**.

(i) Explain why the petroleum industry processes straight chain alkanes into cyclic hydrocarbons.

.....

..... **[1]**

(ii) Hydrocarbon **C** can be processed into the cyclic hydrocarbon **B**.

Construct an equation for this reaction.

[1]

(e) Explain why hydrocarbon **D** has a higher boiling point than hydrocarbon **C**.

.....

.....

.....

..... **[2]**

(f) Hydrocarbons **G** and **H** are stereoisomers of each other.

Explain what is meant by the term *stereoisomerism*.

.....

.....

.....

..... **[2]**

(g) Construct the equation for the **complete** combustion of hydrocarbon **C**.

..... **[2]**


© OCR 2012

(h) A hydrocarbon molecule, $C_{16}H_{34}$, is cracked to form an octane molecule and two molecules of but-2-ene.

Construct the equation for this reaction.

..... **[1]**

(i) Compound **I** is 3-methylheptane. It does not contain a functional group.



I

(i) What is meant by the term *functional group*?

.....

.....

..... **[1]**

(ii) Compound **I** reacts with chlorine in the presence of ultraviolet radiation to give several structural isomers of $C_8H_{17}Cl$.

How many **structural** isomers could be formed in this reaction?

..... **[1]**

(iii) The mechanism of the reaction involves radicals.

What is meant by the term *radical*?

.....

..... **[1]**

[Total: 16]

© OCR 2012

2 Ethanoic acid, CH_3COOH, is used to make esters.

Some information about two of the processes used to make ethanoic acid is given below.

Process 1

This is a one-step process that involves the reaction of methanol with carbon monoxide.

$$CH_3OH + CO \rightarrow CH_3COOH$$

The conditions used are 180 °C and 30 atmospheres pressure. A rhodium/iodine catalyst is used.

The percentage yield for this process is 99%.

Process 2

This involves the oxidation of naphtha, a fraction obtained from crude oil.

Liquid naphtha is oxidised using air at a temperature of 180 °C and 50 atmospheres pressure. No catalyst is needed.

A large variety of other products are also formed in this oxidation.

(a) Suggest **three** advantages of making ethanoic acid using **Process 1** rather than **Process 2**.

.....

.....

.....

.....

.....

.....

..... **[3]**


© OCR 2012

(b) The other products formed in **Process 2** are carboxylic acids, aldehydes and ketones.

A research chemist investigates some of these other products of **Process 2**.

(i) The research chemist isolates product, **J**.

The infrared spectrum of **J** is shown below.



The chemist also finds that 0.172 g of a pure sample of **J** contains 2.00×10^{-3} mol of **J**.

Suggest, with reasons, **one** possible structure for **J**.

In your answer you should link the evidence with your explanation.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... **[5]**

© OCR 2012

(ii) The chemist isolates another product, the carboxylic acid, **K**.

K has the molecular formula $C_4H_8O_2$.

Suggest a possible structure and name for **K**.

structure

name ..... **[2]**

(c) Ethanoic acid is used in the manufacture of the ester, propyl ethanoate.

Describe how ethanoic acid is converted into propyl ethanoate.
Include an equation in your answer.

.....

.....

.....

.....

.....

..... **[4]**

[Total: 14]

© OCR 2012

3 Enthalpy changes can be determined directly or indirectly.

(a) A student investigates the reaction between magnesium and dilute hydrochloric acid.

$$Mg(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$$

The student determines the enthalpy change for this reaction.

In her experiment, she reacts 0.486 g of magnesium with 50.0 cm^3 of 2.00 mol dm^{-3} HCl(aq). The HCl(aq) is in excess.

The temperature of the solution changes from 19.2 °C to 32.0 °C.

(i) Calculate the energy released, in kJ, during this reaction.

The specific heat capacity of the solution = 4.18 J g^{-1} K^{-1}.

The density of the solution is 1.00 g cm^{-3}.

energy = ...kJ **[2]**

(ii) Calculate the amount, in moles, of magnesium used by the student.

amount = ... mol **[1]**

(iii) Calculate the enthalpy change of reaction.

Give your answer to **three** significant figures.

enthalpy change of reaction = ...kJ mol^{-1} **[3]**

© OCR 2012

(b) The student wants to determine the enthalpy change of formation of calcium carbonate, $CaCO_3(s)$.

$$Ca(s) + C(s) + 1\tfrac{1}{2}O_2(g) \rightarrow CaCO_3(s)$$

(i) What is meant by the term *standard enthalpy change of formation*?
You should state the standard conditions in your answer.

.....

.....


.....

..... **[3]**

(ii) Using the following data and enthalpy cycle,

- fill in the boxes on the enthalpy cycle with the correct enthalpy change values
- calculate the enthalpy change of formation, ΔH_f, of $CaCO_3(s)$.

reaction	enthalpy change, ΔH / kJ mol^{-1}
$C(s) + O_2(g) \rightarrow CO_2(g)$	–393
$H_2(g) + \frac{1}{2}O_2(g) \rightarrow H_2O(l)$	–285
$CaCO_3(s) + 2HCl(aq) \rightarrow CaCl_2(aq) + H_2O(l) + CO_2(g)$	–54
$Ca(s) + 2HCl(aq) \rightarrow CaCl_2(aq) + H_2(g)$	–168



ΔH_f = ... kJ mol^{-1} **[3]**

[Total: 12]

© OCR 2012

4 Hydrogen and chlorine are reacted together to form hydrogen chloride.

$$H_2(g) + Cl_2(g) \rightarrow 2HCl(g) \qquad \Delta H = -184 \text{ kJ mol}^{-1}$$

(a) Calculate the bond enthalpy for the H–C*l* bond using the information in the table below.

bond	bond enthalpy / kJ mol^{-1}
H–H	+436
C*l*–C*l*	+243

bond enthalpy = ... kJ mol^{-1} **[2]**

(b) The reaction is repeated at a **higher** pressure.

Describe and explain what happens to the rate of the reaction between $H_2(g)$ and $Cl_2(g)$.

.....

.....

.....

.....

..... **[2]**

© OCR 2012

(c) The reaction is repeated again. This time the temperature is **decreased**.

Describe and explain, by drawing appropriately labelled Boltzmann distributions, what happens to the rate of reaction between $H_2(g)$ and $Cl_2(g)$.

.....

.....

.....

.....

..... **[5]**

(d) The mechanism of the reaction between $H_2(g)$ and $Cl_2(g)$ involves initiation, propagation and termination.

(i) The initiation step is the homolytic fission of the covalent bond in a chlorine molecule.

Write an equation to show this homolytic fission.

..... **[1]**

(ii) Complete the following equations which show the propagation steps.

$Cl + H_2 \rightarrow$ ..... + ...

$H + Cl_2 \rightarrow$ ..... + ... **[2]**

(iii) Suggest equations for **two** termination steps.

.....

..... **[2]**

[Total: 14]

© OCR 2012

5 Sulfuric acid is made from sulfur, oxygen and water in a three-stage process.
This can be represented by the following overall equation.

$$2S(s) + 3O_2(g) + 2H_2O(l) \rightarrow 2H_2SO_4(l)$$

(a) Explain why the overall process to make sulfuric acid has an atom economy of 100%.

.....

.....

..... **[1]**

(b) A factory uses 51.4 tonnes of sulfur to manufacture 147 tonnes of H_2SO_4.

What is the percentage yield of H_2SO_4?

Give your answer to **two** significant figures. (1 tonne = 1×10^6 g)

percentage yield = ... % **[3]**

© OCR 2012

(c) One of the reactions involved in making sulfuric acid converts sulfur dioxide, SO_2, into sulfur trioxide, SO_3.

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \quad \Delta H = -197 \text{ kJ mol}^{-1}$$

This reaction can be carried out at 450 °C and 3 atmospheres pressure in the presence of a V_2O_5 catalyst.

Under these conditions the position of equilibrium is almost completely on the right-hand side.

(i) A research chemist investigates this reaction.
He uses a temperature of 450 °C and 3 atmospheres pressure.
The research chemist does **not** use the catalyst.

Predict the changes, if any, on each of the following.

position of equilibrium .....

rate of backward reaction ..... **[2]**

(ii) The temperature of the reaction mixture is **increased** to 600 °C.

State and explain what will happen to the position of equilibrium.

.....

.....

.....

..... **[1]**

(iii) The pressure of the reaction mixture is **decreased** to 2 atmospheres.

State and explain what will happen to the position of equilibrium.

.....

.....

.....

..... **[1]**

© OCR 2012

(d) Concentrated H_2SO_4 is used as an acid catalyst in the elimination of water from alcohols.

There are several alcohols that are structural isomers with the formula $C_5H_{11}OH$. When these alcohols are heated with H_2SO_4 they form alkenes.

$$C_5H_{11}OH \rightarrow C_5H_{10} + H_2O$$

(i) Pentan-1-ol is a structural isomer of $C_5H_{11}OH$ that is a primary alcohol.

Draw the structure of another structural isomer of $C_5H_{11}OH$ that is a primary alcohol.

[1]

(ii) Pentan-2-ol is a structural isomer of $C_5H_{11}OH$ that is a secondary alcohol.
Pentan-2-ol is heated with H_2SO_4.

Three alkenes are formed, **L**, **M** and **N**.

- **L** and **M** are stereoisomers.
- **N** is a structural isomer of the stereoisomers **L** and **M**.

Draw the structures for alkenes **L**, **M** and **N**.

alkene **L**

alkene **M**

alkene **N**

[3]

© OCR 2012

(iii) One structural isomer of $C_5H_{11}OH$ is an alcohol that **cannot** be oxidised by heating with acidified potassium dichromate(VI).

Draw the structure of this alcohol.

[1]


[Total: 13]

© OCR 2012

6 Mass spectrometry is used in analysis.

(a) Compound **O** contains carbon, hydrogen and oxygen.

The mass spectrum of compound **O** is shown below.



(i) Identify the *m*/*z* value that corresponds to the molecular ion.

..... **[1]**

(ii) Write the formula of the ion that gives rise to the peak at *m*/*z* = 31.

..... **[1]**

(iii) Suggest the molecular formula for **O**.

..... **[1]**

© OCR 2012

(b) A scientist analyses a sample of Moon rock. She uses mass spectrometry to find out which metal the sample contains.

The mass spectrum of the sample shows *m*/*z* peaks as shown in the table.

***m*/*z* value**	**percentage abundance (%)**
63	72.2
65	27.8

Positive ions, $\mathbf{X}^+$, of the metal were responsible for the two *m*/*z* peaks.

Identify the metal **X** by calculating its relative atomic mass to **one decimal place**.

relative atomic mass of **X** = ...

metal **X** = ... **[3]**

[Total: 6]

© OCR 2012

7 Chlorofluorocarbons, CFCs, were once used as propellants in aerosols. CFCs contribute to ozone depletion in the upper atmosphere.

(a) A CFC has the formula CF_2Cl_2.

State the three-dimensional shape of a CF_2Cl_2 molecule and the F–C–Cl bond angle.

shape .....

bond angle ..... **[2]**

(b) Two reasons that CF_2Cl_2 was used as an aerosol propellant are that it has low reactivity and will not hydrolyse in water.

(i) State **one** other reason why CF_2Cl_2 was developed for use as an aerosol.

.....

..... **[1]**

(ii) Suggest why CF_2Cl_2 does **not** hydrolyse in water.

.....

.....

..... **[1]**

(c) Explain, with the aid of equations, how the presence of CFCs in the upper atmosphere leads to ozone depletion.

.....

.....

.....

.....

..... **[3]**

(d) Why are scientists concerned about ozone depletion?

.....

.....

..... **[1]**

© OCR 2012

(e) International agreements have reduced the use of CFCs. However the concentration of atmospheric CFCs has hardly changed.

Suggest **two** reasons why.

.....

.....

.....

.....

..... **[2]**


[Total: 10]

© OCR 2012

8 Cyclopentene is a cyclic alkene.

(a) The flowchart shows some reactions involving cyclopentene and cyclopentanol.

Complete the partial structures in the boxes to show compounds **P**, **Q** and **R**, the main organic products of the reactions.



[3]

(b) What would be the colour change in the reaction between cyclopentene and bromine?

..... to ..... **[1]**

© OCR 2012

(c) Cyclopentene can be polymerised to give poly(cyclopentene).


Draw a section of poly(cyclopentene) to show **two** repeat units.

[1]

(d) Cyclopentene reacts with HC*l* by electrophilic addition.

Use the curly arrow model to complete the mechanism for this reaction.

In your answer include any relevant dipoles, the intermediate and the product.



intermediate


product

[5]

TURN OVER FOR QUESTION 8(e)

© OCR 2012


(e) Chlorocyclopentane can be hydrolysed by heating with aqueous sodium hydroxide.



Use the curly arrow model to complete the mechanism for this hydrolysis reaction.

Include in your answer, relevant dipoles, the name of the mechanism and the type of bond fission.

*In your answer you should use the correct technical terms, spelled correctly***.**



HO^- **products**

name of mechanism .....

type of bond fission ..... **[5]**

[Total: 15]

END OF QUESTION PAPER

© OCR 2012

ADDITIONAL PAGE

If additional space is required, you should use the lined pages below. The question number(s) must be clearly shown.

© OCR 2012

ADDITIONAL PAGE

RECOGNISING ACHIEVEMENT

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© OCR 2012