OCR
Oxford Cambridge and RSA

A Level Chemistry A

H432/03 Unified chemistry

Tuesday 27 June 2017 – Morning

Time allowed: 1 hour 30 minutes

You must have:
- the Data Sheet for Chemistry A (sent with general stationery)

You may use:
- a scientific or graphical calculator
- a ruler

First name									
Last name									
Centre number						Candidate number				

INSTRUCTIONS
- Use black ink. You may use an HB pencil for graphs and diagrams.
- Complete the boxes above with your name, centre number and candidate number.
- Answer **all** the questions.
- Write your answer to each question in the space provided. If additional space is required, use the lined page(s) at the end of this booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.

INFORMATION
- The total mark for this paper is **70**.
- The marks for each question are shown in brackets **[ ]**.
- Quality of extended responses will be assessed in questions marked with an asterisk (*****).
- This document consists of **20** pages.

© OCR 2017 [601/5255/2]
DC (NF/JG) 139140/3
OCR is an exempt Charity

Answer **all** the questions.

1 Within the permafrost in Arctic regions of the Earth, large amounts of methane are trapped within ice as 'methane hydrate', $CH_4 \bullet xH_2O$. Methane makes up about 13.4% of the mass of 'methane hydrate'.

Scientists are concerned that global warming will melt the permafrost, releasing large quantities of methane into the atmosphere.

(a) The H–O–H bond angle in ice is about 109° but about 105° in gaseous H_2O.

Explain why there is this difference.

.....

.....

.....

.....

.....

..... **[3]**

(b) Why are scientists concerned about the release of methane into the atmosphere?

.....

.....

..... **[1]**

(c) Determine the formula of 'methane hydrate', $CH_4 \bullet xH_2O$.

In the formula, show the value of x to **two** decimal places.

formula = ... **[2]**

© OCR 2017

(d) Calculate the volume of methane, in dm^3, that would be released from the melting of each 1.00 kg of 'methane hydrate' at 101 kPa and 0 °C.

Give your answer to **three** significant figures.

volume = ... dm^3 **[4]**

(e) Suggest why some industries are interested in the presence of 'methane hydrate' in regions of the Earth.

.....

.....

..... **[1]**

© OCR 2017

2 A student plans to determine the enthalpy change of **reaction 3.1** shown below.

$$Na_2O(s) + 2HCl(aq) \rightarrow 2NaCl(aq) + H_2O(l)$$ **reaction 3.1**

This enthalpy change can be determined indirectly using Hess' Law from the enthalpy changes of **reaction 3.2** and **reaction 3.3** shown below.

$$Na_2O(s) + H_2O(l) \rightarrow 2NaOH(aq)$$ **reaction 3.2**

$$HCl(aq) + NaOH(aq) \rightarrow NaCl(aq) + H_2O(l) \quad \Delta_r H = -57.6\,kJ\,mol^{-1}$$ **reaction 3.3**

The student will determine the enthalpy change of **reaction 3.2** as outlined below.

- Weigh a bottle containing $Na_2O(s)$ and weigh a polystyrene cup.
- Add about $25\,cm^3$ of water to the polystyrene cup and measure its temperature.
- Add the $Na_2O(s)$, stir the mixture, and measure the maximum temperature reached.
- Weigh the empty bottle and weigh the polystyrene cup with the final solution.

Mass readings

Mass of bottle + $Na_2O(s)$	= 16.58 g
Mass of empty bottle	= 15.34 g
Mass of empty polystyrene cup	= 21.58 g
Mass of polystyrene cup + final solution	= 47.33 g

Temperature readings

Initial temperature of water	= 20.5 °C
Maximum temperature of final solution	= 55.5 °C

The density and specific heat capacity, *c*, of the solution are the same as for water.

© OCR 2017

(a)* Calculate the enthalpy change of **reaction 3.2** and the enthalpy change of **reaction 3.1**.

Show all your working.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... **[6]**

(b) The uncertainty in each temperature reading is ±0.1 °C.

The uncertainty in each mass reading is ±0.005 g.

Determine whether the mass of Na_2O or the temperature change has the greater percentage uncertainty.

Show all your working.

.....

.....

.....

.....

..... **[2]**

© OCR 2017

(c) Suggest a modification to this experiment, using the **same** apparatus, which would reduce the percentage errors in the measurements.

Explain your reasoning.

.....

.....

.....

..... **[2]**

(d) Sodium oxide, Na_2O, can be prepared by the redox reaction of $NaNO_2$ and sodium metal. Nitrogen gas is also formed.

(i) What is the systematic name for $NaNO_2$?

..... **[1]**

(ii) Using oxidation numbers, with signs, show the element that is oxidised and the element that is reduced in this reaction.

Element oxidised .....

Oxidation number change from ..... to .....

Element reduced .....

Oxidation number change from ..... to .....

[2]

(iii) Construct the equation for this reaction.

Equation ..... **[1]**

© OCR 2017

3 This question is about reactions of hydrogen peroxide, H_2O_2.

(a) Hydrogen peroxide, H_2O_2, iodide ions, I^-, and acid, H^+, react as shown in the equation below.

$$H_2O_2(aq) + 2I^-(aq) + 2H^+(aq) \rightarrow I_2(aq) + 2H_2O(l)$$

A student carries out several experiments at the same temperature, using the initial rates method, to determine the rate constant, *k*, for this reaction.

The results are shown below.

Experiment	Initial concentrations			Rate / 10^{-6} mol dm^{-3} s^{-1}
	[H_2O_2(aq)] / mol dm^{-3}	[I^-(aq)] / mol dm^{-3}	[H^+(aq)] / mol dm^{-3}	
1	0.0100	0.0100	0.100	2.00
2	0.0100	0.0200	0.100	4.00
3	0.0200	0.0100	0.100	4.00
4	0.0200	0.0100	0.200	4.00

(i) Determine the rate equation and calculate the rate constant, *k*, including units.

k = ... units ..... **[3]**

(ii) The rate constant, *k*, for this reaction is determined at different temperatures, *T*.

Explain how the student could determine the activation energy, E_a, for the reaction graphically using values of *k* and *T*.

.....

.....

.....

.....

.....

..... **[3]**

© OCR 2017

(b) Solutions of hydrogen peroxide decompose slowly into water and oxygen:

$$2H_2O_2(aq) \rightarrow 2H_2O(l) + O_2(g)$$

This reaction is catalysed by manganese dioxide, $MnO_2(s)$.

Standard electrode potentials are shown below.

$O_2(g) + 2H^+(aq) + 2e^-$	$\rightleftharpoons$	$H_2O_2(aq)$	$E^\ominus = +0.70\,V$
$MnO_2(s) + 4H^+(aq) + 2e^-$	$\rightleftharpoons$	$Mn^{2+}(aq) + 2H_2O(l)$	$E^\ominus = +1.51\,V$
$H_2O_2(g) + 2H^+(aq) + 2e^-$	$\rightleftharpoons$	$2H_2O(l)$	$E^\ominus = +1.78\,V$

Using the electrode potentials, explain how MnO_2 is able to act as a catalyst for the decomposition of hydrogen peroxide.

You answer should include relevant equations.

.....

.....

.....

.....

.....

.....

.....

.....

..... **[4]**

© OCR 2017

(c) Peroxycarboxylic acids are organic compounds with the COOOH functional group.

Peroxyethanoic acid, CH_3COOOH, is used as a disinfectant.

(i) Suggest the structure for CH_3COOOH.

The COOOH functional group must be clearly displayed.

[1]

(ii) Peroxyethanoic acid can be prepared by reacting hydrogen peroxide with ethanoic acid. This is a heterogeneous equilibrium.

$$H_2O_2(aq) + CH_3COOH(aq) \rightleftharpoons CH_3COOOH(aq) + H_2O(l) \quad K_c = 0.37\,dm^3\,mol^{-1}$$

A 250 cm^3 equilibrium mixture contains concentrations of 0.500 $mol\,dm^{-3}$ $H_2O_2(aq)$ and 0.500 $mol\,dm^{-3}$ $CH_3COOH(aq)$.

Calculate the amount, in mol, of peroxyethanoic acid in the equilibrium mixture.

amount = ... mol **[3]**

© OCR 2017

4 This question is about weak acids.





(a) Compound **A** is a weak monobasic acid.

A student is supplied with a 250.0 cm^3 solution prepared from 2.495 g of **A**.

The student titrates 25.0 cm^3 samples of this solution with 0.0840 mol dm^{-3} NaOH in the burette.

The student carries out a trial, followed by the three further titrations. The diagrams show the initial burette readings and the final burette readings for the student's three **further** titrations.

All burette readings are measured to the nearest 0.05 cm^3.

Titration 1		**Titration 2**		**Titration 3**	
Initial reading	Final reading	Initial reading	Final reading	Initial reading	Final reading
					


(i) Record the student's readings and the titres in an appropriate format.

Calculate the mean titre that the student should use for analysing the results.

mean titre = ... cm^3 **[4]**

© OCR 2017

(ii) The structure of compound **A** is shown below.



compound A

Compound **A** has four optical isomers.

Using this information and the student's results, answer the following.


- Determine the molar mass of **A** and the formula of the alkyl group R.
- Draw the structure of compound **A** and label any chiral carbon atoms with an asterisk*.

Show all your working.

[6]

© OCR 2017

(b) The structural formula of compound **A** is repeated below.



compound A

Two reactions of compound **A** are carried out.

Suggest an equation for each reaction and state the type of reaction.

In your equations, draw structures for organic compounds.
You can use R for the alkyl group.

(i) Magnesium ribbon is added to a solution of compound **A**.
Gas bubbles are seen and the magnesium slowly dissolves.

Equation

Type of reaction ..... **[3]**


(ii) Compound **A** is heated with a few drops of concentrated sulfuric acid as a catalyst.
A cyclic 'dimer' of compound **A** forms.

Equation

Type of reaction ..... **[3]**

© OCR 2017

(c) Chromium(III) picolinate, shown below, is a neutral complex that can be prepared from the weak acid, picolinic acid.



Chromium(III) picolinate is used in tablets as a nutritional supplement for chromium.

(i) Draw the structure of the ligand in chromium(III) picolinate.

[1]

(ii) A typical tablet of chromium(III) picolinate contains 200 μg of chromium.

Calculate the mass, in g, of chromium(III) picolinate in a typical tablet.
1 μg = 10^{-6} g.

Give your answer to **three** significant figures.


mass = ... g **[2]**

© OCR 2017

5 This question is about organic molecules that have a strong smell.

(a) Thiols are foul-smelling, organic sulfur compounds with the functional group –SH.

Butane-1-thiol, shown below, contributes to the strong smell of skunks.



butane-1-thiol

(i) Thiols are weak acids.

Write the expression for the acid dissociation constant, K_a, for butane-1-thiol.

[1]

(ii) Thiols react with carboxylic acids to form thioesters.

Write an equation for the reaction of butane-1-thiol with ethanoic acid.

Use structures for all organic compounds with the functional groups clearly displayed.

[2]

(iii) When beer is exposed to light, 3-methylbut-2-ene-1-thiol is formed, which gives an unpleasant smell and flavour to the beer.

Draw the **skeletal** formula for 3-methylbut-2-ene-1-thiol.

[1]

© OCR 2017

(iv) Propane-1,3-dithiol reacts with carbonyl compounds in a condensation reaction to form a cyclic organic sulfur product.


Write an equation for the reaction of propane-1,3-dithiol with propanone.

Use structures for organic compounds.


[2]

© OCR 2017


(b)* The structures for six naturally occurring organic compounds with pleasant smells, **B**–**G**, are shown below. The common names in brackets relate to their source and smell.




B
(linalool)




C
(citral)




D
(geraniol)



E
(camphor)



F
(menthol)



G
(ionone)

© OCR 2017

Explain how chemical tests would allow each compound to be distinguished from the other compounds.

In your answer, include essential details for all test procedures and observations.

Details of apparatus and quantities are **not** required.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... **[6]**

END OF QUESTION PAPER

© OCR 2017

ADDITIONAL ANSWER SPACE

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).

© OCR 2017

© OCR 2017

OCR
Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© OCR 2017