



Mark Scheme (Results)

January 2018

Pearson Edexcel International Advanced
Level In Chemistry (WCH06) Paper 01
Chemistry Laboratory Skills II

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2018

Publications Code WCH06_01_1801_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question Number	Acceptable Answer	Reject	Mark
1(a)	(Contains) a transition metal (ion) / a transition element ALLOW a d-block element (Might contain) chromate(VI) ions/ CrO_4^{2-} / Iron(III) (ions)/ Fe^{3+} / PbI_2 / AgI	Dichromate (ions) / $\text{Cr}_2\text{O}_7^{2-}$ / Cr^{6+}	(1)

Question Number	Acceptable Answer	Reject	Mark
1(b)	Na^+ (ion) / sodium (cat)ion present ALLOW Na ion Na_2CrO_4	Just 'Na' / sodium $\text{Na}_2\text{Cr}_2\text{O}_7$ NaCl / any other sodium compounds	(1)

Question Number	Acceptable Answer	Reject	Mark
1(c)	$\text{Cr}_2\text{O}_7^{2-}$ $(\text{Cr}_2\text{O}_7)^{2-}$	CrO_4^{2-} $\text{Na}_2\text{Cr}_2\text{O}_7$	(1)

Question Number	Acceptable Answer	Reject	Mark
1(d)	1st mark: $\text{Cr}_2\text{O}_7^{2-}$ / dichromate(VI) / chromium(VI) reduced ALLOW Chromium is reduced OR ethanol is oxidised OR ethanol forms ethanoic acid / ethanol forms ethanal (1) 2nd mark: Cr^{3+} / chromium(III) / Cr(III) (ions) formed (1) Mark scoring points independently	Cr^{6+} is reduced Just 'oxidation' or 'reduction' or 'redox'	(2)

Question Number	Acceptable Answer	Reject	Mark
1(e)	1st mark: (Green precipitate formula is) $\text{Cr}(\text{OH})_3(\text{H}_2\text{O})_3$ / $\text{Cr}(\text{OH})_3$ (1) 2nd mark: (Ion responsible for green colour of solution) $\text{Cr}(\text{OH})_6^{3-}$ ALLOW $\text{Cr}(\text{OH})_4^-$ / $\text{Cr}(\text{OH})_5^{2-}$ / CrO_2^- / CrO_3^{3-} (1) IGNORE Any number of H_2O ligands included with $\text{Cr}(\text{OH})_4^-$ / $\text{Cr}(\text{OH})_5^{2-}$	Cr_2O_3 Cr^{3+} / $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$	(2)

Question Number	Acceptable Answer	Reject	Mark
1(f)	1st mark: (Ion responsible for pale blue colour of solution) $\text{Cr}(\text{H}_2\text{O})_6^{2+} / \text{Cr}^{2+}$ (1) 2nd mark: (Role of the Zn) Reducing agent / reduces / reduction / loses electrons / donates electrons (1) IGNORE References to redox Mark independently	Cu^{2+}	(2)

Question Number	Acceptable Answer	Reject	Mark
1(g)	1st mark: (Ion responsible for green colour of the solution) $\text{Cr}(\text{H}_2\text{O})_6^{3+} ((\text{aq})) / \text{Cr}^{3+} ((\text{aq}))$ (1) 2nd mark: (Type of reaction) Oxidation ALLOW 'redox' (1) Mark independently	Other numbers of water ligands Reduction Ligand exchange	(2)

(Total for Question 1 = 11 marks)

Question Number	Acceptable Answer	Reject	Mark
2(a)(i)	EITHER (Y contains an) OH (group) / hydroxy(l) (group) OR (Y is an) alcohol or a carboxylic acid OR (Y contains) an OH or a COOH (group) OR (Y is) propanol or propanoic acid / propan-1-ol or propan-2-ol or propanoic acid Note The significance of 'or'	"hydroxide/ OH ⁻ "	(1)

Question Number	Acceptable Answer	Reject	Mark
2(a)(ii)	(Y is) NOT a (carboxylic) acid / does not contain a carboxyl group ALLOW (Y is) an alcohol	Does not contain a carboxylate group	(1)

Question Number	Acceptable Answer	Reject	Mark
2(a)(iii)	(Y contains a) CH₃CH(OH) group / (Y is a) secondary methyl alcohol / 2° alcohol with methyl group next to C-OH ALLOW Secondary alcohol with methyl group next to the functional group OR (Y is) propan-2-ol IGNORE References to methyl ketone / ethanol / just 'secondary alcohol' / just 'CHOH group'		(1)

Question Number	Acceptable Answer	Reject	Mark
2(a)(iv)	(Y is) propan-2-ol / CH₃CH(OH)CH₃ ALLOW Skeletal / displayed formula	propanol Just C₃H₈O	(1)


Question Number	Acceptable Answer	Reject	Mark
2(a)(v)	1st mark: (Peak caused by) CH₃CHOH⁺ / C₂H₅O⁺ ALLOW TE on structure in (iv) e.g. CH₂CH₂OH⁺ if propan-1-ol given in (iv) (1) 2nd mark: (Species is formed by) (Fragmentation of molecular ion by) loss of a CH₃ group / •CH₃ / CH₃ radical / methyl group / methyl radical ALLOW Breaking a single carbon to carbon bond Loss of CH₃⁺ (1)	MP1 if + sign missing CH₃CH₂O⁺ Loss of CH₃⁻	(2)

Question Number	Acceptable Answer	Reject	Mark
2(b)(i)	Hydrogen chloride / HCl / HCl(g) / HCl(gas) ALLOW Hydrochloric acid/ HCl(aq)		(1)

Question Number	Acceptable Answer	Reject	Mark
2(b)(ii)	(Z contains the functional group) Carboxylic acid / (-)COOH / (-)CO ₂ H ALLOW Carboxylic group / carboxyl group IGNORE Just 'acid' Carbon dioxide is produced	'Carboxylate'	(1)

Question Number	Acceptable Answer	Reject	Mark
2(b)(iii)	(Z is) Propanoic acid / CH ₃ CH ₂ COOH / CH ₃ CH ₂ CO ₂ H / C ₂ H ₅ COOH / C ₂ H ₅ CO ₂ H ALLOW Skeletal or displayed formula		(1)

(Total for Question 2 = 9 marks)

Question Number	Acceptable Answer	Reject	Mark
3(a)(i)	 1st mark: Both axes labelled, with units ALLOW units in brackets (1) 2nd mark: 'Sensible' scale, covering more than half the grid in each direction (1) 3rd mark: All six points plotted correctly (1) 4th mark: Straight line of best fit ALLOW If a non-uniform scale is used (1)	If units missing, no MP1 Non-uniform scale Reject if a non-uniform scale is used in MP2 point-to-point' if misplot/ curve of any description	(4)

Question Number	Acceptable Answer	Reject	Mark
3(a)(ii)	The iodine concentration is proportional to the titre / volume of thiosulfate IGNORE Use of 1:2 ratio		(1)

Question Number	Acceptable Answer	Reject	Mark
3(a)(iii)	The concentrations of the propanone and hydrochloric acid are (effectively) constant (during the reaction) (1) (So that) only the iodine concentration changes (during the reaction) ALLOW Iodine concentration is the only variable (So) the order is that with respect to iodine alone (1)		(2)

Question Number	Acceptable Answer	Reject	Mark
3(a)(iv)	Zero (order) / 0 (order) / $c = 0$ (1) (The straight line graph shows) rate is independent of I_2 concentration / rate (of decrease of I_2 concentration) is constant / gradient is constant (1)	Just 'the graph is a straight line'	(2)

Question Number	Acceptable Answer	Reject	Mark
3(b)(i)	Starch (solution) (1) From blue-black / from blue / from black to colourless (1) IGNORE 'Clear' Colour change is needed for this mark	Purple /Pale blue Off white	(2)

Question Number	Acceptable Answer	Reject	Mark
3(b)(ii)	(When solution/reaction mixture is) pale-yellow / 'straw' coloured (1) IGNORE 'Near end-point' To prevent the formation of an insoluble complex/ insoluble compound / insoluble substance / insoluble starch-iodine complex (1)	Just 'yellow' Just 'goes pale'	(2)

Question Number	Acceptable Answer	Reject	Mark
3(c)	Titre is small(est) (so the (relative) uncertainty/(relative) error is greatest)		(1)

Question Number	Acceptable Answer	Reject	Mark
3(d)	(Order is) first / 1st / 1 (1) Doubling the concentration (of the acid) doubles the rate (1)	'Speed' instead of rate	(2)

Question Number	Acceptable Answer	Reject	Mark
3(e)	Any ONE of the following matching pairs: Colorimetry (1) I_2 colour decreases / changes ALLOW Iodine / I_2 is coloured Brown to colourless (1) ALLOW Conductivity (1) Increase in ion / H^+ / I^- concentration (1) OR pH (1) Increase in H^+ concentration (1) OR Acid-base titration (1) Increase in H^+ concentration (1) OR $AgNO_3$ titration (1) Increase in I^- concentration (1)	Increase in pH	(2)

(Total for Question 3 = 18 Marks)

Question Number	Acceptable Answer	Reject	Mark
4(a)	1st mark: (As liquid boils / vaporises), it / vapour is cooled and condensed (back to liquid) (1) 2nd and 3rd marks: Any TWO from • Prevents escape of reactants / products / flammable liquids / flammable vapours / volatile liquids • Enables (in)flammable / harmful liquids to be heated safely • Allows time for reaction / allows for a complete reaction / increases rate of reaction / increases yield (of product) / overcome the activation energy (for the reaction) (2)		(3)

Question Number	Acceptable Answer	Reject	Mark
4(b)(i)	To remove / react with the (excess/unreacted) ethanoyl chloride ALLOW Hydrolysis of the ethanoyl chloride	Cool down / dilute the mixture / solution	(1)

Question Number	Acceptable Answer	Reject	Mark
4(b)(ii)	The reaction (with water) is (very) exothermic / gives out heat / is violent / is vigorous IGNORE explosive		(1)

Question Number	Acceptable Answer	Reject	Mark
4(c)	(Filtration is) faster / quicker (1) Dries the solid / dries the precipitate / dries the crystals OR Removes the maximum amount of solution IGNORE References to 'higher yield' / 'less product lost' / fewer 'transfer losses' (1)		(2)

Question Number	Acceptable Answer	Reject	Mark
4(d)	'Sharp' melting temperature / melts over a 1-2°C range / melts over a 'narrow' temp range (1) Value (of melting temperature is) consistent with literature value / Internet value ALLOW 'Correct value' (obtained) (1)	Sharper melting temperature narrower	(2)

Question Number	Acceptable Answer	Reject	Mark
4(e)	SCROLL DOWN TO CHECK WHOLE SPACE TE is allowed at each stage <u>Correct answer with no working scores 3</u> IGNORE SF except 1 SF Moles of $C_7H_7NO_2$ $(= \frac{3.70}{137}) = 0.027007299$ $= 0.0270 \text{ (mol)}$ (1) Actual moles of product, $C_9H_9NO_3$ $(= \frac{2.42}{179}) = 0.013519553$ $= 0.0135 \text{ (mol)}$ (1) % yield $(= \frac{0.0135}{0.0270} \times 100\%)$ $= 50\%$ (1) OR Alternative route for MP2 and MP3 Expected mass of product, $C_9H_9NO_3$ $(= 0.0270 \times 179) = 4.833 \text{ (g)}$ (1) % yield $(= \frac{2.42}{4.833} \times 100\%)$ $= 50.07241879\%$ $= 50.1\%$ (1)	Just $\frac{2.42}{3.70} \times 100$ $= 65.4\%$	3

	OR FURTHER ALTERNATIVE ROUTE Expected moles of product, $\text{C}_9\text{H}_9\text{NO}_3$ $\frac{3.70}{137} = 0.027007299$ (1) Expected mass of product, $\text{C}_9\text{H}_9\text{NO}_3$ $0.027007299 \times 179 = 4.834(\text{g})$ (1) % yield ($= \frac{2.42}{4.834} \times 100\%$) $= 50.06206041\%$ $= 50.1\%$ (1) NOTE Final answer will depend on figures held in calculator by candidate – all working must be checked e.g if all numbers held in calculator, value equals 50.0588857 %		
--	---	--	--

(Total for Question 4 = 12 Marks)

TOTAL FOR PAPER = 50 MARKS