UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2011 question paper

for the guidance of teachers

9701 CHEMISTRY

9701/22 Paper 2 (AS Structured Questions), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

UNIVERSITY of CAMBRIDGE
International Examinations

1 **(a)** $K_c = \dfrac{[CH_3CH_2R][H_20]}{[CH_3CH_2H][ROH]}$ (1)

no units (1) [2]

(b) **(i)** $n(NaOH) = \dfrac{22.5 \times 2.00}{1000} = 0.045$ (1)

(ii) $n(NaOH) = n(HCl) = 0.005$ (1)

(iii) $CH_3CO_2H + NaOH \rightarrow CH_3CO_2Na + H_2O$ (1)

(iv) $n(NaOH) = 0.045 - 0.005 = 0.04$
allow ecf on **(i)** and/or **(ii)** (1) [4]

(c) **(i)** $n(NaOH)$ **and** $n(CH_3CO_2H) = 0.04$ (1)
$n(CH_3CO_2R)$ **and** $n(H_2O) = 0.06$ (1)

(ii) $K_c = \dfrac{0.06 \times 0.06}{0.04 \times 0.04} = 2.25$

allow ecf on wrong values in **(b)(i)**
allow ecf on wrong expression in **(a)** (1) [3]

(d) E_a for reaction with ester is high **or**
E_a for reaction with acid is low
or
reaction with ester is slow **or**
reaction with acid is fast (1) [1]

(e) equilibrium moves to RHS/more ester would be formed (1)
to maintain value of K_c **or**
to restore system to equilibrium (1) [2]

[Total: 12]

© University of Cambridge International Examinations 2011

2 (a) $CH_2=CH_2 + HF \rightarrow CH_3CH_2F$

bonds	4 C-H	1640	bonds	5 C-H	2050
broken	1 C=C	610	made	1 C-C	350
/kJ mol^{-1}	1 H-F	562	/kJ mol^{-1}	1 C-F	*E*
		2812			(2400 + *E*)

breaking reactant bonds requires
$4 \times 410 + 610 + 562 = 2812$ kJ mol^{-1} (1)

making product bonds gives

$5 \times 410 + 350 + E = (2400 + E)$ kJ mol^{-1} (1)

$\Delta H^{\ominus}_{reaction} = -(2400 + E) + 2812 = -73$ kJ mol^{-1} (1)

$(2400 + E) = 2812 + 73 = 2885$ kJ mol^{-1}

$E = 2885 - 2400 = 485$ kJ mol^{-1} (1)

allow ecf on wrong bond energy values and/or incorrect arithmetic [4]

(b) any **two** from
non-toxic
unreactive
volatile
non-flammable
easily liquefied (1 + 1) [2]

(c) in CCl_2F_2
C-C*l* bond energy is 340 kJ mol^{-}1 and is weaker than C-F or C-H bonds (1)
C-C*l* bond is broken by uvl **or**
C*l* free radicals are formed (1) [2]

(d) (i) the trapping of reflected heat from the Earth in the lower atmosphere producing global warming

(ii) CO_2/carbon dioxide (1) [3]

(e) octahedral (1) [1]

[Total: 12]

© University of Cambridge International Examinations 2011

3



(a) **R** BaO (1)
S $BaCl_2$ (1)
T $Ba(OH)_2$ (1)
U $BaSO_4$ (1)
V $BaCO_3$ (1)
W $Ba(NO_3)_2$ (1) [6]

(b) **(i)** **T** to **W**
$Ba(OH)_2 + 2HNO_3 \rightarrow Ba(NO_3)_2 + 2H_2O$ (1)

heat on **V**
$BaCO_3 \rightarrow BaO + CO_2$ (1)

(ii) **T** to **V**
CO_2 (1)
$Ba(OH)_2 + CO_2 \rightarrow BaCO_3 + H_2O$ (1) [4]

(c) $Na_2SO_4(aq)/K_2SO_4(aq)$ or any soluble sulfate (1) [1]

© University of Cambridge International Examinations 2011

(d) (i) Ba:O = $\frac{81.1}{137} : \frac{18.9}{16}$ (1)

= 0.59 : 1.18
= 1 : 2
gives BaO_2 (1)

(ii) $BaSO_4$ (1)

(iii) $BaO_2 + H_2SO_4 \rightarrow BaSO_4 + H_2O_2$ (1) [4]

[Total: 15]

4 (a) (i)



titanium/graphite anode identified correctly (1)
steel cathode identified correctly (1)
diaphragm identified correctly (1)
all three outlets correctly shown (1) [4]

(ii) **anode** $2Cl^-(aq) \rightarrow Cl_2(g) + 2e^-$ (1)

cathode $2H^+(aq) + 2e^- \rightarrow H_2(g)$

or $2H_2O(l) + 2e^- \rightarrow H_2(g) + 2OH^-(aq)$ (1) [2]

(iii) sodium hydroxide (1) [1]

[Total: 7]

© University of Cambridge International Examinations 2011

5 **(a)**



all three alcohol groups must be esterified (1) [1]

(b) dilute HC*l* **or** dilute H_2SO_4 **or** dilute mineral acid
or NaOH(aq) **followed by** dilute acid (1) [1]

(c)



(1) [1]

(d) **(i)** fatty acid that contains more than one C=C bond (1)

(ii) hydrogen (1)
nickel/Raney nickel/platinum/palladium (1) [3]

(e) **(i)** $CH_3(CH_2)_7CHO$ (1)
$OHC(CH_2)_7CX$ (1)

(ii) 2,4-dinitrophenylhydrazine (1)
yellow/orange/red precipitate (1)

(iii) Tollens' reagent **or** Fehling's/Benedict's solution (1)
silver mirror/ **or** brick red ppt.
grey precipitate (1) [6]

(f) **(i)** two (1)

(ii) ester (1) [2]

[Total: 14]

© University of Cambridge International Examinations 2011