UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

*0337350796*

**CHEMISTRY** **9701/53**

Paper 5 Planning, Analysis and Evaluation **October/November 2012**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams, graphs or rough working.
Do not use staples, paper clips, highlighters, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.
You may lose marks if you do not show your working or if you do not use appropriate units.
Use of a Data Booklet is unnecessary.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.

| For Examiner's Use | |
|---|---|
| 1 | |
| 2 | |
| 3 | |
| Total | |

This document consists of **8** printed pages.


© UCLES 2012

For Examiner's Use

**1** The particles of a gas are considered to be in constant random motion. For a fixed mass of gas in a container, the particles collide with the container walls and it is these collisions that are responsible for the pressure exerted by the gas.

The more frequent the collisions, the greater the pressure. If the temperature is kept constant, the average kinetic energy of the gas remains constant. A temperature increase means an increase in this average kinetic energy.

**(a) (i)** Predict how the pressure of a fixed mass of gas held at a constant temperature varies as the volume of the gas decreases. Explain this prediction in terms of the frequency of the gas collisions with the container walls.

Predict how the pressure will change .....

.....

Explanation .....

.....

.....

**(ii)** Display your prediction in the form of a sketch graph, labelling clearly the axes.




**(iii)** Sketch a second graph showing how your projected relationship will change if the temperature of the fixed mass of gas is increased. Label this second graph clearly.
[3]

**(b)** If you were to carry out an experiment to investigate how the **pressure** of a fixed mass of gas varies as the **volume decreases**, name

**(i)** the independent variable .....

**(ii)** the dependent variable .....
[2]

[Total: 5]

© UCLES 2012

**2** Copper has two oxides, $Cu_2O$ and CuO.
Copper(II) carbonate, $CuCO_3$, decomposes on heating to form one of these oxides. Separate equations can be written showing the two possible decompositions.

**2.1** $2CuCO_3(s) \rightarrow Cu_2O(s) + 2CO_2(g) + \frac{1}{2}O_2(g)$

**2.2** $CuCO_3(s) \rightarrow CuO(s) + CO_2(g)$

The following information gives some of the hazards associated with copper(II) carbonate.

**Copper(II) carbonate**; Harmful if swallowed.
Dispose of by reacting no more than 60 g in 1 $dm^3$ of warm 1 mol $dm^{-3}$ ethanoic acid before pouring down a foul-water drain. This procedure should be kept to a minimum.

You are to plan an experiment to investigate the decomposition of copper(II) carbonate on heating and hence decide which of the two equations represents the actual decomposition. The volume of gas liberated per mole of copper(II) carbonate depends on which equation is correct and this should be used as the basis of the plan.

**(a)** Draw a diagram of the apparatus and experimental set up you would use in the experiment. Your apparatus should use only standard items found in a school or college laboratory and should show clearly

**(i)** how the copper(II) carbonate will be heated,

**(ii)** how the volume of the gas evolved will be collected and its volume measured.

Label each piece of apparatus used, indicating its size or capacity.

[3]

© UCLES 2012

For Examiner's Use

**(b)** Using the apparatus shown in **(a)**, design a laboratory experiment which will enable you to determine the way in which copper(II) carbonate decomposes on heating.

In addition to the standard apparatus present in a laboratory, you are provided with the following material.

a sample of copper(II) carbonate

Give a step-by-step description of how you would carry out the experiment by

**(i)** calculating the volume of gas evolved by decomposing 1 mole of copper(II) carbonate according to equation **2.1**.

**(ii)** calculating the volume of gas evolved by decomposing 1 mole of copper(II) carbonate according to equation **2.2**.

**(iii)** using these results to calculate a mass of copper(II) carbonate that would give volumes of gas that could be collected using the apparatus proposed, under either decomposition.

**(iv)** stating how you would ensure that the decomposition is complete.

**(v)** stating how you would use your results to reach a conclusion.

[$A_r$: C, 12.0; O, 16.0; Cu, 63.5; the molar gas volume at 25 °C is 24.0 $dm^3$]

[5]

© UCLES 2012

*For Examiner's Use*

**(c)** State one hazard that must be considered when planning the experiment and describe a precaution that should be taken to keep risks to a minimum.

.....

.....

.....

.....

..... [2]

[Total: 10]

© UCLES 2012



**[Turn over**

*For Examiner's Use*

**3 (a)** The concentration of aqueous sulfuric acid can be determined by measuring the density of the solution.

In order to establish a calibration graph, various solutions of known percentage composition by mass were set up using very accurate apparatus.

The preparation of aqueous sulfuric acid is exothermic. Once each of the solutions had cooled, the actual density was determined using a hydrometer.

The table below shows some of the calculations involved, together with the measured densities.

To calculate the actual volumes of sulfuric acid and water mixed, the mass was used in conjunction with the appropriate density.

The density of pure water is 0.997 g $cm^{-3}$ and the density of pure sulfuric acid is 1.826 g $cm^{-3}$.

All measurements were carried out at 25 °C.

| percentage by mass of sulfuric acid | mass of sulfuric acid / g | mass of water / g | volume of sulfuric acid / $cm^3$ | volume of water / $cm^3$ | total volume of 100 g of solution / $cm^3$ | calculated density of the solution / g $cm^{-3}$ | measured density of the solution / g $cm^{-3}$ |
|---|---|---|---|---|---|---|---|
| 0 | 0.000 | 100.000 | 0.000 | 100.301 | 100.301 | 0.997 | 0.997 |
| 10 | 10.000 | 90.000 | 5.476 | 90.271 | 95.747 | 1.044 | 1.064 |
| 20 | 20.000 | 80.000 | | 80.241 | 91.194 | | 1.137 |
| 30 | 30.000 | 70.000 | 16.429 | 70.211 | 86.640 | 1.154 | 1.215 |
| 40 | 40.000 | 60.000 | | | 82.087 | 1.218 | 1.299 |
| 50 | 50.000 | | 27.382 | | 77.532 | | 1.391 |
| 60 | 60.000 | 40.000 | 32.859 | 40.120 | | 1.370 | 1.494 |
| 70 | 70.000 | 30.000 | 38.335 | 30.090 | | | 1.606 |
| 80 | 80.000 | 20.000 | 43.812 | | 63.872 | 1.566 | 1.722 |
| 90 | 90.000 | 10.000 | | 10.030 | | 1.686 | 1.809 |
| 100 | 100.000 | 0.000 | 54.765 | 0.000 | 54.765 | 1.826 | 1.826 |

Complete the table. You may use the space below for any working for your calculations.

[4]

© UCLES 2012

**(b)** On the grid below, using appropriate scales and labelled axes, plot **two** curves to show the variation of **both** the calculated densities **and** the measured densities against the percentage of sulfuric acid. Label each curve.



[4]

*For Examiner's Use*

**(c) (i)** Using your curves, deduce the difference between the calculated and measured densities for the 45 % by mass of sulfuric acid mixture. State which of the two has the larger density.

**(ii)** By considering the molecular structures of the two liquids, suggest an explanation for the difference.

.....

.....

.....

.....

[3]

**(d)** You have 100.000 g of a 60 % by mass sulfuric acid mixture of a **calculated density** of 1.370 g $cm^{-3}$. Calculate the mass of water you would need to add to the 100.000 g in order to give a final **calculated density** of 1.154 g $cm^{-3}$.

[2]

**(e)** Attempting to determine the measured density of aqueous sulfuric acid in a school / college laboratory would involve using less accurate apparatus than in **(a)**.
Calculate the errors that could arise when measuring both the mass and volume of 100 g of sulfuric acid using a balance accurate to the nearest 0.01 g and a 100 $cm^3$ measuring cylinder accurate to the nearest 0.25 $cm^3$.

mass error

volume error

[2]

[Total: 15]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2012