Cambridge International AS & A Level

Cambridge International Examinations
Cambridge International Advanced Subsidiary and Advanced Level

PHYSICS **9702/21**

Paper 2 AS Level Structured Questions **May/June 2017**

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE®, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

® IGCSE is a registered trademark.

© UCLES 2017

Question	Answer	Marks
1(a)	(stress =) force / area **or** kg m s^{-2} / m^2	**B1**
	= kg m^{-1} s^{-2}	**A1**
1(b)(i)	$0.58 = 2\pi \times [(4 \times 0.500 \times 0.600^3) / (E \times 0.0300 \times 0.00500^3)]^{0.5}$	**C1**
	$E = [4\pi^2 \times 4 \times 0.500 \times (0.600)^3] / [(0.58)^2 \times 0.0300 \times (0.00500)^3]$ $= 1.35 \times 10^{10}$ (Pa)	**C1**
	= 14 (13.5) GPa	**A1**
1(b)(ii)1.	(accuracy determined by) the closeness of the value(s)/measurement(s) to the true value	**B1**
	(precision determined by) the range of the values/measurements	**B1**
1(b)(ii)2.	l is (cubed so) 3 × (percentage/fractional) uncertainty **and** T is (squared so) 2 × (percentage / fractional) uncertainty **and** (so) l contributes more	**B1**

Question	Answer	Marks
2(a)	resultant force (in any direction) is zero	**B1**
	resultant torque/moment (about any point) is zero	**B1**
2(b)(i)	$a = (v - u) / t$ **or** gradient **or** $\Delta v / (\Delta) t$	**C1**
	e.g. $a = (8.8 - 4.6) / (7.0 - 4.0) = 1.4\ \text{m s}^{-2}$	**A1**
2(b)(ii)	$s = 4.6 \times 4 + [(8.8 + 4.6) / 2] \times 3$	**C1**
	$= 18.4 + 20.1$ $= 39\ (38.5)\ \text{m}$	**A1**
2(b)(iii)	$\Delta E = ½ \times 95\ [(8.8)^2 - (4.6)^2]$	**C1**
	$= 3678 - 1005$ $= 2700\ (2673)\ \text{J}$	**A1**
2(b)(iv)1.	weight $= 95 \times 9.81$ (= 932 N)	**C1**
	vertical tension force = 280 sin 25° **or** 280 cos 65° (=118.3 N)	**C1**
	$F = 932 + 118$ $= 1100\ (1050)\ \text{N}$	**A1**
2(b)(iv)2.	horizontal tension force = 280 cos 25° **or** 280 sin 65° (= 253.8 N)	**C1**
	resultant force $= 95 \times 1.4$ (= 133 N)	**C1**
	$133 = 253.8 - R$ $R = 120\ (120.8)\ \text{N}$	**A1**

© UCLES 2017

Question	Answer	Marks
3(a)	$\rho = m / V$	**C1**
	$V = \pi d^2 L / 4$ **or** $\pi r^2 L$	**C1**
	weight = $2.7 \times 10^3 \times \pi\ (1.2 \times 10^{-2})^2 \times 5.0 \times 10^{-2} \times 9.81 = 0.60$ N	**A1**
3(b)(i)	the point from where (all) the weight (of a body) seems to act	**B1**
3(b)(ii)	$W \times 12$	**C1**
	$(0.25 \times 8) + (0.6 \times 38)$	**C1**
	W = (2 + 22.8) / 12 = 2.1 (2.07) N	**A1**
3(c)(i)	pressure changes with depth (in water) **or** pressure on bottom (of cylinder) different from pressure on top	**B1**
	pressure on bottom of cylinder <u>greater than</u> pressure on top **or** force (up) on bottom of cylinder <u>greater than</u> force (down) on top	**B1**
3(c)(ii)	anticlockwise moment reduced and reducing the weight of X reduces clockwise moment **or** anticlockwise moment reduced so clockwise moment now greater than (total) anticlockwise moment	**B1**

© UCLES 2017

Question	Answer	Marks
4(a)	(two) waves travelling (at same speed) in opposite directions overlap	**B1**
	waves (are same type and) have same frequency/wavelength	**B1**
4(b)(i)	λ = 12 / 250 (= 0.048 m)	**C1**
	distance = 1.5 × 0.048 = 0.072 m	**A1**
4(b)(ii)	T = 1 / 250 = 0.004 (s) or 4 (ms)	**C1**
	1. curve drawn is mirror image of that in Fig. 4.2 and labelled P	**A1**
	2. horizontal line drawn between A and B and labelled Q	**A1**

Question	Answer	Marks
5(a)	observed frequency is different to source frequency when source moves relative to observer	**B1**
5(b)	360 = (400 × 340) / (340 ± v)	**C1**
	v = 38 (37.8) m s^{-1}	**A1**
	away (from the observer)	**B1**

Question	Answer	Marks
6(a)	volt / ampere	**B1**
6(b)(i)	$R_T = [1/3.0 + 1/6.0]^{-1} + 4.0$ (= 6.0 Ω)	**C1**
	I = 1.5 / 6.0	**C1**
	= 0.25 A	**A1**
6(b)(ii)	V_B = 0.5 V I = 0.5 / 3.0 = 0.17 (0.167) A	**A1**
6(b)(iii)	$P = I^2R$ **or** VI **or** V^2/R	**C1**
	ratio = $(0.167^2 \times 3.0) / (0.25^2 \times 4.0)$ = 0.33	**A1**
6(c)(i)	vary/change/different radius/diameter/cross-sectional area (of wire)	**B1**
6(c)(ii)	$v = I / Ane$ ratio $= \frac{(I_B / A_B)}{(I_C / A_C)}$ **or** $\frac{I_B}{I_C} \times \frac{A_C}{A_B}$	**C1**
	($R \propto 1/A$ so) ratio $= \frac{I_B}{I_C} \times \frac{R_B}{R_C} = \frac{0.167}{0.25} \times \frac{3.0}{4.0}$ = 0.50	**A1**
6(d)(i)	0.25 A to 0.13 (0.125) A **or** halved	**A1**
6(d)(ii)	no change	**A1**

<table>
<tr><th>Question</th><th>Answer</th><th>Marks</th></tr>
<tr><td>7(a)(i)</td><td>(proton is uud so) $(2/3)e + (2/3)e - (1/3)e = e$</td><td>B1</td></tr>
<tr><td>7(a)(ii)</td><td>(neutron is udd so) $(2/3)e - (1/3)e - (1/3)e = 0$</td><td>B1</td></tr>
<tr><td>7(b)(i)</td><td><table><tr><td></td><td>β^-</td><td>β^+</td></tr><tr><td>nucleon number</td><td>90</td><td>64</td></tr><tr><td>proton number</td><td>39</td><td>28</td></tr></table>all correct</td><td>B1</td></tr>
<tr><td>7(b)(ii)</td><td>weak (nuclear force/interaction)</td><td>B1</td></tr>
<tr><td>7(b)(iii)</td><td>β^- decay: electron and (electron) antineutrino
β^+ decay: positron and (electron) neutrino
all correct</td><td>B1</td></tr>
</table>

© UCLES 2017