Version 1.0



General Certificate of Education (A-level)
June 2013

Mathematics MPC1

(Specification 6360)

Pure Core 1

Final

Mark Scheme

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2013 AQA and its licensors. All rights reserved.

Copyright
AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334).
Registered address: AQA, Devas Street, Manchester M15 6EX.

Key to mark scheme abbreviations

M	mark is for method
m or dM	mark is dependent on one or more M marks and is for method
A	mark is dependent on M or m marks and is for accuracy
B	mark is independent of M or m marks and is for method and accuracy
E	mark is for explanation
√or ft or F	follow through from previous incorrect result
CAO	correct answer only
CSO	correct solution only
AWFW	anything which falls within
AWRT	anything which rounds to
ACF	any correct form
AG	answer given
SC	special case
OE	or equivalent
A2,1	2 or 1 (or 0) accuracy marks
–*x* EE	deduct *x* marks for each error
NMS	no method shown
PI	possibly implied
SCA	substantially correct approach
c	candidate
sf	significant figure(s)
dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.


Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.



Q	Solution	Marks	Total	Comments
1(a)	$3p-4(p+2)+5=0$	M1		condone omission of brackets or one sign error
	$(\Rightarrow p=)\ -3$	A1	2	
(b)	$y=\frac{3}{4}x+\frac{5}{4}$	M1		rearranging into form $y=\pm\frac{3}{4}x+c$
	(gradient AB =) $\frac{3}{4}$	A1	2	condone slips in rearranging if gradient is correct .
(c)	(gradient AC =) $\frac{k-2}{-5-1}$	M1		*or* $\frac{2-k}{1--5}$ (condone one sign error)
	"their" $\frac{(k-2)}{-6}\times\frac{3}{4}=-1$ *OE*	m1		product of grads = –1 in terms of k
	$(\Rightarrow k=)\ 10$	A1	3	
(d)	$3x-4y+5=0$ and $2x-5y=6$			must use "correct" pair of equations **and**
	$\Rightarrow Px=Q$ or $Ry=S$	M1		attempt to eliminate y (or x) (generous)
	$x=-7$	A1		
	$y=-4$	A1	3	
				(–7, – 4)
	Total		**10**	

Q	Solution	Marks	Total	Comments
2(a)(i)	$\left(\sqrt{48}\ =\right)4\sqrt{3}$	B1	1	condone $n = 4$. No ISW .
(ii)	$\sqrt{12}\ = 2\sqrt{3}$ and $\sqrt{48}\ = 4\sqrt{3}$	M1		(FT 'their'n) $2x\sqrt{3} = 7\sqrt{3} - 4\sqrt{3}$
	$\left(x =\right)\dfrac{7\sqrt{3} - 4\sqrt{3}}{2\sqrt{3}}$	A1		correct quotient unsimplified or correct equation in integers eg $6x = 21 - 12$
	$= \dfrac{3}{2}$	A1cso	3	accept 1.5 but not $\dfrac{9}{6}$ etc **alternative 1** $x = \dfrac{7\sqrt{3} - \sqrt{48}}{\sqrt{12}} \times \dfrac{\sqrt{12}}{\sqrt{12}}$ M1 integer terms $= \dfrac{42 - 24}{12}$ A1 $= \dfrac{3}{2}$ A1
(b)	$\dfrac{11\sqrt{3} + 2\sqrt{5}}{2\sqrt{3} + \sqrt{5}} \times \dfrac{2\sqrt{3} - \sqrt{5}}{2\sqrt{3} - \sqrt{5}}$	M1		
	(numerator =) $22 \times 3 + 4\sqrt{15} - 11\sqrt{15} - 2 \times 5$	A1		correct unsimplified but must simplify $\left(\sqrt{3}\right)^2$, $\left(\sqrt{5}\right)^2$ and $\sqrt{3} \times \sqrt{5}$ correctly
	(denominator = 12 – 5 =) 7	B1		must be seen or identified as denominator giving $\dfrac{56 - 7\sqrt{15}}{7}$
	(Answer =) $8 - \sqrt{15}$	A1cso	4	$m = 8$
	Total		**8**	

Q	Solution	Marks	Total	Comments
3(a)	$(x-5)^2+(y+7)^2$	M1		one term correct
		A1		both terms correct and added
	$(x-5)^2+(y+7)^2=49$	A1cao	3	must see 49 not just 7^2 condone $(x-5)^2+(y--7)^2=49$
(b)(i)	(Centre is) $(5, -7)$	B1√	1	correct or FT their a and b
(ii)	Radius = 7	B1√	1	condone $\sqrt{49}$ but **not** ± 7 or $\pm\sqrt{49}$ correct or FT their $\sqrt{k}$ provided $k>0$
(c)(i)		M1		freehand circle with centre in correct quadrant or FT from their **(b)(i)** must have both axes shown clearly
		A1	2	correct position cutting negative y-axis twice and touching x-axis at $x=5$ 5 must be marked on x-axis or centre clearly marked as $(5, -7)$ must have correct centre and radius in **(b)**
(ii)	$x=5$	B1		
	$y=-14$	B1	2	$(5, -14)$
(d)	Translation	E1		and no other transformation
	through $\begin{bmatrix}6*\end{bmatrix}$	M1		
	$\begin{bmatrix}6\\-7\end{bmatrix}$	A1cso	3	both components correct for A1; may describe in words or use a column vector
	Total		**12**	

Q	Solution	Marks	Total	Comments
4(a)	$f(-3) = (-3)^3 - 4\times(-3) + 15$ $f(-3) = -27 + 12 + 15$	M1		f(–3) attempted **not** long division
	$= 0 \Rightarrow x + 3$ is a factor	A1	2	shown = 0 plus statement
(ii)	Quadratic factor $(x^2 - 3x + 5)$	M1		$-3x$ or + 5 term by inspection *or* full long division attempt
	$(f(x) =) \quad (x+3)(x^2 - 3x + 5)$	A1	2	must see correct product
(b) (i)	$\left(\frac{dy}{dx} =\right) 4x^3 - 16x + 60$	M1 A1		one of these terms correct another term correct
		A1	3	all correct (no +*c* etc)
(ii)	$4x^3 - 16x + 60 = 0$			must see this line OE
	$\Rightarrow x^3 - 4x + 15 = 0$	B1	1	**AG**
(iii)	Discriminant of quadratic = $(-3)^2 - 4\times 5$	M1		discriminant of " their" quadratic or correct use of quad eqn "formula"
	$b^2 - 4ac = -11$ (*or* $b^2 - 4ac < 0$) therefore quadratic has no (real)roots			**correct discriminant** evaluated correctly (or shown to be < 0) with appropriate conclusion
	Hence only stationary point is when $x = -3$	A1	2	**plus** final statement
(iv)	$\left(\frac{d^2y}{dx^2} =\right) 12x^2 - 16$	B1√		
	$= 12(-3)^2 - 16$ (*or* $12\times 9 - 16$ etc)	M1		sub $x = -3$ into "their" $\frac{d^2y}{dx^2}$
	$= 92$	A1	3	
(v)	Minimum **since** $\frac{d^2y}{dx^2} > 0$ (*or* 92 > 0 etc)	E1√	1	FT appropriate conclusion from their value from **(iv) plus** reason treat parts **(iv) & (v)** holistically
	Total		**14**	

Q	Solution	Marks	Total	Comments
5(a)(i)	$2(x+1.5)^2$	M1		OE
	$2(x+1.5)^2+0.5$	A1	2	$2\left(x+\frac{3}{2}\right)^2+\frac{1}{2}$ OE
(ii)	(Minimum value is) 0.5	B1√	1	ft their q
(b)(i)	$\left(AB^2=\right)\ (x+3)^2+(3x+9-5)^2$	M1		condone one sign error inside one bracket
	$(3x+4)^2=9x^2+24x+16$	B1		OE
	$AB^2=x^2+6x+9+9x^2+24x+16=10x^2+30x+25$ $\Rightarrow AB^2=5(2x^2+6x+5)$	A1cso	3	**AG**
(ii)	*Either* $\sqrt{5\times\text{'}their\text{'}}$ **(a)(ii)** *or* $5\times\text{'}their\text{'}$ **(a)(ii)**	M1		using their minimum value from **(a)(ii)** and 5 provided "their" **(a)(ii)** > 0
	(Minimum length of AB =) $\frac{1}{2}\sqrt{10}$	A1cso	2	
	Total		**8**	
6(a)	$\frac{dy}{dx}=5x^4-4x$	M1 A1		one of these terms correct all correct (no +c etc)
	$\left(=5(-1)^4-4(-1)\right)\quad=9$	A1		
	Tangent has equation $y=\text{'}their\ 9\text{'}x+c$ **and** $6=\text{'}their\ 9\text{'}(-1)+c\ \Rightarrow c=...$	m1		tangent using 'their' gradient, and attempt to find c using $x = –1$ and $y = 6$
	$\Rightarrow y=9x+15$	A1	5	equation must be seen in this form
(b)(i)	When $x=2$, $y=2^5-2\times2^2+9=32-8+9=33$ $(k=)$ 33	B1	1	be convinced that they are using **curve** equation **NMS** $k = 33$ scores B0
(ii)	When $x=2$, $y=9\times2+15=33$ so lies on tangent	B1	1	be convinced that they are using **tangent** equation **and** have statement

Q	Solution	Marks	Total	Comments
6(c)(i)	$\frac{x^6}{6}-\frac{2x^3}{3}+9x$	M1 A1 A1		one of these terms correct another term correct all correct (may have +c)
	$\left[\frac{2^6}{6}-\frac{2\times2^3}{3}+9\times2\right]-\left[\frac{(-1)^6}{6}-\frac{2\times(-1)^3}{3}+9\times(-1)\right]$	m1		F(2) – F(–1) unsimplified FT "their terms" from integration
	$\left[\frac{64}{6}-\frac{16}{3}+18\right]-\left[\frac{1}{6}+\frac{2}{3}-9\right]$ $= \frac{70}{3}-\left(-\frac{49}{6}\right)$			
	$= 31.5$ $\left(or\ \frac{189}{6}\ etc\right)$	A1	5	condone single fraction
(ii)	Area of trapezium = $\frac{1}{2}\times3\times(6+'their'k)$	B1√		= 58.5 when $k = 33$
	Shaded area = **Trapezium** – 'their' **(c)(i)** value	M1		
	$= 27$	A1	3	OE $eg\ \frac{162}{6}$
	Total		**15**	
7(a)	$(k-2)^2-4\times(2k-7)(k-3)$	M1		discriminant – condone one slip –condone omission of brackets
	$k^2-4k+4-4(2k^2-6k-7k+21)$	A1		
	"their" $-7k^2+48k-80\geqslant0$	B1		real roots condition ; f(k) $\geqslant$ 0 must appear before final line
	$7k^2-48k+80\leqslant0$	A1cso	4	**AG** (all working correct with no missing brackets etc)
(b)	$7k^2-48k+80=(7k-20)(k-4)$	M1		correct factors
				(or roots unsimplified) $\frac{48\pm\sqrt{64}}{14}$
	critical values are 4 and $\frac{20}{7}$	A1		accept $\frac{56}{14}$, $\frac{40}{14}$ etc here
		M1		sketch or sign diagram including values
				
	$\frac{20}{7}\leqslant k\leqslant4$	A1cao	4	fractions must be simplified here
	Take their final line as their answer			
	Total		**8**	
	TOTAL		**75**	