Version 1.0



**General Certificate of Education (A-level)**
**June 2013**

# Mathematics MPC4

# (Specification 6360)

# Pure Core 4

# Final

# *Mark Scheme*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of students' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2013 AQA and its licensors. All rights reserved.

**Copyright**
AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334).
Registered address: AQA, Devas Street, Manchester M15 6EX.

**Key to mark scheme abbreviations**

| | |
|---|---|
| M | mark is for method |
| m or dM | mark is dependent on one or more M marks and is for method |
| A | mark is dependent on M or m marks and is for accuracy |
| B | mark is independent of M or m marks and is for method and accuracy |
| E | mark is for explanation |
| √or ft or F | follow through from previous incorrect result |
| CAO | correct answer only |
| CSO | correct solution only |
| AWFW | anything which falls within |
| AWRT | anything which rounds to |
| ACF | any correct form |
| AG | answer given |
| SC | special case |
| OE | or equivalent |
| A2,1 | 2 or 1 (or 0) accuracy marks |
| –*x* EE | deduct *x* marks for each error |
| NMS | no method shown |
| PI | possibly implied |
| SCA | substantially correct approach |
| c | candidate |
| sf | significant figure(s) |
| dp | decimal place(s) |

**No Method Shown**

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| **1(a)(i)** | $5-8x=A(1-3x)+B(2+x)$<br>$x=-2 \qquad x=\frac{1}{3}$<br>$A=3 \qquad B=1$ | M1<br>m1<br>A1 | 3 | Two values of $x$ used to find values for $A$ and $B$ |
| **(ii)** | $\int_{-1}^{0}\frac{3}{2+x}+\frac{1}{1-3x}\,\mathrm{d}x$<br>$=3\ln(2+x)-\frac{1}{3}\ln(1-3x)$<br>$=\left(3\ln 2-\frac{1}{3}\ln 1\right)-\left(3\ln 1-\frac{1}{3}\ln 4\right)$<br>$=3\ln 2+\frac{1}{3}\ln 4$<br>$=\frac{11}{3}\ln 2$ | <br>M1<br>m1<br>A1ft<br>A1ft | <br><br><br><br>4 | <br>$a\ln(2+x)+b\ln(1-3x)$ where $a$ and $b$ are constants<br>$\mathrm{f}(0)-\mathrm{f}(-1)$ used<br>ft $A$ and $B$<br>ft $\left(A+\frac{2}{3}B\right)\ln 2$ |
| **(b)(i)** | $(C=)2$ | B1 | 1 | |
| **(ii)** | $\int\frac{9-18x-6x^2}{2-5x-3x^2}\,\mathrm{d}x=\int C\,\mathrm{d}x+\int\frac{5-8x}{2-5x-3x^2}\,\mathrm{d}x$<br>$\int_{-1}^{0}\frac{9-18x-6x^2}{2-5x-3x^2}\,\mathrm{d}x=2+\frac{11}{3}\ln 2$ | M1<br>A1ft | <br>2 | Seen or implied.<br>Allow $\pm C+\int\frac{5-8x}{2-5x-3x^2}\,\mathrm{d}x$<br>Accept $2+3\ln 2+\frac{1}{3}\ln 4$<br>ft $2+$ candidate's answer to part (a)(ii) if exact. |
| **(a)(i)** | **Alternative**<br>$5-8x=A(1-3x)+B(2+x)$<br>$5=A+2B$<br>$-8=-3A+B$<br>$A=3 \quad B=1$ | <br>(M1)<br>(m1)<br>(A1) | <br><br><br>(3) | <br><br>Set up simultaneous equations and solve. |
| | **Total** | | **10** | |

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| **2(a)(i)** | $h^2 = 2^2 + \sqrt{5}^2 = 9 \Rightarrow h = 3 \Rightarrow \sin\alpha = \frac{2}{3}$ | B1 | | Pythagoras used or all of $2,\ \sqrt{5},\ 3$ seen correctly on triangle |
| | $\cos\alpha = \frac{\sqrt{5}}{3}$ | B1 | 2 | **AG** $\frac{\sqrt{5}}{3}$ or $\sqrt{\frac{5}{9}}$ or $\frac{5}{3\sqrt{5}}$ seen |
| **(ii)** | $\sin 2\alpha = 2\sin\alpha\cos\alpha$ | M1 | | Correct formula seen or implied |
| | $= \left(2 \times \frac{2}{3} \times \frac{\sqrt{5}}{3}\right) = \frac{4}{9}\sqrt{5}$ | A1 | 2 | Must see $\frac{\sqrt{5}}{3}$ here or in part (a)(i) Accept $\frac{4}{3}\sqrt{\frac{5}{9}}$ |
| **(b)** | $\cos\beta = \frac{2}{\sqrt{5}}$ or $\sin\beta = \frac{1}{\sqrt{5}}$ | B1 | | Either correct. Accept $\sqrt{\frac{4}{5}}$ , $\frac{\sqrt{5}}{5}$ |
| | $\cos(\alpha - \beta) = \cos\alpha\cos\beta + \sin\alpha\sin\beta$ | M1 | | Correct formula seen or implied. |
| | $= \frac{\sqrt{5}}{3} \times \frac{2}{\sqrt{5}} + \frac{2}{3} \times \frac{1}{\sqrt{5}}$ | A1 | | All correct |
| | $= \frac{2}{15}\left(5 + \sqrt{5}\right)$ | A1 | 4 | $k = 5$ with previous A mark awarded |
| **(a)(i)** | **Alternative** | | | |
| | $\text{cosec}^2\alpha = 1 + \cot^2\alpha = 1 + \frac{5}{4} = \frac{9}{4}$ | | | |
| | $\text{cosec}\,\alpha = \frac{3}{2} \quad \sin\alpha = \frac{2}{3}$ | (B1) | | Must be positive |
| | $\sec^2\alpha = 1 + \tan^2\alpha = 1 + \frac{4}{5} = \frac{9}{5}$ | | | |
| | $\sec\alpha = \frac{3}{\sqrt{5}} \quad \cos\alpha = \frac{\sqrt{5}}{3}$ | (B1) | | Must be posiitve |
| | **Total** | | **8** | |

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| **3(a)** | $(1+6x)^{-\frac{1}{3}} = 1+\left(-\frac{1}{3}\right)6x+kx^2$ | M1 | | |
| | $=1-2x+8x^2$ | A1 | 2 | |
| **(b)(i)** | $(27+6x)^{-\frac{1}{3}} = 27^{-\frac{1}{3}}\left(1+\frac{6}{27}x\right)^{-\frac{1}{3}}$ | B1 | | |
| | $\left(1+\frac{6}{27}x\right)^{-\frac{1}{3}} = 1+\left(-\frac{1}{3}\times\frac{6}{27}x\right)+\left(-\frac{1}{3}\times-\frac{4}{3}\right)\frac{1}{2}\left(\frac{6}{27}x\right)^2$ | M1 | | Condone missing brackets and one error |
| | $(27+6x)^{-\frac{1}{3}} = \frac{1}{3}-\frac{2}{81}x+\frac{8}{2187}x^2$ | A1 | 3 | |
| **(ii)** | $\left(\sqrt[3]{\frac{2}{7}} = \frac{2}{\sqrt[3]{28}} \Rightarrow 27+6x=28 \Rightarrow x=\frac{1}{6}\right)$ | | | |
| | $\sqrt[3]{\frac{1}{28}} = \frac{1}{3}-\frac{2}{81}\times\frac{1}{6}+\frac{8}{2187}\times\left(\frac{1}{6}\right)^2 \left(\approx 0.3293..\right)$ | M1 | | Substitute $x=\frac{1}{6}$ into expansion from (b)(i) |
| | $\left(\sqrt[3]{\frac{2}{7}} \approx 2\times 0.3293197.. = 0.6586394...\right)$ | | | |
| | $= 0.658639 \quad (6\text{dp})$ | A1 | 2 | CSO |
| | **Alternatives** | | | |
| **(b)(i)** | $(27+6x)^{-\frac{1}{3}} = 27^{-\frac{1}{3}}\left(1+\frac{6}{27}x\right)^{-\frac{1}{3}}$ | (B1) | | |
| | $\left(1+\frac{6}{27}x\right)^{-\frac{1}{3}} = 1-2\times\frac{1}{27}x+8\times\left(\frac{1}{27}\right)^2x^2$ | (M1) | | Replace $x$ with $\frac{1}{27}x$, **not** $\frac{6}{27}x$, in expansion from (a); condone missing brackets and one error |
| | $(27+6x)^{-\frac{1}{3}} = \frac{1}{3}-\frac{2}{81}x+\frac{8}{2187}x^2$ | (A1) | (3) | |
| **(b)(i)** | $(27+6x)^{-\frac{1}{3}} = 27^{-\frac{1}{3}}+\left(-\frac{1}{3}\right)27^{-\frac{4}{3}}\times 6x$ | (M1) | | Use result from formula book; Condone missing brackets and one error |
| | $+\left(-\frac{1}{3}\right)\times\left(-\frac{4}{3}\right)\frac{1}{2}27^{-\frac{7}{3}}\times(6x)^2$ | | | |
| | $(27+6x)^{-\frac{1}{3}} = \frac{1}{3}-\frac{2}{81}x+\frac{8}{2187}x^2$ | (A2) | (3) | A1 not available |
| | **Total** | | **7** | |

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| **4(a)** | $\left(\frac{dx}{dt}=\right)-16e^{-2t}$ $\left(\frac{dy}{dt}=\right)4e^{2t}$ | B1 | | Both derivatives correct |
| | $\frac{dy}{dx}=\frac{\text{candidate's }\frac{dy}{dt}}{\text{candidate's }\frac{dx}{dt}}$ | M1 | | chain rule used correctly |
| | $\frac{dy}{dx}=\frac{4e^{2t}}{-16e^{-2t}}$ $\left(=-\frac{1}{4}e^{4t}\right)$ | A1 | 3 | Simplification not required<br>$4e^{2t}$ and $-16e^{-2t}$ must be seen.<br>ISW. |
| **(b)** | | | | |
| **(i)** | $t=\ln 2$ gradient at $P=-4$ | B1ft | 1 | B0 if ISW result is used. |
| **(ii)** | coordinates of $P$ $x=-2$ | B1 | | |
| | $y=12$ | B1 | 2 | |
| **(iii)** | gradient of normal $=\frac{1}{4}$ | B1ft | | ft gradient at $P$ |
| | equation of normal $\frac{y-12}{x--2}=\frac{1}{4}$ | M1 | | Set up equation of normal |
| | at $y=0$ $x=-50$ | A1 | 3 | $(-50,0)$ CSO |
| **(c)** | $xy+4y-4x=(8e^{-2t}-4)(2e^{2t}+4)$ | | | |
| | $+4(2e^{2t}+4)-4(8e^{-2t}-4)$ | M1 | | Write $xy+4y-4x$ in terms of $t$. |
| | $=16+32e^{-2t}-8e^{2t}-16$ | | | |
| | $+8e^{2t}+16-32e^{-2t}+16$ | m1 | | Multiply out and simplify using $e^{-2t}e^{2t}=1$ PI |
| | $(xy+4y-4x)=32$ | A1 | 3 | Correct working to $k=32$<br>$k=32$ NMS; SC1 |
| **(c)** | **Alternative** | | | |
| | $e^{-2t}=\frac{x+4}{8}$ or $e^{2t}=\frac{y-4}{2}$ | (M1) | | Write $e^{-2t}$ in terms of $x$ or $e^{2t}$ in terms of $y$ . Condone sign errors |
| | $e^{-2t}e^{2t}=\left(\frac{x+4}{8}\right)\left(\frac{y-4}{2}\right)$ | | | |
| | $=\frac{xy+4y-4x-16}{16}=1$ | (m1) | | Multiply out and use $e^{-2t}e^{2t}=1$ |
| | $xy+4y-4x=32$ | (A1) | (3) | All correct with $k=32$ |
| | Other alternatives are possible | | | |
| | **Total** | | **12** | |

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| **5(a)** | $f\left(-\frac{3}{2}\right)=4\left(-\frac{3}{2}\right)^3-11\left(-\frac{3}{2}\right)-3$<br>$=-4\times\frac{27}{8}+\frac{33}{2}-3=0\Rightarrow$ factor | M1<br>A1 | 2 | $x=-\frac{3}{2}$ substituted<br>Processing, $=0$ and conclusion |
| **(b)** | $2x^2-3x-1$ | M1A1 | 2 | M1 for any two of $a,b,c$ correct |
| **(c)(i)** | $2\cos 2\theta\sin\theta+9\sin\theta+3$<br>$=2\left(1-2\sin^2\theta\right)\sin\theta+9\sin\theta+3$<br>$=2\sin\theta-4\sin^3\theta+9\sin\theta+3$<br>$\sin\theta=x\Rightarrow 4x^3-11x-3=0$ | M1<br>m1<br>A1 | 3 | $\cos 2\theta$ expanded ; ACF and substituted<br>All in terms of $\sin\theta$ or $x$ and simplified to a cubic expression.<br>Reverse signs and express in $x$ correctly **AG** |
| **(c)(ii)** | $2x^2-3x-1=0\Rightarrow x=\frac{3\pm\sqrt{17}}{4}$<br>$x=\frac{3-\sqrt{17}}{4}$ or $-0.28...$<br>$\theta=196°$ and $344°$<br>$x=\frac{3+\sqrt{17}}{4}$ no solutions for $\sin\theta$<br>$x=-\frac{3}{2}$ no solutions for $\sin\theta$ | M1<br>A1<br>A1<br><br>E1 | 4 | Use formula correctly to solve $ax^2+bx+c=0$ from part (b)<br><br>Both required and no others in range; condone greater accuracy<br>Ignore solutions out of range.<br>Must have three correct roots and reject both other roots from cubic equation. |
| | **Total** | | **11** | |

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| **6(a)** | | | | |
| | $\lambda = -1$ | B1 | | $\lambda = -1$ seen or implied |
| | $\lambda = -1$ verified in all three components | B1 | 2 | Shown |
| **(b)** | $\pm\begin{bmatrix}-2\\-3\\2\end{bmatrix}$ | B1 | | $\overrightarrow{AB}$ or $\overrightarrow{BA}$ correct |
| | $\mathbf{r} = \overrightarrow{OA} + \mu\overrightarrow{AB} = \begin{bmatrix}3\\-2\\4\end{bmatrix} + \mu\begin{bmatrix}-2\\-3\\2\end{bmatrix}$ | M1 | | $\mathbf{a} + \mu\mathbf{d}$ |
| | | A1ft | 3 | OE; ft on $\overrightarrow{AB}$ or $\overrightarrow{BA}$ |
| **(c)** | | | | |
| | $\overrightarrow{CD} = \overrightarrow{OD} - \overrightarrow{OC}$ | | | |
| | $= \begin{bmatrix}3-2\mu\\-2-3\mu\\4+2\mu\end{bmatrix} - \begin{bmatrix}-4\\5\\-1\end{bmatrix} \quad \left(= \begin{bmatrix}7-2\mu\\-7-3\mu\\5+2\mu\end{bmatrix}\right)$ | B1 | | $\pm\overrightarrow{CD}$ in terms of $\mu$<br>OE |
| | $\overrightarrow{CD}.\overrightarrow{AB} = 0$ or $\overrightarrow{CD}.\overrightarrow{AD} = 0$ | | | |
| | $= \left(\begin{bmatrix}3-2\mu\\-2-3\mu\\4+2\mu\end{bmatrix} - \begin{bmatrix}-4\\5\\-1\end{bmatrix}\right) . \begin{bmatrix}-2\\-3\\2\end{bmatrix} = 0$ | M1 | | Candidate's $\overrightarrow{CD}$ sp with candidate's $\overrightarrow{AB}$ or $\overrightarrow{AD}$<br>$= 0$ PI by a solution for $\mu$ |
| | $-14 + 4\mu + 21 + 9\mu + 10 + 4\mu = 0$ | | | |
| | $17 + 17\mu = 0$ | | | |
| | $\mu = -1$ | m1A1 | | Expand sp to an equation in $\mu$ and solve for $\mu$ |
| | $D$ is at $(5,1,2)$ | A1 | 5 | Accept as a column vector |
| **(d)** | $\overrightarrow{OE} = \overrightarrow{OA} + \overrightarrow{AE} = \overrightarrow{OA} + 3\overrightarrow{AD}$ | M1 | | Accept $AE = 3AD$ |
| | $\overrightarrow{OE} = \begin{bmatrix}3\\-2\\4\end{bmatrix} + 3\begin{bmatrix}2\\3\\-2\end{bmatrix}$ $E$ is at $(9,7,-2)$ | A1 | | Accept as a column vector |
| | Or | | | |
| | $\overrightarrow{OE} = \overrightarrow{OA} + \overrightarrow{AE} = \overrightarrow{OA} + 3\overrightarrow{DA}$ | M1 | | Accept $AE = 3DA$ |
| | $\overrightarrow{OE} = \begin{bmatrix}3\\-2\\4\end{bmatrix} + 3\begin{bmatrix}-2\\-3\\2\end{bmatrix}$ $E$ is at $(-3,-11,10)$ | A1 | 4 | Accept as a column vector. |

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| 6(c) | **Alternative using Pythagoras**<br>$\overrightarrow{CD} = \overrightarrow{OD} - \mu\overrightarrow{OC}$<br>$= \begin{bmatrix} 3-2\mu \\ -2-3\mu \\ 4+2\mu \end{bmatrix} - \begin{bmatrix} -4 \\ 5 \\ -1 \end{bmatrix} \quad \left( = \begin{bmatrix} 7-2\mu \\ -7-3\mu \\ 5+2\mu \end{bmatrix} \right)$ | (B1) | | $\pm\overrightarrow{CD}$ in terms of $\mu$ |
| | $AC^2 = AD^2 + CD^2$<br>$\left(7^2+7^2+5^2\right) = \mu^2\left(2^2+3^2+2^2\right)$<br>$+\left((7-2\mu)^2+(7+3\mu)^2+(5+2\mu)^2\right)$ | (M1) | | $\overrightarrow{AC} = \begin{bmatrix} -7 \\ 7 \\ -5 \end{bmatrix} \quad \overrightarrow{AD} = \begin{bmatrix} -2\mu \\ -3\mu \\ 2\mu \end{bmatrix}$<br>Correct Pythagoras expression in terms of $\mu$; |
| | $123 = 17\mu^2 + 123 + 34\mu + 17\mu^2$<br>$0 = 34\mu^2 + 34\mu$ | (m1) | | Multiply out and solve to find a value for $\mu$ |
| | $\mu = -1 \quad (\mu = 0$ is point $A)$ | (A1) | | $\mu = -1$ |
| | $D$ is at $(5,1,2)$ | (A1) | (5) | |
| 6(d) | **Alternative**<br>$\left\|\overrightarrow{DE}\right\| = 2\left\|\overrightarrow{AD}\right\| \Rightarrow \overrightarrow{OE} = \overrightarrow{OD} + 2\overrightarrow{AD}$ | (M1) | | |
| | $\overrightarrow{OE} = \begin{bmatrix} 5 \\ 1 \\ 2 \end{bmatrix} + 2\begin{bmatrix} 2 \\ 3 \\ -2 \end{bmatrix} \quad E$ is at $(9,7,-2)$ | (A1) | | |
| | $\left\|DE\right\| = 4\left\|DA\right\| \Rightarrow \overrightarrow{OE} = \overrightarrow{OD} + 4\overrightarrow{DA}$ | (M1) | | |
| | $\overrightarrow{OE} = \begin{bmatrix} 5 \\ 1 \\ 2 \end{bmatrix} + 4\begin{bmatrix} -2 \\ -3 \\ 2 \end{bmatrix} \quad E$ is at $(-3,-11,10)$ | (A1) | (4) | |
| | **Total** | | **14** | |

| Q | Solution | Marks | Total | Comments |
|---|---|---|---|---|
| **7** | $\frac{\mathrm{d}h}{\mathrm{d}t}$ | B1 | 1 | $\frac{\mathrm{d}h}{\mathrm{d}t}$ seen |
| | $a = 1.3$ or $a = -1.3$ | B1 | 1 | |
| | $k = \frac{\pi}{6}$ or $k = \frac{2\pi}{12}$ | B1 | 1 | |
| | **Total** | | **3** | |
| **8 (a)** | $\int t\cos\left(\frac{\pi}{4}t\right)\mathrm{d}t$ | M1 | | Clear attempt to use parts<br>$u = t$ $\quad \frac{\mathrm{d}v}{\mathrm{d}t} = \cos\left(\frac{\pi}{4}t\right)$<br>$\frac{\mathrm{d}u}{\mathrm{d}t} = 1$ $\quad v = k\sin\left(\frac{\pi}{4}t\right)$ |
| | $= t \times \frac{4}{\pi}\sin\left(\frac{\pi}{4}t\right) - \frac{4}{\pi}\int \sin\left(\frac{\pi}{4}t\right)(\mathrm{d}t)$ | A1 | | Must be in terms of $\pi$ |
| | $= pt\sin\left(\frac{\pi}{4}t\right) + q\cos\left(\frac{\pi}{4}t\right)$ | m1 | | Correct form, any non-zero values for $p$, $q$ |
| | $= t \times \frac{4}{\pi}\sin\left(\frac{\pi}{4}t\right) + \frac{4}{\pi} \times \frac{4}{\pi}\cos\left(\frac{\pi}{4}t\right)$ | A1 | 4 | Any correct unsimplified form.<br>Constant not required |
| **(b)** | | | | |
| | $\int 32x\ \mathrm{d}x = \int t\cos\left(\frac{\pi}{4}t\right)\mathrm{d}t$ | B1 | | Correct separation and notation. |
| | $16x^2 =$ | B1 | | $\frac{x^2}{2}$ if 32 not brought over; allow $32 \times \frac{x^2}{2}$ |
| | $t \times \frac{4}{\pi}\sin\left(\frac{\pi}{4}t\right) + \frac{16}{\pi^2}\cos\left(\frac{\pi}{4}t\right) + C$ | M1 | | Equate to result from part (a) with constant and use $(0,4)$ to find a value for the constant |
| | $C = 256 - \frac{16}{\pi^2}$ | A1 | | Accept $C = 254$ or better $(254.37886...)$ |
| | $t = 45$ | | | Substitute $t = 45$ into $kx^2 = pt\sin\left(\frac{\pi}{4}t\right) + q\cos\left(\frac{\pi}{4}t\right) + C$<br>$p \neq 0$ , $q \neq 0$ |
| | $16x^2 = -40.514... - 1.146... + 254.378...$<br>$= 212.718...$<br>$x^2 = 13.294...$<br>$x = 3.646... = 3.65$ m<br>or (height =) 365 cm | m1A1 | 6 | and calculate $x$.<br>CSO |
| | **Total** | | **10** | |
| | **TOTAL** | | **75** | |