Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	9	/	0	1	Signature	

Paper Reference(s)

6679/01

Edexcel GCE

Mechanics M3

Advanced/Advanced Subsidiary

Wednesday 13 May 2015 – Morning

Time: 1 hour 30 minutes

Examiner's use only

Team Leader's use only

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
Total	

Materials required for examination
Mathematical Formulae (Pink)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer to each question in the space following the question.
Whenever a numerical value of g is required, take $g = 9.8$ m s^{-2}, and give your answer to either two significant figures or three significant figures.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 7 questions in this question paper. The total mark for this paper is 75.
There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2015 Pearson Education Ltd.

Printer's Log. No.
P44514A

5/1/

P 4 4 5 1 4 A 0 1 2 8

Turn over

PEARSON

Leave blank

1. A particle P of mass 0.5 kg is attached to one end of a light elastic spring, of natural length 1.2 m and modulus of elasticity λ newtons. The other end of the spring is attached to a fixed point A on a ceiling. The particle is hanging freely in equilibrium at a distance 1.5 m vertically below A.

(a) Find the value of λ.

(3)

The particle is now raised to the point B, where B is vertically below A and $AB = 0.8$ m. The spring remains straight. The particle is released from rest and first comes to instantaneous rest at the point C.

(b) Find the distance AC.

(4)

Leave blank

Question 1 continued

Q1

(Total 7 marks)

Leave blank

2. The finite region bounded by the x-axis, the curve with equation $y = 2e^x$, the y-axis and the line $x = 1$ is rotated through one complete revolution about the x-axis to form a uniform solid.

Use algebraic integration to

(a) show that the volume of the solid is $2\pi(e^2 - 1)$,

(4)

(b) find, in terms of e, the x coordinate of the centre of mass of the solid.

(6)

Leave blank

Question 2 continued

Leave blank

Question 2 continued

Leave blank

Question 2 continued

Q2

(Total 10 marks)

Leave blank

3.




Figure 1

A small ball P of mass m is attached to the midpoint of a light inextensible string of length $4l$. The ends of the string are attached to fixed points A and B, where A is vertically above B. Both strings are taut and AP makes an angle of 30° with AB, as shown in Figure 1. The ball is moving in a horizontal circle with constant angular speed ω.

(a) Find, in terms of m, g, l and ω,

(i) the tension in AP,

(ii) the tension in BP.

(8)

(b) Show that $\omega^2 \geqslant \dfrac{g\sqrt{3}}{3l}$.

(2)

Leave blank

Question 3 continued

Leave blank

Question 3 continued

Leave blank

Question 3 continued

Q3

(Total 10 marks)

Leave blank

4. A vehicle of mass 900 kg moves along a straight horizontal road. At time t seconds the resultant force acting on the vehicle has magnitude $\frac{63000}{kt^2}$ N, where k is a positive constant. The force acts in the direction of motion of the vehicle. At time t seconds, $t \geqslant 1$, the speed of the vehicle is v m s^{-1} and the vehicle is a distance x metres from a fixed point O on the road. When $t = 1$ the vehicle is at rest at O and when $t = 4$ the speed of the vehicle is 10.5 m s^{-1}.

(a) Show that $v = 14 - \frac{14}{t}$

(7)

(b) Hence deduce that the speed of the vehicle never reaches 14 m s^{-1}.

(1)

(c) Use the trapezium rule, with 4 equal intervals, to estimate the value of x when $v = 7$

(4)

Leave blank

Question 4 continued

Leave blank

Question 4 continued

Leave blank

Question 4 continued

Q4

(Total 12 marks)

Leave blank

5.




Figure 2

Figure 2 shows a uniform solid spindle which is made by joining together the circular faces of two right circular cones. The common circular face has radius r and centre O. The smaller cone has height h and the larger cone has height kh. The point A lies on the circumference of the common circular face. The spindle is suspended from A and hangs freely in equilibrium with AO at an angle of 30° to the vertical.

Show that $k = \dfrac{4r}{h\sqrt{3}} + 1$

(6)

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Q5

(Total 6 marks)

Leave blank

6.




Figure 3

Two points A and B are 6 m apart on a smooth horizontal floor. A particle P of mass 0.5 kg is attached to one end of a light elastic spring, of natural length 2.5 m and modulus of elasticity 20 N. The other end of the spring is attached to A. A second light elastic spring, of natural length 1.5 m and modulus of elasticity 18 N, has one end attached to P and the other end attached to B, as shown in Figure 3. Initially P rests in equilibrium at the point O, where AOB is a straight line.

(a) Find the length of AO.

(4)

The particle P now receives an impulse of magnitude 6 N s acting in the direction OB and P starts to move towards B.

(b) Show that P moves with simple harmonic motion about O.

(4)

(c) Find the amplitude of the motion.

(4)

(d) Find the time taken by P to travel 1.2 m from O.

(3)

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Q6

(Total 15 marks)

Leave blank

7. A solid smooth sphere, with centre O and radius r, is fixed to a point A on a horizontal floor. A particle P is placed on the surface of the sphere at the point B, where B is vertically above A. The particle is projected horizontally from B with speed $\frac{\sqrt{gr}}{2}$ and starts to move on the surface of the sphere. When OP makes an angle θ with the upward vertical and P remains in contact with the sphere, the speed of P is v.

(a) Show that $v^2 = \frac{gr}{4}(9 - 8\cos\theta)$.

(4)

The particle leaves the surface of the sphere when $\theta = \alpha$.

(b) Find the value of $\cos\alpha$.

(4)

After leaving the surface of the sphere, P moves freely under gravity and hits the floor at the point C.

Given that $r = 0.5\,\text{m}$,

(c) find, to 2 significant figures, the distance AC.

(7)

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Q7

(Total 15 marks)

TOTAL FOR PAPER: 75 MARKS

END