Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	5	/	0	1	Signature	

Paper Reference(s)

6665/01

Edexcel GCE

Core Mathematics C3

Advanced

Monday 16 June 2014 – Morning

Time: 1 hour 30 minutes

Examiner's use only		

Team Leader's use only		

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
9	
Total	

Materials required for examination
Mathematical Formulae (Pink)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer for each question in the space following the question.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 9 questions in this question paper. The total mark for this paper is 75.
There are 32 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2014 Pearson Education Ltd.

Printer's Log. No.
P43164A
W850/R6665/57570 5/5/5/1/

P 4 3 1 6 4 A 0 1 3 2

Turn over

PEARSON

Leave blank

1. The curve C has equation $y = \mathrm{f}(x)$ where

$$\mathrm{f}(x) = \frac{4x+1}{x-2}, \quad x > 2$$

(a) Show that

$$\mathrm{f}'(x) = \frac{-9}{(x-2)^2}$$

(3)

Given that P is a point on C such that $\mathrm{f}'(x) = -1$,

(b) find the coordinates of P.

(3)

Leave blank

Question 1 continued

Q1

(Total 6 marks)

Leave blank

2. Find the exact solutions, in their simplest form, to the equations

(a) $2 \ln (2x + 1) - 10 = 0$

(2)

(b) $3^x e^{4x} = e^7$

(4)

Leave blank

Question 2 continued

Q2

(Total 6 marks)

Leave blank

3. The curve C has equation $x = 8y \tan 2y$

The point P has coordinates $\left(\pi, \frac{\pi}{8}\right)$

(a) Verify that P lies on C.

(1)

(b) Find the equation of the tangent to C at P in the form $ay = x + b$, where the constants a and b are to be found in terms of π.

(7)

Leave blank

Question 3 continued

Q3

(Total 8 marks)

Leave blank

4.




Figure 1

Figure 1 shows part of the graph with equation $y = \mathrm{f}(x)$, $x \in \mathbb{R}$.

The graph consists of two line segments that meet at the point $Q(6, -1)$.

The graph crosses the y-axis at the point $P(0, 11)$.

Sketch, on separate diagrams, the graphs of

(a) $y = |\mathrm{f}(x)|$

(2)

(b) $y = 2\mathrm{f}(-x) + 3$

(3)

On each diagram, show the coordinates of the points corresponding to P and Q.

Given that $\mathrm{f}(x) = a|x - b| - 1$, where a and b are constants,

(c) state the value of a and the value of b.

(2)

Leave blank

Question 4 continued

Leave blank

Question 4 continued

Leave blank

Question 4 continued

Q4

(Total 7 marks)

Leave blank

5.

$$g(x) = \frac{x}{x+3} + \frac{3(2x+1)}{x^2+x-6}, \quad x > 3$$

(a) Show that $g(x) = \frac{x+1}{x-2}, \quad x > 3$

(4)

(b) Find the range of g.

(2)

(c) Find the exact value of a for which $g(a) = g^{-1}(a)$.

(4)

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Q5

(Total 10 marks)

Leave blank

6.




Figure 2

Figure 2 shows a sketch of part of the curve with equation

$$y = 2\cos\left(\frac{1}{2}x^2\right) + x^3 - 3x - 2$$

The curve crosses the x-axis at the point Q and has a minimum turning point at R.

(a) Show that the x coordinate of Q lies between 2.1 and 2.2

(2)

(b) Show that the x coordinate of R is a solution of the equation

$$x = \sqrt{1 + \frac{2}{3}x\sin\left(\frac{1}{2}x^2\right)}$$

(4)

Using the iterative formula

$$x_{n+1} = \sqrt{1 + \frac{2}{3}x_n \sin\left(\frac{1}{2}x_n^{\,2}\right)}, \quad x_0 = 1.3$$

(c) find the values of x_1 and x_2 to 3 decimal places.

(2)

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Q6

(Total 8 marks)

Leave blank

7. (a) Show that

$$\text{cosec } 2x + \cot 2x = \cot x, \qquad x \neq 90n°, \qquad n \in \mathbb{Z}$$

(5)

(b) Hence, or otherwise, solve, for $0 \leqslant \theta < 180°$,

$$\text{cosec}\,(4\theta + 10°) + \cot\,(4\theta + 10°) = \sqrt{3}$$

You must show your working.

(*Solutions based entirely on graphical or numerical methods are not acceptable.*)

(5)

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Q7

(Total 10 marks)

Leave blank

8. A rare species of primrose is being studied. The population, P, of primroses at time t years after the study started is modelled by the equation

$$P = \frac{800e^{0.1t}}{1 + 3e^{0.1t}}, \qquad t \geqslant 0, \quad t \in \mathbb{R}$$

(a) Calculate the number of primroses at the start of the study.

(2)

(b) Find the exact value of t when $P = 250$, giving your answer in the form $a \ln(b)$ where a and b are integers.

(4)

(c) Find the exact value of $\frac{dP}{dt}$ when $t = 10$. Give your answer in its simplest form.

(4)

(d) Explain why the population of primroses can never be 270

(1)

Leave blank

Question 8 continued

Leave blank

Question 8 continued

Leave blank

Question 8 continued

Q8

(Total 11 marks)

Leave blank

9. (a) Express $2\sin\theta - 4\cos\theta$ in the form $R\sin(\theta - \alpha)$, where R and α are constants, $R > 0$ and $0 < \alpha < \frac{\pi}{2}$

Give the value of α to 3 decimal places.

(3)

$$H(\theta) = 4 + 5(2\sin 3\theta - 4\cos 3\theta)^2$$

Find

(b) (i) the maximum value of $H(\theta)$,

(ii) the smallest value of θ, for $0 \leqslant \theta < \pi$, at which this maximum value occurs.

(3)

Find

(c) (i) the minimum value of $H(\theta)$,

(ii) the largest value of θ, for $0 \leqslant \theta < \pi$, at which this minimum value occurs.

(3)

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Q9

(Total 9 marks)

TOTAL FOR PAPER: 75 MARKS

END