Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	8	0	/	0	1	Signature	

Paper Reference(s)

6680/01

Edexcel GCE

Mechanics M4

Advanced/Advanced Subsidiary

Monday 16 June 2014 – Morning

Time: 1 hour 30 minutes

Examiner's use only

Team Leader's use only

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
Total	

Materials required for examination
Mathematical Formulae (Pink)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer to each question in the space following the question.
Whenever a numerical value of g is required, take $g = 9.8$ m s^{-2} and give your answer to either two significant figures or three significant figures.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 7 questions in this question paper. The total mark for this paper is 75.
There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2014 Pearson Education Ltd.

Printer's Log. No.
P43159A
W850/R6680/57570 5/5/5/1/1/

P 4 3 1 5 9 A 0 1 2 4

Turn over

PEARSON

Leave blank

1. A particle A has constant velocity $(3\mathbf{i} + \mathbf{j})$ m s^{-1} and a particle B has constant velocity $(\mathbf{i} - \mathbf{k})$ m s^{-1}. At time $t = 0$ seconds, the position vectors of the particles A and B with respect to a fixed origin O are $(-6\mathbf{i} + 4\mathbf{j} - 3\mathbf{k})$ m and $(-2\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$ m respectively.

(a) Show that, in the subsequent motion, the minimum distance between A and B is $4\sqrt{2}$ m.

(6)

(b) Find the position vector of A at the instant when the distance between A and B is a minimum.

(2)

Leave blank

Question 1 continued

Q1

(Total 8 marks)

Leave blank

2. A car of mass 1000 kg is moving along a straight horizontal road. The engine of the car is working at a constant rate of 25 kW. When the speed of the car is v m s^{-1}, the resistance to motion has magnitude $10v$ newtons.

(a) Show that, at the instant when $v = 20$, the acceleration of the car is 1.05 m s^{-2}.

(3)

(b) Find the distance travelled by the car as it accelerates from a speed of 10 m s^{-1} to a speed of 20 m s^{-1}.

(8)

Leave blank

Question 2 continued

Q2

(Total 11 marks)

Leave blank

3. A small ball is moving on a smooth horizontal plane when it collides obliquely with a smooth plane vertical wall. The coefficient of restitution between the ball and the wall is $\frac{1}{3}$. The speed of the ball immediately after the collision is half the speed of the ball immediately before the collision.

Find the angle through which the path of the ball is deflected by the collision.

(8)

Leave blank

Question 3 continued

Q3

(Total 8 marks)

Leave blank

4. At noon two ships A and B are 20 km apart with A on a bearing of 230° from B. Ship B is moving at 6 km h^{-1} on a bearing of 015°. The maximum speed of A is 12 km h^{-1}. Ship A sets a course to intercept B as soon as possible.

(a) Find the course set by A, giving your answer as a bearing to the nearest degree.

(4)

(b) Find the time at which A intercepts B.

(4)

Leave blank

Question 4 continued

Leave blank

Question 4 continued

Leave blank

Question 4 continued

Q4

(Total 8 marks)

Leave blank

5.




Figure 1

Two smooth uniform spheres A and B have equal radii. The mass of A is m and the mass of B is $3m$. The spheres are moving on a smooth horizontal plane when they collide obliquely. Immediately before the collision, A is moving with speed $3u$ at angle α to the line of centres and B is moving with speed u at angle β to the line of centres, as shown in Figure 1. The coefficient of restitution between the two spheres is $\frac{1}{5}$. It is given that $\cos\alpha = \frac{1}{3}$ and $\cos\beta = \frac{2}{3}$ and that α and β are both acute angles.

(a) Find the magnitude of the impulse on A due to the collision in terms of m and u.

(8)

(b) Express the kinetic energy lost by A in the collision as a fraction of its initial kinetic energy.

(4)

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Q5

(Total 12 marks)

Leave blank

6. A particle of mass m kg is attached to one end of a light elastic string of natural length a metres and modulus of elasticity $5ma$ newtons. The other end of the string is attached to a fixed point O on a smooth horizontal plane. The particle is held at rest on the plane with the string stretched to a length $2a$ metres and then released at time $t = 0$. During the subsequent motion, when the particle is moving with speed v m s^{-1}, the particle experiences a resistance of magnitude $4mv$ newtons. At time t seconds after the particle is released, the length of the string is $(a + x)$ metres, where $0 \leqslant x \leqslant a$.

(a) Show that, from $t = 0$ until the string becomes slack,

$$\frac{\mathrm{d}^2x}{\mathrm{d}t^2} + 4\frac{\mathrm{d}x}{\mathrm{d}t} + 5x = 0$$

(3)

(b) Hence express x in terms of a and t.

(6)

(c) Find the speed of the particle at the instant when the string first becomes slack, giving your answer in the form ka, where k is a constant to be found correct to 2 significant figures.

(4)

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Q6

(Total 13 marks)

Leave blank

7.




Figure 2

A bead B of mass m is threaded on a smooth circular wire of radius r, which is fixed in a vertical plane. The centre of the circle is O, and the highest point of the circle is A. A light elastic string of natural length r and modulus of elasticity kmg has one end attached to the bead and the other end attached to A. The angle between the string and the downward vertical is θ, and the extension in the string is x, as shown in Figure 2.

Given that the string is taut,

(a) show that the potential energy of the system is

$$2mgr\{(k-1)\cos^2\theta - k\cos\theta\} + \text{constant}$$

(6)

Given also that $k = 3$,

(b) find the positions of equilibrium and determine their stability.

(9)

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Q7

(Total 15 marks)

TOTAL FOR PAPER: 75 MARKS

END