# GCE

# Mathematics A

## H240/01: Pure Mathematics

Advanced GCE

# Mark Scheme for Autumn 2021

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2021

## Text Instructions

### 1. Annotations and abbreviations

| Annotation in RM assessor | Meaning |
|---|---|
| ✓and ✗ | |
| BOD | Benefit of doubt |
| FT | Follow through |
| ISW | Ignore subsequent working |
| M0, M1 | Method mark awarded 0, 1 |
| A0, A1 | Accuracy mark awarded 0, 1 |
| B0, B1 | Independent mark awarded 0, 1 |
| SC | Special case |
| ^ | Omission sign |
| MR | Misread |
| BP | Blank Page |
| Seen | |
| Highlighting | |
| | |
| **Other abbreviations in mark scheme** | **Meaning** |
| dep* | Mark dependent on a previous mark, indicated by *. The * may be omitted if only one previous M mark |
| cao | Correct answer only |
| oe | Or equivalent |
| rot | Rounded or truncated |
| soi | Seen or implied |
| www | Without wrong working |
| AG | Answer given |
| awrt | Anything which rounds to |
| BC | By Calculator |
| DR | This question included the instruction: In this question you must show detailed reasoning. |

## 2. Subject-specific Marking Instructions for A Level Mathematics A

a Annotations must be used during your marking. For a response awarded zero (or full) marks a single appropriate annotation (cross, tick, M0 or ^) is sufficient, but not required.

For responses that are not awarded either 0 or full marks, you must make it clear how you have arrived at the mark you have awarded and all responses must have enough annotation for a reviewer to decide if the mark awarded is correct without having to mark it independently.

It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

Award NR (No Response)

- if there is nothing written at all in the answer space and no attempt elsewhere in the script
- OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
- OR if there is a mark (e.g. a dash, a question mark, a picture) which isn't an attempt at the question.

Note: Award 0 marks only for an attempt that earns no credit (including copying out the question).

If a candidate uses the answer space for one question to answer another, for example using the space for 8(b) to answer 8(a), then give benefit of doubt unless it is ambiguous for which part it is intended.

b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not always be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.
If you are in any doubt whatsoever you should contact your Team Leader.

c The following types of marks are available.

**M**
A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.
A method mark may usually be implied by a correct answer unless the question includes the DR statement, the command words "Determine" or "Show that", or some other indication that the method must be given explicitly.

**A**
Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

**B**
Mark for a correct result or statement independent of Method marks.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep*' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.

e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case please, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.
Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

f We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so.

- When a value **is given** in the paper only accept an answer correct to at least as many significant figures as the given value.

- When a value **is not given** in the paper accept any answer that agrees with the correct value to **3 s.f.** unless a different level of accuracy has been asked for in the question, or the mark scheme specifies an acceptable range.
  NB for Specification B (MEI) the rubric is not specific about the level of accuracy required, so this statement reads "2 s.f".

Follow through should be used so that only one mark in any question is lost for each distinct accuracy error.
Candidates using a value of 9.80, 9.81 or 10 for *g* should usually be penalised for any final accuracy marks which do not agree to the value found with 9.8 which is given in the rubric.

g Rules for replaced work and multiple attempts:

- If one attempt is clearly indicated as the one to mark, or only one is left uncrossed out, then mark that attempt and ignore the others.
- If more than one attempt is left not crossed out, then mark the last attempt unless it only repeats part of the first attempt or is substantially less complete.
- if a candidate crosses out all of their attempts, the assessor should attempt to mark the crossed out answer(s) as above and award marks appropriately.

h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question. Marks designated as cao may be awarded as long as there are no other errors.
If a candidate corrects the misread in a later part, do not continue to follow through. Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

i If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers, provided that there is nothing in the wording of the question specifying that analytical methods are required such as the bold "In this question you must show detailed reasoning", or the command words "Show" or "Determine". Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.

j If in any case the scheme operates with considerable unfairness consult your Team Leader.

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| 1 | | | $16 - 4(k + 3)$<br>$-4k - 12 + 16 > 0$<br>$4k - 4 < 0$<br><br>$k < 1$ | **M1***<br>**A1**<br>**M1dep***<br><br>**A1**<br>**[4]** | **1.1**<br>**2.3**<br>**1.1**<br><br>**1.1** | Attempt discriminant<br>Obtain correct inequality<br>Attempt to solve their inequality or equation for $k$<br><br>Obtain $k < 1$ | Allow $b^2 + 4ac$ for M1, but nothing else<br>Not necessarily expanded<br><br><br><br>**OR** (completing the square or differentiating)<br>**M1*** – attempt to complete the square, or differentiate, and link minimum point to 0<br>**A1** – obtain $(k + 3) - 4 < 0$<br>**M1d*** – solve their inequality or equation<br>**A1** – obtain $k < 1$<br><br>**OR** (using perfect square)<br>**M1*** – link $k + 3$ to 4<br>**A1** – obtain $k + 3 < 4$<br>**M1d*** – solve their inequality or equation<br>**A1** – obtain $k < 1$ |
| 2 | **(a)** | | $(C =)\ 4000 + 4m$<br>$(C =)\ 6(m - 100)$ | **B1**<br>**B1**<br>**[2]** | **3.3**<br>**3.3** | Correct equation / expression for $A$<br>Correct equation / expression for $B$ | Or $40 + 0.04m$<br>Or $0.06(m - 100)$<br>**B1B0** if units inconsistent in two equations<br>**SC B1** for both $44 + 0.04m$ and $0.06m$ (or $4400 + 4m$ and $6m$) – from using $m = 0$ at 100 minutes |
| | **(b)** | | $4000 + 4m = 6(m - 100)$<br>$2m = 4600$<br><br><br>$m = 2300$ | **M1**<br><br><br><br>**A1**<br>**[2]** | **1.1**<br><br><br><br>**3.4** | Attempt to solve simultaneously, from two linear equations in $m$<br><br><br><br>Obtain 2300 (minutes) | At least one equation must have constant term<br>Could be implied by final answer of 38hrs 20 mins<br>isw once 2300 seen |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| 3 | (a) | | $x = ky^2\sqrt{z}$<br>$30 = k \times 4 \times 3$<br>$k = 2.5$ | M1 | 3.1a | Attempt to find value for $k$ | From $x = ky^2\sqrt{z}$ or $x = kz^2\sqrt{y}$ only<br>Using sum, not product, is M0 but watch for $+\sqrt{z}$ being used for positive square root |
| | | | $x = 2.5y^2\sqrt{z}$ | A1<br>[2] | 1.1 | Correct equation | Ignore modulus sign if used around $\sqrt{z}$<br>Allow BOD if initial equation stated explicitly, $k$ found correctly but then final equation not seen or seen as now incorrect |
| | (b) | | $x = 2.5 \times 9 \times 5$<br>$x = 112.5$ | M1<br>A1<br>[2] | 1.1<br>1.1 | Attempt to find $x$, from equation in terms of $y$, $z$ and numerical $k$<br>Obtain 112.5 | Could be from using direct proportion and not their equation from (a)<br>Or any exact equiv |
| 4 | (a) | | **DR**<br>f(0.5) = 0.25 – 0.75 – 5.5 + 6 = 0 | B1<br>[1] | 2.1 | Attempt f(0.5) and show equal to 0<br>Must be using factor theorem so B0 for alternative methods | B0 for just f(0.5) = 0<br>Condone $2(0.5)^3 - 3(0.5)^2 - 11(0.5) + 6 = 0$ |
| | (b) | | **DR**<br>f($x$) = $(2x - 1)(x^2 - x - 6)$ | M1<br>A1 | 1.1<br>1.1 | Attempt complete division by $(2x - 1)$<br>Obtain correct quadratic factor | **DR** so need to see quadratic factor<br>Allow equivalent complete methods eg coefficient matching / inspection / grid method<br>Condone slip(s) in otherwise correct method<br>Seen in division / correct coeffs eg $A = 1$ etc / at top of grid |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | $f(x) = (2x – 1)(x – 3)(x + 2)$ | **A1**<br><br>**[3]** | **1.1** | Obtain correct fully factorised $f(x)$ | Must be seen as a product of all 3 factors<br>**SC B1** for correct factorisation with no DR |
| | **(c)** | | **DR**<br>$x = 2^y$<br><br>$2^y = 0.5, y = –1$<br>$2^y = 3, y = 1.58$<br>$2^y = –2$, no solutions as $2^y > 0$ for all $y$<br>Hence $y = –1, y = \log_2 3$ | **B1**<br><br>**M1**<br><br>**A1**<br><br>**[3]** | **3.1a**<br><br>**1.1**<br><br>**2.4** | State or imply that $x = 2^y$<br><br>Attempt to find at least one value of $y$<br>Obtain both correct values, and no others<br><br>Must give **reason** for $2^y = –2$ having no solution | Could be implied by equating $2^y$ to at least one of their roots<br>Exact or decimal<br><br>1.58 or better, or $\frac{\log_n 3}{\log_n 2}$ for $\log_2 3$<br><br>eg cannot log a negative number<br>$2^y$ always greater than 0 |
| **5** | **(a)** | **(i)** | (2, 9) | **B1**<br>**[1]** | **1.1** | Correct coordinate | And no others |
| | | **(ii)** | (1, 12) | **B1**<br>**B1**<br><br>**[2]** | **3.1a**<br>**1.1** | Correct $x$-coordinate<br>Correct $y$-coordinate | If more than one solution given then award B1 if either co-ordinate is consistent in all solutions |
| | | **(iii)** | (6, 2) | **B1**<br>**[1]** | **1.2** | Correct coordinate | And no others |
| | **(b)** | **(i)** | $x = –2, x = 4$ | **B1**<br><br>**[1]** | **3.1a** | Both $x$-coordinates correct, and no others | Ignore any attempt at $y$ value*s* |
| | | **(ii)** | $x < –2$ | **B1**<br><br>**[1]** | **1.2** | Correct inequality, and no others | Allow ≤<br>Could be written in set notation |
| | | **(iii)** | $x = 4$ | **B1** | **1.2** | Correct $x$-coordinate, and no others | Ignore any attempt at $y$ values |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | | [1] | | | |
| 6 | (a) | | $(1-\frac{3}{8}x)^{\frac{1}{3}} = 1+(\frac{1}{3})(-\frac{3}{8}x)+\frac{(\frac{1}{3})(-\frac{2}{3})(-\frac{3}{8}x)^2}{2!}$ | B1 | 1.1 | Obtain correct first two terms | Allow unsimplified second term, including product of two fractions |
| | | | | M1 | 1.1 | Attempt third term in expansion of $(1-\frac{3}{8}x)^{\frac{1}{3}}$ | Allow BOD if no brackets, even if never recovered<br>Allow BOD if no negative sign |
| | | | $=1-\frac{1}{8}x-\frac{1}{64}x^2$ | A1 | 1.1 | Correct third term | Allow unsimplified fraction as coefficient, but must be single term |
| | | | $(8-3x)^{\frac{1}{3}} = 8^{\frac{1}{3}}(1-\frac{3}{8}x)^{\frac{1}{3}} = 2(1-\frac{3}{8}x)^{\frac{1}{3}}$<br>$(8-3x)^{\frac{1}{3}} = 2-\frac{1}{4}x-\frac{1}{32}x^2$ | B1FT | 1.1 | Correct expansion of $(8-3x)^{\frac{1}{3}}$ | FT as 2 x their expansion (at least two terms)<br>Bracket expanded and fractions simplified |
| | | | | [4] | | | |
| | (b) | | $\lvert x\rvert < \frac{8}{3}$ | B1 | 1.2 | Allow any equivalent eg $-\frac{8}{3} < x < \frac{8}{3}$ | Must be strict inequality<br>Must be condition for $x$, so B0 for $\lvert 3x\rvert < 8$ |
| | | | | [1] | | | |
| | (c) | | $(1+2x)^{-2} = 1+(-2)(2x)+\frac{(-2)(-3)}{2!}(2x)^2$ | M1 | 3.1a | Attempt first three terms of expansion | Must be expanding $(1+2x)^{-2}$<br>Allow BOD if no brackets on $2x$, even if never recovered |
| | | | $=1-4x+12x^2$ | A1 | 1.1 | Obtain correct first three terms | Allow unsimplified fraction for coeff of third term |
| | | | $(2\times 12)+(-\frac{1}{4}\times -4)+(-\frac{1}{32}\times 1)$ | M1 | 1.1 | Attempt all 3 relevant products | Finding 3 appropriate terms from the product of two 3-term quadratics<br>If part of full expansion then M1 when reqd 3 products and no others are combined |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | $\frac{799}{32}$ or $24\frac{31}{32}$ | **A1**<br><br>**[4]** | **1.1** | Any exact equivalent, including 24.96875 | Condone $x^2$ still present |
| **7** | **(a)** | | $2x\ln x+\frac{x^2-2}{x}$<br><br>$2x\ln x+\frac{x^2-2}{x}=0$<br>$2x^2\ln x+x^2-2=0$ **A.G.** | **M1**<br><br>**A1**<br><br>**[2]** | **3.1a**<br><br>**1.1** | Attempt differentiation using product rule<br><br>Equate to 0 and obtain given answer | May expand first to give $2x\ln x+\frac{x^2}{x}-\frac{2}{x}$<br>(allow middle term as just $x$)<br>Must be equated to 0 **before** clearing the fractions<br>Must be equation ie … = 0 |
| | **(b)** | | $f'(x)=4x\ln x+2x^2.\frac{1}{x}+2x$<br><br>$x_{n+1}=x_n-\frac{2x_n^2\ln x_n+x_n^2-2}{4x_n\ln x_n+2x_n^2.\frac{1}{x_n}+2x_n}$<br><br>$x_{n+1}=\frac{x_n(4x_n\ln x_n+4x_n)-(2x_n^2\ln x_n+x_n^2-2)}{4x_n\ln x_n+4x_n}$<br><br>$x_{n+1}=\frac{4x_n^2\ln x_n+4x_n^2-2x_n^2\ln x_n-x_n^2+2}{4x_n\ln x_n+4x_n}$<br><br>$x_{n+1}=\frac{2x_n^2\ln x_n+3x_n^2+2}{4x_n(\ln x_n+1)}$ **A.G.** | **B1**<br><br>**M1**<br><br>**M1**<br><br>**A1**<br><br>**[4]** | **1.1**<br><br>**1.1**<br><br>**1.1**<br><br>**2.1** | Correct derivative seen<br><br>Use correct Newton-Raphson formula, with numerator correct and their derivative in the denominator<br><br>Attempt rearrangement into single fraction with brackets expanded<br><br>Obtain given answer, with no errors seen | Allow simplified middle term of $2x$<br><br>Allow fractional term without subscripts<br>**SC** Condone use of N-R on $(x^2-2)\ln x$<br><br>Allow without subscripts<br>N-R not necessarily correct, but must be recognisable attempt<br>**SC** Rearrange their N-R on $(x^2-2)\ln x$<br><br>Subscripts needed on RHS at least one step before AG<br>LHS needs $x_{n+1}$ seen |
| | **(c)** | | $x_2=1.25, x_3=1.2075$ | **B1**<br><br>**[1]** | **1.1** | Condone 1.21, or better, for $x_3$ | $x_3 = 1.207515437…$ |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | **(d)** | | (1.206, – 0.102) | **B1**<br>**B1**<br><br><br><br>**[2]** | **2.2a**<br>**2.2a** | Correct $x$-coordinate<br>Correct $y$-coordinate | Must be 3dp or better<br>Could be given as single coordinate or $x = 1.206$, $y = -0.102$<br>Allow BOD if 1.206 given but not identified as $x$-value |
| **8** | **(a)** | **(i)** | $f(x) \in \mathbb{R}$ | **B1**<br><br><br>**[1]** | **2.5** | Allow alternative notation, or worded equivalent<br>Allow $y$, or just f, but not $x$ | Accept just $\mathbb{R}$<br>Allow $(-\infty, \infty)$ |
| | | **(ii)** | $g(x) \in (-\infty, -1] \cup [1, \infty)$ | **B1**<br><br><br>**[1]** | **2.5** | Allow alternative notation, or worded equivalent<br>Allow $y$, or just g, but not $x$ | Or $(-\infty, \infty)$ with $(-1, 1)$ clearly excluded |
| | **(b)** | **(i)** | cos (0.6) = 0.8253, so<br>$\sec(0.6) = \frac{1}{0.8253} = 1.2116$<br>$2\tan(1.2116) = 2 \times 2.6634 = 5.3269$<br>hence fg(0.6) = 5.33 **A.G.** | **M1**<br><br><br>**A1**<br>**[2]** | **2.1**<br><br><br>**2.1** | Attempt correct composition of functions<br><br><br>Conclude with 5.33 | At least one interim value required<br><br><br><br>**SC B1** for stating $2\tan(1 \div \cos 0.6) = 5.33$ |
| | | **(ii)** | f($x$) is a many to one function so has no inverse | **B1** | **2.4** | Must refer to inverse of f not existing, with reason | Must be clear that referring to the function f |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | | [1] | | | |
| | (c) | | **DR**<br>$4\tan^2 x + 6\sec x = 0$<br>$4(\sec^2 x - 1) + 6\sec x = 0$ | **M1** | **3.1a** | Attempt use of identity in their equation to obtain quadratic | Allow $\tan^2 x = \pm \sec^2 x \pm 1$<br>Award M1 when reduced to single trig ratio |
| | | | $4\sec^2 x + 6\sec x - 4 = 0$ | **A1** | **1.1** | Obtain correct equation in sec$x$ – possibly still with brackets | Or correct quadratic in cos$x$ – possibly still with brackets but with no fractions ($4\cos^2 x - 6\cos x - 4 = 0$) |
| | | | $\sec x = -2$, $\sec x = \frac{1}{2}$ | **M1** | **1.1** | Solve 3 term quadratic and attempt to find at least one value for $x$ | Could solve quadratic BC<br>Must be using root that would give solution for $x$ |
| | | | $x = \frac{2}{3}\pi,\ \frac{4}{3}\pi$ | **A1** | **1.1** | Obtain at least one correct value | Allow decimals or in degrees<br>Must be from correct solution method of correct quadratic (condone second root not being seen – but must be correct if seen) |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | $\sec x = \frac{1}{2}$ has no solutions as $\lvert\sec x\rvert \geq 1$ | **A1**<br><br>**[5]** | **2.3** | Obtain both correct values, and no others, and explain that sec$x$ = 0.5 has no solutions as outside range | Now exact and in radians<br>Or equiv explanation for cos$x$ |
| **9** | **(a)** | | $e^{kt} > 0$ for all $t$, so $y > 0$ for $t \geq 0$ (or for all $t$) hence never crosses $x$-axis | **B1**<br><br>**[1]** | **2.4** | Or show that $2e^{-3t} = 0$ has no solutions | Need to see $y \neq 0$ or $y > 0$ and reason relating to exponential (or logarithmic) function<br>Must clearly be referring to $y$ or $2e^{-3t}$ |
| | **(b)** | | $e^{2t} - 4e^t + 3 = 0$<br>$(e^t - 1)(e^t - 3) = 0$<br>$e^t = 1,\ e^t = 3$ | **M1** | **3.1a** | Equate to 0 and attempt to solve disguised quadratic | 'determine' so some evidence of method needed |
| | | | $t = 0,\ t = \ln 3$ | **A1**<br>**[2]** | **1.1** | Obtain both correct values | A0 for ln1 and not 0 |
| | **(c)** | | $\frac{dx}{dt} = 2e^{2t} - 4e^t$ | **B1** | **1.1** | Correct $\frac{dx}{dt}$ | Mark derivative and condone no/wrong label |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | $\frac{dy}{dt} = -6e^{-3t}$ | **B1** | **1.1** | Correct $\frac{dy}{dt}$ | Mark derivative and condone no/wrong label |
| | | | $\frac{dy}{dx} = \frac{-6e^{-3t}}{2e^{2t} - 4e^{t}}$ | **M1** | **2.4** | Attempt correct method to combine derivatives | Combine their derivatives correctly |
| | | | $\frac{dy}{dx} = \frac{-3e^{-3t}}{e^{2t} - 2e^{t}} = \frac{3}{e^{3t}\left(2e^{t} - e^{2t}\right)} = \frac{3}{2e^{4t} - e^{5t}}$ **A.G.** | **A1** | **2.1** | Show manipulation to given answer | Need to see some evidence of how $e^{-3t}$ is dealt with<br>**AG** so method must be fully correct |
| | | | | **[4]** | | | |
| | **(d)** | | $2e^{4t} - e^{5t} = 0$ | **M1** | **3.1a** | Equate denominator to 0 | Or $\frac{dx}{dy} = 0$ or $\frac{dx}{dt} = 0$ |
| | | | $e^{4t}\left(2 - e^{t}\right) = 0$ | **A1** | **1.1** | Solve for $t$ to obtain $t = \ln 2$ | No need to see $e^{4t} = 0$ discounted |
| | | | $t = \ln 2$<br>$(-1, \frac{1}{4})$ | **A1** | **1.1** | Obtain correct coordinate | Or $x = -1$, $y = \frac{1}{4}$ |
| | | | | **[3]** | | | |
| **10** | **(a)** | **(i)** | $\cos y = \frac{CD}{a}$ hence $CD = a\cos y$ | **B1** | **2.4** | Justification for $CD$ | Need to see either $\cos y = \frac{CD}{a}$ or<br>adj = hyp × cos$\theta$ before given answer |
| | | | | **[1]** | | | |
| | | **(ii)** | area $= \frac{1}{2}AC.CD\sin x = \frac{1}{2}b\left(a\cos y\right)\sin x$<br>$= \frac{1}{2}ab\sin x\cos y$ **A.G.** | **B1** | **2.4** | Use area of triangle to show given answer | Could quote general expression for area and then show clear substitution<br>If not, then sides being used need to be clearly identified through statement or diagram<br>Could also use right-angled triangle, with base as $AD$<br>Condone not being rearranged to given expression |
| | | | | **[1]** | | | |
| | | **(iii)** | $CD = b\cos x$ | **B1** | **2.1** | Correct $CD$ in terms of $b$ and $x$ | |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | Area $BCD$ = $\frac{1}{2}BC.CD\sin y = \frac{1}{2}a(b\cos x)\sin y$ $= \frac{1}{2}ab\cos x\sin y$ | **B1** | **2.1** | Correct area of triangle $BCD$ | B0 B1 if correct area stated with no justification |
| | | | Area $ABC$ = $\frac{1}{2}AC.BC\sin(x+y) = \frac{1}{2}ab\sin(x+y)$ | **B1** | **1.1** | Correct area of triangle $ABC$ | |
| | | | $\frac{1}{2}ab\sin(x+y) = \frac{1}{2}ab\sin x\cos y + \frac{1}{2}ab\cos x\sin y$ $\sin(x+y) = \sin x\cos y + \cos x\sin y$ | **B1** | **2.1** | Equate area of $ABC$ to the sum of the areas of the two small triangles and complete proof convincingly | Allow alternative proofs eg using lengths |
| | | | | **[4]** | | | |
| | **(b)** | | $\sin 30\cos\alpha + \cos 30\sin\alpha =$ $\cos 45\cos\alpha + \sin 45\sin\alpha$ | **B1** | **1.1** | Correct use of compound angle formulae | Could be implied if exact values used immediately – allow BOD for RHS<br>May be seen as two separate expressions, not yet equated |
| | | | $\frac{1}{2}\cos\alpha + \frac{1}{2}\sqrt{3}\sin\alpha = \frac{1}{2}\sqrt{2}\cos\alpha + \frac{1}{2}\sqrt{2}\sin\alpha$ | **M1** | **1.1** | Use exact trig values | In either equation or two expressions<br>Must see all 4 values, but expansions may not be fully correct |
| | | | $\left(\sqrt{3}-\sqrt{2}\right)\sin\alpha = \left(\sqrt{2}-1\right)\cos\alpha$ $\frac{\sin\alpha}{\cos\alpha} = \tan\alpha = \frac{\sqrt{2}-1}{\sqrt{3}-\sqrt{2}}$ | **M1** | **3.1a** | Gather like terms and attempt $\tan\alpha$<br>May still have fractions in the fraction | $\tan\alpha$ does not yet need to be the subject, but must only appear once for M1 |
| | | | $= \frac{\left(\sqrt{2}-1\right)\left(\sqrt{3}+\sqrt{2}\right)}{\left(\sqrt{3}-\sqrt{2}\right)\left(\sqrt{3}+\sqrt{2}\right)} = \frac{\sqrt{6}+2-\sqrt{2}-\sqrt{3}}{3-2}$ | **M1** | **3.1a** | Attempt to rationalise their denominator | Clear intention seen to multiply throughout by the conjugate of their denominator |
| | | | $\tan\alpha = 2+\sqrt{6}-\sqrt{3}-\sqrt{2}$ **A.G.** | **A1** | **2.1** | Obtain given answer | With full detail, including (at least) 3 – 2 in denominator |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | | [5] | | | |
| 11 | (a) | | $2u\mathrm{d}u = 2x\mathrm{d}x$ | B1 | 1.1a | Any correct expression linking d$u$ and d$x$ | Could be $\mathrm{d}u = \frac{1}{2}2x\left(x^2+3\right)^{-\frac{1}{2}}\mathrm{d}x$ or equiv in terms of $u$ |
| | | | $\int \frac{4u\left(u^2-3\right)}{\sqrt{u^2}}\mathrm{d}u$ | M1* | 2.1 | Attempt to rewrite integrand in terms of $u$ | Not just d$x$ = d$u$, unless from a clear attempt at d$u$ eg using $u = x+\sqrt{3}$ |
| | | | $\int\left(4u^2-12\right)\mathrm{d}u$ | A1 | 1.1 | Obtain correct integrand | Allow unsimplified expression |
| | | | $\frac{4}{3}u^3-12u(+c)$ | M1dep* | 1.1 | Attempt integration | Simplify to form that can be integrated, then increase all powers by 1 |
| | | | $\frac{4}{3}u(u^2-9)+c = \frac{4}{3}\left(x^2-6\right)\sqrt{x^2+3}+c$ **A.G.** | A1 | 2.1 | Obtain given answer, with at least one intermediate step seen | Need evidence of common factor (in terms of $u$ or $x$) being taken out<br>Condone omission of $+c$ |
| | | | | [5] | | | |
| | (b) | | **DR**<br>$\frac{4}{3}\left(\left(-5\times2\right)-\left(-6\times\sqrt{3}\right)\right)$ | M1 | 2.1 | Attempt to use limits $x = 0$ and $x = 1$, or $u = \sqrt{3}$ and $u = 2$ in integral in terms of $u$ | Correct order and subtraction<br>Attempt to use both limits in their integral to give two terms<br>DR so just stating decimal area is M0<br>Either using answer from (a) or their integration attempt with +2 or +3 |
| | | | $= \frac{4}{3}\left(6\sqrt{3}-10\right)$ or 0.523 | A1 | 1.1 | Obtain correct area under curve | Accept exact (inc unsimplified) or decimal<br>Using +2 gives $\frac{4}{3}\left(4\sqrt{2}-3\sqrt{3}\right)$ or 0.614 |
| | | | $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{12x^2\left(x^2+3\right)^{\frac{1}{2}}-4x^3.2x.\frac{1}{2}\left(x^2+3\right)^{-\frac{1}{2}}}{x^2+3}$ | M1 | 3.1a | Attempt derivative using the quotient rule | Or equiv with product rule<br>Need difference of two terms in numerator, at least one term correct, but allow subtraction in incorrect order<br>Using either +2 or +3 equation |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | | **A1** | **1.1** | Obtain correct, unsimplified, derivative | With either +2 or +3 |
| | | | at $x = 1$, $m = \frac{11}{2}$ hence $m' = -\frac{2}{11}$ | **M1** | **2.1** | Attempt gradient of normal at $x = 1$ | Substitute $x = 1$ and use negative reciprocal<br>Using +2 gives $m' = -\frac{3}{32}\sqrt{3}$<br>Can be with $m$ found BC |
| | | | $y - 2 = -\frac{2}{11}(x - 1)$<br>when $y = 0$, $x = 12$ | **M1** | **1.1** | Attempt to find point of intersection of normal with $x$-axis | Attempt equation of normal with their gradient and either (1, 2) or $\left(1, \frac{4}{3}\sqrt{3}\right)$, and then use $y = 0$ to find $x$ intersection |
| | | | area $= 8\sqrt{3} - \frac{40}{3} + 11$<br>$= 8\sqrt{3} - \frac{7}{3}$ | **A1**<br><br>**[7]** | **3.1a** | Obtain correct area<br>Allow any exact (including unsimplified) or decimal equivalent | From combining a correct area under curve and a correct area of triangle (either 11 or $\frac{64}{9}\sqrt{3}$), even if inconsistent<br>Can still get A1 following M0 for area under curve BC and/or $m$ found BC |
| **12** | **(a)** | **(i)** | $\frac{d\theta}{dt} = -k$ | **B1**<br><br>**[1]** | **3.3** | Allow $\frac{d\theta}{dt} = k$ **or** $\frac{d\theta}{dt} = -3.5$ | Both sides of differential equation required |
| | | **(ii)** | $\theta = -3.5t + c$ | **M1** | **3.4** | Obtain equation of the form $\theta = \pm 3.5t + c$, where $c$ could already be numerical and possibly incorrect | Not dependent on correct differential equation in **(i)** |
| | | | $\theta = 160 - 3.5t$ | **A1**<br>**[2]** | **1.1** | Obtain correct equation | **Alt method**<br>For M1, integrate to get $\theta = kt + c$, then use (0, 160) and (10, 125) to attempt $c$ and hence $k$ |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | (iii) | The model would predict that the temperature would fall below room temperature, and eventually below freezing point | B1<br><br>[1] | 3.5b | Any sensible comment | Cooling rate unlikely to be linear<br>Identify that limit (ie room temperature) will be reached |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | (b) | (i) | $\frac{d\theta}{dt} = -k(\theta - 20)$ | B1<br><br>[1] | 3.3 | Allow $\frac{d\theta}{dt} = k(\theta - 20)$ | Both sides of differential equation required<br>ISW if $k = -3.5$ used once correct equation seen (but B0 if only ever seen with – 3.5) |
| | | (ii) | $\int \frac{1}{\theta - 20} d\theta = \int -k \, dt$ | M1 | 3.1a | Separate variables (or invert each side) and attempt integration | Allow M1 for integration of a differential equation not of this form eg $\frac{d\theta}{dt} = \frac{-k}{(\theta - 20)}$<br>, as long as $t$ and/or $\theta$ are involved – must be attempt at correct rearrangement of their diff eqn |

| Question | | | Answer | Marks | AO | Guidance | |
|---|---|---|---|---|---|---|---|
| | | | $\ln\lvert\theta-20\rvert=-kt+c$ | **A1** | **1.1** | Obtain correct integral | Or $\ln\lvert\theta-20\rvert=kt+c$<br>Condone brackets not modulus |
| | | | $\ln140=c$ | **M1** | **3.4** | Use $t=0$, $\theta=160$ in an equation involving both $k$ and $c$ | Equation must be from integration attempt, but could follow M0<br>As far as numerical $c$ or $k$<br>Using both pairs of values as limits in a definite integral is M2 |
| | | | $\ln105=-10k+\ln140$<br>$k=-0.1\ln0.75$ | **M1** | **1.1a** | Use $t=10$, $\theta=125$ in an equation involving both $k$ and $c$ ($c$ possibly now numerical) | As far as numerical $c$ and $k$ |
| | | | $\ln\lvert\theta-20\rvert=(0.1\ln0.75)t+\ln140$<br>$\theta-20=e^{(0.1\ln0.75)t+\ln140}=140e^{(0.1\ln0.75)t}$ | **M1** | **1.1** | Attempt to make $\theta$ the subject | As far as correctly removing logs<br>Equation must now be of the correct form<br>ie $\ln\lvert a\theta+b\rvert=ct+d$<br>Could still be in terms of $c$ and $k$ to give eg $\theta=Ae^{kt}+20$ |
| | | | $\theta=20+140e^{(0.1\ln0.75)t}$ | **A1** | **1.1** | Obtain correct equation<br>Allow – 0.0288 (or better) for $0.1\ln0.75$ and/or 4.94 (or better) for $\ln140$ | Allow $\theta=20+e^{(0.1\ln0.75)t+\ln140}$<br>Could see $\theta=140(0.75)^{0.1t}+20$ |
| | | | | **[6]** | | | |
| | **(c)** | | $25=160-3.5t \Rightarrow t=38.6$ mins<br>$\ln5=(0.1\ln0.75)t+\ln140 \Rightarrow t=115.8$ mins | **M1** | **3.4** | Use $\theta=25$ in both of their equations to find values for $t$ | As far as two numerical values for $t$ |
| | | | 77 minutes | **A1** | **3.4** | Obtain 77 minutes | Accept any answer rounding to 77, with no errors seen |
| | | | | **[2]** | | | |

**OCR (Oxford Cambridge and RSA Examinations)**
**The Triangle Building**
**Shaftesbury Road**
**Cambridge**
**CB2 8EA**

**OCR Customer Contact Centre**

**Education and Learning**
Telephone: 01223 553998
Facsimile: 01223 552627
Email: general.qualifications@ocr.org.uk

**www.ocr.org.uk**

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Cambridge Assessment