FOR EDEXCEL

GCE Examinations

Advanced Subsidiary

Core Mathematics C1

Paper K

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

C1 Paper K – Marking Guide

1. $(2^2)^{y+3} = 2^3$ M1
$2y + 6 = 3$ M1
$y = -\frac{3}{2}$ A1 **(3)**

---


2. $= \int (3x^2 + \frac{1}{2}x^{-2})\ dx$ B1
$= x^3 - \frac{1}{2}x^{-1} + c$ M1 A2 **(4)**

---

3. $\frac{EH}{AD} = \frac{EF}{AB} \ \therefore\ \frac{EH}{\sqrt{5}} = \frac{1+\sqrt{5}}{3-\sqrt{5}}$ M1
$\frac{1+\sqrt{5}}{3-\sqrt{5}} = \frac{1+\sqrt{5}}{3-\sqrt{5}} \times \frac{3+\sqrt{5}}{3+\sqrt{5}} = \frac{3+\sqrt{5}+3\sqrt{5}+5}{9-5} = 2 + \sqrt{5}$ M2 A1
$\therefore EH = \sqrt{5}\,(2 + \sqrt{5}\,) = 5 + 2\sqrt{5}$ M1 A1 **(6)**

---

4. *(a)*



B2
B2

(b) 3 solutions B1
$x^2 - 4x + \frac{1}{x} = 0 \ \Rightarrow\ x^2 - 4x = -\frac{1}{x}$
and the graphs of $y = x^2 - 4x$ and $y = -\frac{1}{x}$ intersect at 3 points B1 **(6)**

---

5. *(a)* $(x + k)^2 - k^2 + 4 = 0$ M1
$(x + k)^2 = k^2 - 4$ A1
$x + k = \pm\sqrt{k^2 - 4}$ M1
$x = -k \pm \sqrt{k^2 - 4}$ A1

(b) $k = 3 \ \ \therefore x = -3 \pm \sqrt{3^2 - 4}$ M1
$= -3 \pm \sqrt{5}$ A1 **(6)**

---

6. *(a)* AP: $a = 77, l = -70$ B1
$S_{50} = \frac{50}{2}[77 + (-70)] = 25 \times 7 = 175$ M1 A1

(b) AP: $a = 2, d = \frac{1}{2}$ B2
$S_n = \frac{n}{2}[4 + \frac{1}{2}(n - 1)]$ M1
$= \frac{1}{4}n[8 + (n - 1)] = \frac{1}{4}n(n + 7) \qquad [\ k = \frac{1}{4}\ ]$ A1 **(7)**

---

7. $x - 3y + 7 = 0 \ \Rightarrow \quad x = 3y - 7$ M1
sub. into $x^2 + 2xy - y^2 = 7$
$(3y - 7)^2 + 2y(3y - 7) - y^2 = 7$ M1
$y^2 - 4y + 3 = 0$ A1
$(y - 1)(y - 3) = 0$ M1
$y = 1, 3$ A1
$\therefore\ x = -4,\ y = 1$ or $x = 2,\ y = 3$ M1 A1 **(7)**

© *Solomon Press*

8. *(a)* $\frac{dy}{dx} = 1 - 4x^{-3}$ B1

$\frac{d^2y}{dx^2} = 12x^{-4}$ M1 A1

(b) $y = \int (1 - 4x^{-3})\ dx$

$y = x + 2x^{-2} + c$ M1 A2

$x = -1, y = 0 \ \therefore \ 0 = -1 + 2 + c$

$c = -1$ M1

$y = x + 2x^{-2} - 1$

when $x = 2$, $y = 2 + \frac{1}{2} - 1 = \frac{3}{2}$ M1 A1 **(9)**

---

9. *(a)* $y = x - 6\sqrt{x} + 9$ M1 A1

$\frac{dy}{dx} = 1 - 3x^{-\frac{1}{2}} = 1 - \frac{3}{\sqrt{x}}$ M1 A1

(b) $x = 4 \ \therefore y = 1$

grad of tangent $= 1 - \frac{3}{2} = -\frac{1}{2}$ M1

grad of normal $= \frac{-1}{-\frac{1}{2}} = 2$ M1 A1

$\therefore \ y - 1 = 2(x - 4)$ M1

$y = 2x - 7$ A1

(c) at intersect: $x - 6\sqrt{x} + 9 = 2x - 7$

$x + 6\sqrt{x} - 16 = 0$ M1

$(\sqrt{x} + 8)(\sqrt{x} - 2) = 0$ M1

$\sqrt{x} = -8, 2$ A1

$\sqrt{x} = 2 \Rightarrow x = 4$ (at P)

$\sqrt{x} = -8 \Rightarrow$ no real solutions $\therefore$ normal does not intersect again A1 **(13)**

---

10. *(a)* $y - 4 = 3(x + 6)$ M1

$y = 3x + 22$ A1

(b) at B, $x = 0 \ \therefore y = 2 \quad \Rightarrow \quad B\ (0, 2)$ B1

at C, $x - 7(3x + 22) + 14 = 0$ M1

$x = -7$ A1

$\therefore \ C\ (-7, 1)$ A1

(c) grad $AB = \frac{2-4}{0-(-6)} = -\frac{1}{3}$ M1 A1

grad $AC = \frac{1-4}{-7-(-6)} = 3$

grad $AB \times$ grad $AC = -\frac{1}{3} \times 3 = -1$ M1

$\therefore$ AB perp to $AC \ \therefore \angle BAC = 90°$ A1

(d) $AB = \sqrt{(0+6)^2 + (2-4)^2} = \sqrt{36+4} = \sqrt{40} = 2\sqrt{10}$ M1 A1

$AC = \sqrt{(-7+6)^2 + (1-4)^2} = \sqrt{1+9} = \sqrt{10}$

area $= \frac{1}{2} \times 2\sqrt{10} \times \sqrt{10} = 10$ M1 A1 **(14)**

Total **(75)**

© *Solomon Press*

Performance Record – C1 Paper K

Question no.	**1**	**2**	**3**	**4**	**5**	**6**	**7**	**8**	**9**	**10**	Total
Topic(s)	indices	integr.	surds	curve sketch	compl. square	AP	simul. eqn	diff., integr.	diff., normal	straight lines	
Marks	3	4	6	6	6	7	7	9	13	14	75
Student											

© *Solomon Press*