FOR EDEXCEL

GCE Examinations

Advanced Subsidiary

Core Mathematics C3

Paper K

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks could be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

C3 Paper K – Marking Guide

1. *(a)* $\arctan(x-2) = -\frac{\pi}{3}$ M1

$x - 2 = \tan(-\frac{\pi}{3}) = -\sqrt{3}$ M1

$x = 2 - \sqrt{3}$ A1

(b) $1 - 2\sin^2\theta - \sin\theta - 1 = 0$ M1

$2\sin^2\theta + \sin\theta = 0$

$\sin\theta(2\sin\theta + 1) = 0$ M1

$\sin\theta = 0$ or $-\frac{1}{2}$ A1

$\theta = 0$ or $-\frac{\pi}{6}, -\pi + \frac{\pi}{6}$

$\theta = -\frac{5\pi}{6}, -\frac{\pi}{6}, 0$ A2 **(8)**

---

2. *(a)* $= \frac{4x}{(x+3)(x-3)} - \frac{2}{x+3}$ M1

$= \frac{4x-2(x-3)}{(x+3)(x-3)}$ M1

$= \frac{2x+6}{(x+3)(x-3)} = \frac{2(x+3)}{(x+3)(x-3)}$ M1

$= \frac{2}{x-3}$ A1

(b) $2^3 - 8 = 0 \ \therefore (x-2)$ is a factor of $(x^3 - 8)$ B1

$$\begin{array}{r|l} x-2 & x^2 + 2x + 4 \\ & \overline{x^3 + 0x^2 + 0x - 8} \\ & x^3 - 2x^2 \\ & \overline{\quad 2x^2 + 0x} \\ & \quad 2x^2 - 4x \\ & \overline{\qquad\quad 4x - 8} \\ & \qquad\quad 4x - 8 \end{array}$$

M1 A1

$\therefore x^3 - 8 = (x-2)(x^2 + 2x + 4)$

$\therefore \frac{x^3-8}{3x^2-8x+4} = \frac{(x-2)(x^2+2x+4)}{(3x-2)(x-2)} = \frac{x^2+2x+4}{3x-2}$ M1 A1 **(9)**

---

3. *(a)* $= -\operatorname{cosec}^2 x^2 \times 2x = -2x\operatorname{cosec}^2 x^2$ M1 A1

(b) $= 2x \times e^{-x} + x^2 \times (-e^{-x}) = xe^{-x}(2 - x)$ M1 A2

(c) $= \frac{\cos x \times (3 + 2\cos x) - \sin x \times (-2\sin x)}{(3 + 2\cos x)^2}$ M1 A1

$= \frac{3\cos x + 2\cos^2 x + 2\sin^2 x}{(3 + 2\cos x)^2} = \frac{3\cos x + 2}{(3 + 2\cos x)^2}$ M1 A1 **(9)**

---

4. *(a)* $(e^x - 3)(e^x - 5) = 0$ M1 A1

$e^x = 3, 5$

$x = \ln 3, \ln 5$ M1 A1

(b) assume $\log_2 3$ is rational B1

$\therefore \log_2 3 = \frac{p}{q}$ where p and q are integers and $q \neq 0$ M1

$\Rightarrow \quad 2^{\frac{p}{q}} = 3$ M1

$\Rightarrow \quad 2^p = 3^q$ A1

2 and 3 are co-prime $\therefore$ only solution is $p = q = 0$ M1

but $q \neq 0 \ \therefore$ contradiction $\therefore \log_2 3$ is irrational A1 **(10)**

© Solomon Press

5. *(a)* $f(x) > 0$ B1

(b) $y = 3e^{x-1}, \quad x - 1 = \ln \frac{y}{3}$ M1

$x = 1 + \ln \frac{y}{3}$

$f^{-1}(x) = 1 + \ln \frac{x}{3}, \ x \in \mathbb{R}, \ x > 0$ M1 A2

(c) $f(\ln 2) = 3e^{\ln 2 - 1} = 3e^{-1}e^{\ln 2} = 6e^{-1}$ M1 A1

$gf(\ln 2) = g(6e^{-1}) = 30e^{-1} - 2$ A1

(d) $f^{-1}g(x) = f^{-1}(5x - 2) = 1 + \ln \frac{5x-2}{3}$ M1 A1

$\therefore \ 1 + \ln \frac{5x-2}{3} = 4, \quad \frac{5x-2}{3} = e^3$ M1

$x = \frac{1}{5}(3e^3 + 2)$ A1 **(12)**

---

6. *(a)* $2x^2 + 3 \ln (2 - x) = 0 \ \Rightarrow \ 3 \ln (2 - x) = -2x^2$

$\ln (2 - x) = -\frac{2}{3}x^2$ M1

$2 - x = e^{-\frac{2}{3}x^2}$ M1

$x = 2 - e^{-\frac{2}{3}x^2} \quad [\, k = -\frac{2}{3} \,]$ A1

(b) $x_1 = 1.90988, x_2 = 1.91212, x_3 = 1.91262, x_4 = 1.91273$ M1 A1

$\therefore \ \alpha = 1.913$ (3dp) A1

$f(1.9125) = 0.0070, \ f(1.9135) = -0.020$ M1

sign change, $f(x)$ continuous $\therefore$ root A1

(c) $f'(x) = 4x + \frac{3}{2-x} \times (-1) = 4x - \frac{3}{2-x}$ M1 A1


$\therefore \ 4x - \frac{3}{2-x} = 0, \quad 4x = \frac{3}{2-x}, \quad 4x(2 - x) = 3$ M1

$4x^2 - 8x + 3 = 0, \quad (2x - 3)(2x - 1) = 0$ M1


$x = \frac{1}{2}, \frac{3}{2}$ A1 **(13)**

---

7. *(a)* *(i)*



(ii)



B3

M1 A2

(b) $2\sqrt{2} \cos x - 2\sqrt{2} \sin x = R \cos x \cos \alpha - R \sin x \sin \alpha$

$R \cos \alpha = 2\sqrt{2}, \ R \sin \alpha = 2\sqrt{2}, \quad \therefore \ R = \sqrt{8+8} = 4$ M1 A1

$\tan \alpha = 1, \ \alpha = 45$ B1

$\therefore \ f(x) = A + 4 \cos (x + 45)°$

(c) 3 B1

(d) $3 + 4 \cos (x + 45) = 0, \quad \cos (x + 45) = -\frac{3}{4}$ M1

$x + 45 = 180 - 41.4, 180 + 41.4 = 138.6, 221.4$ M1

$x = 93.6, 176.4$ (1dp) A2 **(14)**

Total **(75)**

© *Solomon Press*

Performance Record – C3 Paper K

Question no.	1	2	3	4	5	6	7	Total
Topic(s)	trigonometry	rational expressions	differentiation	exponentials and logarithms, proof	functions	numerical methods, differentiation	functions, trigonometry	
Marks	8	9	9	10	12	13	14	75
Student								

© *Solomon Press*