GCE Examinations

Advanced Subsidiary / Advanced Level

Further Pure Mathematics
Module FP1

Paper D

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong & Chris Huffer

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

FP1 Paper D – Marking Guide

1. *(a)* (3 − i) must also be root — B1
$\therefore [x-(3+\mathrm{i})][x-(3-\mathrm{i})] = x^2-6x+10$ is a factor — M1 A1

(b) linear factor is $(3x+\frac{k}{10})$ — M1 A1
$\therefore$ coeff. of $x = 30-\frac{6k}{10} = 42$ — M1
giving $k = -20$ — A1 **(7)**

---

2. $\frac{x}{x-1}-\frac{2}{3-x}>0 \ \therefore \ \frac{x^2-x-2}{(x-1)(x-3)}>0$ — M1 A1
$\frac{(x-2)(x+1)}{(x-1)(x-3)}>0 \ \therefore$ critical values are $-1, 1, 2, 3$ — M1 A1
considering change of sign of factors gives
$x<-1$ or $1<x<2$ or $x>3$ — M2 A2 **(8)**

---

3. $\frac{\mathrm{d}y}{\mathrm{d}x}+4xy=3x \ \therefore$ int. fac. $= \mathrm{e}^{\int 4x\,\mathrm{d}x} = \mathrm{e}^{2x^2}$ — M1 A1
$\mathrm{e}^{2x^2}\frac{\mathrm{d}y}{\mathrm{d}x}+4xy\,\mathrm{e}^{2x^2} = 3x\,\mathrm{e}^{2x^2}$
$\frac{\mathrm{d}}{\mathrm{d}x}(y\,\mathrm{e}^{2x^2}) = 3x\,\mathrm{e}^{2x^2}$ — M1
$y\,\mathrm{e}^{2x^2} = \int 3x\,\mathrm{e}^{2x^2}\,\mathrm{d}x = \frac{3}{4}\mathrm{e}^{2x^2}+c$ — M1 A1
$x=0, y=\frac{1}{2} \ \therefore c=-\frac{1}{4}$ — M1 A1
$\therefore y = \frac{1}{4}(3-\mathrm{e}^{-2x^2})$ — A1 **(8)**

---

4. *(a)* $\frac{3r+4}{r(r+1)(r+2)} \equiv \frac{A}{r}+\frac{B}{r+1}+\frac{C}{r+2}$ — M1
giving $A=2, B=-1, C=-1 \ \therefore \ \frac{3r+4}{r(r+1)(r+2)} \equiv \frac{2}{r}-\frac{1}{r+1}-\frac{1}{r+2}$ — M1 A1

(b) $\sum_{r=1}^{n}(\frac{2}{r}-\frac{1}{r+1}-\frac{1}{r+2})$
$= (\frac{2}{1}+\frac{2}{2}+\frac{2}{3}+\ldots+\frac{2}{n})-(\frac{1}{2}+\frac{1}{3}+\ldots+\frac{1}{n}+\frac{1}{n+1})-(\frac{1}{3}+\ldots+\frac{1}{n}+\frac{1}{n+1}+\frac{1}{n+2})$ — M1 A2
$= \frac{2}{1}+\frac{2}{2}-\frac{1}{2}-\frac{1}{n+1}-\frac{1}{n+1}-\frac{1}{n+2}$ — M1
$= \frac{5}{2}-\frac{2}{n+1}-\frac{1}{n+2}$ — A1
giving $\frac{n(5n+9)}{2(n+1)(n+2)}$ — M1 A1 **(10)**


---

5. *(a)* $\frac{\mathrm{d}y}{\mathrm{d}x} = 2ax+b \quad \frac{\mathrm{d}^2y}{\mathrm{d}x^2} = 2a$ — M1 A1
$2a+4ax+2b+10ax^2+10bx+10c = 5x^2-13x+1$ — M1
giving $a=\frac{1}{2}, b=-\frac{3}{2}, c=\frac{3}{10}$ — M1 A1

(b) aux. eqn. $m^2+2m+10=0$ — M1
$m=-1\pm 3\mathrm{i}$ — M1 A1
gen. soln. $y=\mathrm{e}^{-x}(A\cos 3x+B\sin 3x)+\frac{1}{2}x^2-\frac{3}{2}x+\frac{3}{10}$ — A2 **(10)**

© Solomon Press

6. *(a)*



B2

3 pts of intersec. ∴ 3 solutions — B1

(b) (i) f(2.5) = −0.292; f(3) = 0.721
f cont. over interval, change of sign ∴ root — M1 A1

(ii) $\alpha \approx 2.5 + \frac{0.292258}{0.292258+0.720585} \times 0.5 = 2.64$ (2dp) — M1 A2

(iii) f(2.635) = −0.0919; f(2.645) = −0.0744
no change of sign ∴ no root ∴ not correct to 2dp — M1 A1

(c) $f'(x) = \frac{2}{3} + 2\cos 2x$ — M1

$x_{n+1} = x_n - \frac{\frac{2}{3}x_n + \sin 2x_n - 1}{\frac{2}{3} + 2\cos 2x_n}$ — A1

$x_1 = 0.399956$, $x_2 = 0.407755$, $x_3 = 0.407798$ — M1 A1

root = 0.408 (3sf) — A1 **(15)**

---

7. *(a)* require $\frac{d(r\sin\theta)}{d\theta} = 0$ — M1

$r\sin\theta = a\sin\theta(1-\cos\theta) = a(\sin\theta - \frac{1}{2}\sin 2\theta)$ — M1

$\therefore \frac{d(r\sin\theta)}{d\theta} = a(\cos\theta - \cos 2\theta)$ — A1

giving $2\cos^2\theta - \cos\theta - 1 = (2\cos\theta + 1)(\cos\theta - 1) = 0$ — M1

$\therefore \cos\theta = -\frac{1}{2}, 1$ so $\theta = (0), \frac{2\pi}{3}, \frac{4\pi}{3}$ — M1 A1

$\therefore P(\frac{3}{2}a, \frac{2\pi}{3}),\ Q(\frac{3}{2}a, \frac{4\pi}{3})$ — A1

(b) area bounded by C and lines OP and OQ

$= \frac{1}{2}\int_{\frac{2\pi}{3}}^{\frac{4\pi}{3}} a^2(1-\cos\theta)^2\ d\theta = \frac{1}{2}a^2\int_{\frac{2\pi}{3}}^{\frac{4\pi}{3}} 1 - 2\cos\theta + \cos^2\theta\ d\theta$ — M1 A1

giving $\frac{1}{4}a^2\int_{\frac{2\pi}{3}}^{\frac{4\pi}{3}} 3 - 4\cos\theta + \cos 2\theta\ d\theta$ — M1 A1

$= \frac{1}{4}a^2[3\theta - 4\sin\theta + \frac{1}{2}\sin 2\theta]_{\frac{2\pi}{3}}^{\frac{4\pi}{3}}$ — A1

giving $\frac{1}{8}a^2(4\pi + 9\sqrt{3})$ — M1 A1

area of triangle $OPQ = \frac{1}{2} \times \frac{3}{2}a \times \frac{3}{2}a \times \sin\frac{2\pi}{3} = \frac{9}{16}a^2\sqrt{3}$ — M1 A1

shaded area $= \frac{1}{8}a^2(4\pi + 9\sqrt{3}) - \frac{9}{16}a^2\sqrt{3} = \frac{1}{16}a^2(8\pi + 9\sqrt{3})$ — A1 **(17)**

Total **(75)**

© Solomon Press

Performance Record – FP1 Paper D

Question no.	**1**	**2**	**3**	**4**	**5**	**6**	**7**	Total
Topic(s)	complex nos.	inequality	1st order diff. eqn	series	2nd order diff. eqn	numerical methods	polar coords	
Marks	7	8	8	10	10	15	17	75
Student								

© *Solomon Press*