GCE Examinations

Advanced Subsidiary / Advanced Level

Further Pure Mathematics
Module FP1

Paper G

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong & Chris Huffer

© Solomon Press

These sheets may be copied for use solely by the purchaser's institute.

FP1 Paper G – Marking Guide

1. $\frac{x^2-12}{x} - 1 \geq 0 \;\therefore\; \frac{x^2-x-12}{x} \geq 0$ M1 A1

$\frac{(x-4)(x+3)}{x} \geq 0 \;\therefore$ critical values are $-3, 0, 4$ M1 A1

considering change of sign of factors and expression undefined at $x = 0$ gives

$-3 \leq x < 0$ or $x \geq 4$ M1 A2 **(7)**

---

2. $5^2 + 9^2 + 13^3 + 17^2 + \ldots = \sum_{r=1}^{n} (4r+1)^2 = \sum_{r=1}^{n} (16r^2 + 8r + 1)$ M1 A2

$= 16 \times \frac{1}{6} n(n+1)(2n+1) + 8 \times \frac{1}{2} n(n+1) + n$ M1 A1

$= \frac{1}{3} n[8(2n^2 + 3n + 1) + 12(n+1) + 3]$ M1

$= \frac{1}{3} n(16n^2 + 36n + 23)$ A1 **(7)**

---

3. *(a)* $f(0) = 2;\; f(1) = -2$

f cont. over interval, change of sign $\therefore$ root M1 A1

(b) $f'(x) = 3x^2 - 10x$ M1

$x_{n+1} = x_n - \frac{x_n^3 - 5x_n^2 + 2}{3x_n^2 - 10x_n}$ M1 A1

giving $\alpha = 0.68$ (2dp) M1 A1

(c) e.g. $f'(0) = 0 \;\therefore$ tangent to curve at $x = 0$ is parallel to x-axis

and N-R uses intersection of tangent with x-axis as next approx. B2 **(9)**

---

4. *(a)* $z = \frac{1+i\sqrt{3}}{1-i\sqrt{3}} \times \frac{1+i\sqrt{3}}{1+i\sqrt{3}} = \frac{-2+2i\sqrt{3}}{4}$ M1 A1

$= \frac{-1+i\sqrt{3}}{2} = -\frac{1}{2}(1 - i\sqrt{3}) \;\therefore\; \lambda = -\frac{1}{2}$ M1 A1

(b) $\text{mod}\, z = \frac{1}{2}\sqrt{(1+3)} = 1$ B1

$\arg z = \tan^{-1}\left(\frac{\sqrt{3}}{-1}\right) = \frac{2\pi}{3}$ M1 A1

(c) $\text{mod}\, z^4 = 1^4 = 1$ M1 A1

$\arg z^4 = 4 \arg z = \frac{8\pi}{3}$ (or $\frac{2\pi}{3}$) M1 A1 **(11)**

---

5. *(a)* $\frac{dy}{dx} = p\cos x - q\sin x, \;\; \frac{d^2y}{dx^2} = -p\sin x - q\cos x$ M1 A1

$-p\sin x - q\cos x + 2p\cos x - 2q\sin x + 5p\sin x + 5q\cos x = \sin x$ M1 A1

$4p - 2q = 1$

$2p + 4q = 0$ A1

giving $p = \frac{1}{5}, \; q = -\frac{1}{10}$ M1 A1

(b) aux. eqn. $m^2 + 2m + 5 = 0$ M1

$m = \frac{-2 \pm \sqrt{4-20}}{2} = -1 \pm 2i$ M1 A1

$\therefore y = e^{-x}(A\cos 2x + B\sin 2x) + \frac{1}{5}\sin x - \frac{1}{10}\cos x$ M1 A1 **(12)**

© *Solomon Press*

6. *(a)* $\int 2\cot x \, dx = \int \frac{2\cos x}{\sin x} \, dx$ M1

$= 2 \ln|\sin x| + c$ A1

$= \ln(\sin^2 x) + c \quad [\sin^2 x \geq 0]$ A1

(b) $\frac{dy}{dx} + 2y \cot x = \operatorname{cosec} x$ M1

int. fac. $= e^{\int 2\cot x \, dx} = e^{\ln(\sin^2 x)} = \sin^2 x$ M1

$\therefore \sin^2 x \frac{dy}{dx} + 2y \sin x \cos x = \sin x$ A1

$\frac{d}{dx}(y \sin^2 x) = \sin x$ M1

$y \sin^2 x = \int \sin x \, dx$

$y \sin^2 x = -\cos x + c$ A1

(c) $x = \frac{\pi}{4}, y = 0 \ \therefore c = \frac{1}{\sqrt{2}}$ so $y \sin^2 x = \frac{1}{\sqrt{2}} - \cos x$ M1 A1

$\therefore$ when $x = \frac{\pi}{3}$, $(\frac{\sqrt{3}}{2})^2 y = \frac{1}{\sqrt{2}} - \frac{1}{2}$ M1

$\therefore \frac{3}{4}y = \frac{1}{2}(\sqrt{2} - 1)$ giving $y = \frac{2}{3}(\sqrt{2} - 1)$ A1 **(12)**

---

7. *(a)* $2(1 + \cos\theta)\cos\theta = \frac{3}{2}$ M1

$4\cos^2\theta + 4\cos\theta - 3 = 0$ A1

$(2\cos\theta - 1)(2\cos\theta + 3) = 0$ M1

$\cos\theta = -\frac{3}{2}$ (no solns) or $\frac{1}{2}$ A1

$\therefore \theta = \pm\frac{\pi}{3}$ A1

giving $A(3, \frac{\pi}{3})$ and $B(3, -\frac{\pi}{3})$ A1

(b) $\angle AOB = \frac{2\pi}{3}$ B1

area of triangle $OAB = \frac{1}{2} \times 3 \times 3 \times \sin\frac{2\pi}{3}$ M1

$= \frac{1}{2} \times 9 \times \frac{\sqrt{3}}{2} = \frac{9\sqrt{3}}{4}$ A1

(c) area between OA, curve and x-axis $= \frac{1}{2}\int_0^{\frac{\pi}{3}} 4(1 + \cos\theta)^2 \, d\theta$ M1

$= \int_0^{\frac{\pi}{3}} 2 + 4\cos\theta + 2\cos^2\theta \, d\theta$ A1

$= \int_0^{\frac{\pi}{3}} 3 + 4\cos\theta + \cos 2\theta \, d\theta$ M1

$= [3\theta + 4\sin\theta + \frac{1}{2}\sin 2\theta]_0^{\frac{\pi}{3}}$ A1

$= (\pi + 4\times\frac{\sqrt{3}}{2} + \frac{1}{2}\times\frac{\sqrt{3}}{2}) - 0 = \pi + \frac{9\sqrt{3}}{4}$ M1 A1

shaded area $= 2(\pi + \frac{9\sqrt{3}}{4}) - \frac{9\sqrt{3}}{4} = 2\pi + \frac{9\sqrt{3}}{4}$ M1 A1 **(17)**

Total **(75)**

© *Solomon Press*

Performance Record – FP1 Paper G

Question no.	**1**	**2**	**3**	**4**	**5**	**6**	**7**	Total
Topic(s)	inequality	series	numerical methods	complex nos.	2nd order diff. eqn	1st order diff. eqn	polar coords	
Marks	7	7	9	11	12	12	17	75
Student								

© *Solomon Press*