

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/13

Paper 1 Pure Mathematics 1

May/June 2020

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

- 1** Find the set of values of m for which the line with equation $y = mx + 1$ and the curve with equation $y = 3x^2 + 2x + 4$ intersect at two distinct points. [4]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

- 2** The equation of a curve is such that $\frac{dy}{dx} = 3x^{\frac{1}{2}} - 3x^{-\frac{1}{2}}$. It is given that the point (4, 7) lies on the curve.


Find the equation of the curve.

[4]

[illegible]

- 3 In each of parts (a), (b) and (c), the graph shown with solid lines has equation $y = f(x)$. The graph shown with broken lines is a transformation of $y = f(x)$.

(a)




State, in terms of f , the equation of the graph shown with broken lines.

[1]

.....

(b)




State, in terms of f , the equation of the graph shown with broken lines.

[1]

.....

(c)



State, in terms of f , the equation of the graph shown with broken lines.

[2]

.....

- 4 (a) Expand $(1 + a)^5$ in ascending powers of a up to and including the term in a^3 . [1]

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence expand $[1 + (x + x^2)]^5$ in ascending powers of x up to and including the term in x^3 , simplifying your answer. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



[6]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school handwriting practice paper. The lines are evenly spaced and run across the entire width of the page. There are no margins, text, or other markings present.

- 6** A point P is moving along a curve in such a way that the x -coordinate of P is increasing at a constant rate of 2 units per minute. The equation of the curve is $y = (5x - 1)^{\frac{1}{2}}$.
- (a)** Find the rate at which the y -coordinate is increasing when $x = 1$. [4]

[illegible]

© UCLES 2020

7 (a) Show that $\frac{\tan \theta}{1 + \cos \theta} + \frac{\tan \theta}{1 - \cos \theta} \equiv \frac{2}{\sin \theta \cos \theta}$. [4]

[illegible]

© UCLES 2020

- 8** The first term of a progression is $\sin^2 \theta$, where $0 < \theta < \frac{1}{2}\pi$. The second term of the progression is $\sin^2 \theta \cos^2 \theta$.
- (a)** Given that the progression is geometric, find the sum to infinity. [3]

[illegible]

It is now given instead that the progression is arithmetic.

- (b) (i) Find the common difference of the progression in terms of $\sin \theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Find the sum of the first 16 terms when $\theta = \frac{1}{3}\pi$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 9 The functions f and g are defined by

$$f(x) = x^2 - 4x + 3 \quad \text{for } x > c, \text{ where } c \text{ is a constant,}$$

$$g(x) = \frac{1}{x+1} \quad \text{for } x > -1.$$

- (a) Express $f(x)$ in the form $(x - a)^2 + b$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is given that f is a one-one function.

- (b) State the smallest possible value of c . [1]

.....

.....

.....

.....

.....

.....

.....

.....

It is now given that $c = 5$.

- (c) Find an expression for $f^{-1}(x)$ and state the domain of f^{-1} . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Find an expression for $gf(x)$ and state the range of gf . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 10 (a)** The coordinates of two points A and B are $(-7, 3)$ and $(5, 11)$ respectively.

Show that the equation of the perpendicular bisector of AB is $3x + 2y = 11$.

[4]


[illegible]

- (b)** A circle passes through A and B and its centre lies on the line $12x - 5y = 70$.

Find an equation of the circle.

[5]

[illegible]



The diagram shows part of the curve with equation $y = x^3 - 2bx^2 + b^2x$ and the line OA , where A is the maximum point on the curve. The x -coordinate of A is a and the curve has a minimum point at $(b, 0)$, where a and b are positive constants.

- (a) Show that $b = 3a$. [4]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dashed lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, which is otherwise blank.

[illegible]

[illegible]

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.