

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/43

Paper 4 Mechanics

May/June 2020

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages. Blank pages are indicated.

- 1 Particles P of mass m kg and Q of mass 0.2 kg are free to move on a smooth horizontal plane. P is projected at a speed of 2 m s^{-1} towards Q which is stationary. After the collision P and Q move in opposite directions with speeds of 0.5 m s^{-1} and 1 m s^{-1} respectively.

Find m .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 A minibus of mass 4000 kg is travelling along a straight horizontal road. The resistance to motion is 900 N.

(a) Find the driving force when the acceleration of the minibus is 0.5 m s^{-2} . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the power required for the minibus to maintain a constant speed of 25 m s^{-1} . [2]

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



[6]

[illegible]

- 4 A car starts from rest and moves in a straight line with constant acceleration $a \text{ m s}^{-2}$ for a distance of 50 m. The car then travels with constant velocity for 500 m for a period of 25 s, before decelerating to rest. The magnitude of this deceleration is $2a \text{ m s}^{-2}$.

(a) Sketch the velocity-time graph for the motion of the car. [1]



(b) Find the value of a . [3]

.....

.....

.....

.....

.....

.....

.....

(c) Find the total time for which the car is in motion. [3]

.....

.....

.....

.....

.....

.....

.....

- (a) Find the decrease in kinetic energy of the block as it moves from P to Q . [2]

.....

.....

.....

.....

.....

- [illegible]

- (c)** At the instant the block reaches Q , the force pushing the block up the slope is removed.

Find the time taken, after this instant, for the block to return to P .

[4]

[illegible]

- 6 A particle travels in a straight line PQ . The velocity of the particle t s after leaving P is $v \text{ m s}^{-1}$, where

$$v = 4.5 + 4t - 0.5t^2.$$

- (a) Find the velocity of the particle at the instant when its acceleration is zero. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

The particle comes to instantaneous rest at Q .

(b) Find the distance PQ .

[6]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school handwriting practice paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Two particles A and B , of masses $3m$ kg and $2m$ kg respectively, are attached to the ends of a light inextensible string. The string passes over a fixed smooth pulley which is attached to the edge of a plane. The plane is inclined at an angle θ to the horizontal. A lies on the plane and B hangs vertically, 0.8 m above the floor, which is horizontal. The string between A and the pulley is parallel to a line of greatest slope of the plane (see diagram). Initially A and B are at rest.

- (a) Given that the plane is smooth, find the value of θ for which A remains at rest. [3]

.....

.....

.....

.....

.....

.....

It is given instead that the plane is rough, $\theta = 30^\circ$ and the acceleration of A up the plane is 0.1 m s^{-2} .

- (b) Show that the coefficient of friction between A and the plane is $\frac{1}{10}\sqrt{3}$. [5]

.....

.....

.....

.....

.....

.....

- [illegible]

This image shows a blank sheet of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no other markings or text on the page.

[illegible]

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.