



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/21

Paper 2 Pure Mathematics 2

May/June 2021

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.



BLANK PAGE

1 Solve the inequality $|3x - 7| < |4x + 5|$.

[4]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dashed lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text present.

- 2** By first expanding $\sin(\theta + 30^\circ)$, solve the equation $\sin(\theta + 30^\circ) \operatorname{cosec} \theta = 2$ for $0^\circ < \theta < 360^\circ$. [6]

[illegible]

- 3 (a) Show that $(\sec x + \cos x)^2$ can be expressed as $\sec^2 x + a + b \cos 2x$, where a and b are constants to be determined. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence find the exact value of $\int_0^{\frac{1}{4}\pi} (\sec x + \cos x)^2 dx$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 A curve has parametric equations

$$x = \ln(2t + 6) - \ln t, \quad y = t \ln t.$$


(a) Find the value of t at the point P on the curve for which $x = \ln 4$. [3]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

(b) Find the exact gradient of the curve at P .

[5]

[illegible]



-
- This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

- (b) Use the equation in part (a) to show by calculation that the x -coordinate of M lies between 3 and 4. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Use an iterative formula, based on the equation in part (a), to find the x -coordinate of M correct to 5 significant figures. Give the result of each iteration to 7 significant figures. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6 (a) Use the trapezium rule with three intervals to find an approximation to $\int_1^4 \frac{6}{1+\sqrt{x}} dx$. Give your answer correct to 5 significant figures. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the exact value of $\int_1^4 2e^{\frac{1}{2}x-2} dx$. [3]

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

(c)



The diagram shows the curves $y = \frac{6}{1 + \sqrt{x}}$ and $y = 2e^{\frac{1}{2}x - 2}$ which meet at a point with x -coordinate 4. The shaded region is bounded by the two curves and the line $x = 1$.

Use your answers to parts (a) and (b) to find an approximation to the area of the shaded region. Give your answer correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

(d) State, with a reason, whether your answer to part (c) is an over-estimate or under-estimate of the exact area of the shaded region. [1]

.....

.....

.....

.....

.....

7 The polynomial $p(x)$ is defined by

$$p(x) = ax^3 - 11x^2 - 19x - a,$$

where a is a constant. It is given that $(x - 3)$ is a factor of $p(x)$.

(a) Find the value of a . [2]

.....

.....

.....

.....

.....

.....

.....

(b) When a has this value, factorise $p(x)$ completely. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

[illegible]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.