

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/62

Paper 6 Probability & Statistics 2

February/March 2020

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages. Blank pages are indicated.

- 1** The booklets produced by a certain publisher contain, on average, 1 incorrect letter per 30 000 letters, and these errors occur randomly. A randomly chosen booklet from this publisher contains 12 500 letters.

Use a suitable approximating distribution to find the probability that this booklet contains at least 2 errors. [3]

[illegible]

- 2 Lengths of a certain species of lizard are known to be normally distributed with standard deviation 3.2 cm. A naturalist measures the lengths of a random sample of 100 lizards of this species and obtains an $\alpha\%$ confidence interval for the population mean. He finds that the total width of this interval is 1.25 cm.

Find α .

[5]

[illegible]

- 3** In the past, the mean time taken by Freda for a particular daily journey was 39.2 minutes. Following the introduction of a one-way system, Freda wishes to test whether the mean time for the journey has decreased. She notes the times, t minutes, for 40 randomly chosen journeys and summarises the results as follows.

$$n = 40 \quad \Sigma t = 1504 \quad \Sigma t^2 = 57\,760$$

- (a) Calculate unbiased estimates of the population mean and variance of the new journey time. [3]

.....

.....

.....

.....

.....

- (b)** Test, at the 5% significance level, whether the population mean time has decreased. [5]

This image shows a full page of a worksheet designed for handwriting practice. It features 15 evenly spaced, horizontal dashed lines across the entire width of the page. The background is plain white, providing a clear guide for letter height and placement. There are no margins, text, or other markings present.

4 The number of accidents on a certain road has a Poisson distribution with mean 0.4 per 50-day period.

(a) Find the probability that there will be fewer than 3 accidents during a year (365 days). [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The probability that there will be no accidents during a period of n days is greater than 0.95.

Find the largest possible value of n . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 5** Bottles of Lanta contain approximately 300 ml of juice. The volume of juice, in millilitres, in a bottle is $300 + X$, where X is a random variable with probability density function given by

$$f(x) = \begin{cases} \frac{3}{4000}(100 - x^2) & -10 \leq x \leq 10, \\ 0 & \text{otherwise.} \end{cases}$$

- (a)** Find the probability that a randomly chosen bottle of Lanta contains more than 305 ml of juice. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b)** Given that 25% of bottles of Lanta contain more than $(300 + p)$ ml of juice, show that

$$p^3 - 300p + 1000 = 0. \quad [4]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Given that $p = 3.47$, and that 50% of bottles of Lanta contain between $(300 - q)$ and $(300 + q)$ ml of juice, find q . Justify your answer. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6** The volumes, in millilitres, of large and small cups of tea are modelled by the distributions $N(200, 30)$ and $N(110, 20)$ respectively.
- (a)** Find the probability that the total volume of a randomly chosen large cup of tea and a randomly chosen small cup of tea is less than 300 ml. [4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dashed lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, leaving ample room for writing practice. There is no text or other markings on the page.

- 7 A national survey shows that 95% of year 12 students use social media. Arvin suspects that the percentage of year 12 students at his college who use social media is less than the national percentage. He chooses a random sample of 20 students at his college and notes the number who use social media. He then carries out a test at the 2% significance level.

(a) Find the rejection region for the test.

[4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- (b) Find the probability of a Type I error.

[1]

.....

.....

.....

.....

- (c) Jimmy believes that the true percentage at Arvin's college is 70%. Assuming that Jimmy is correct, find the probability of a Type II error. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[illegible]

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.