

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/31

Paper 3 Pure Mathematics 3

October/November 2023

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION


- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

- 1** Find the exact coordinates of the points on the curve $y = \frac{x^2}{1-3x}$ at which the gradient of the tangent is equal to 8. [5]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

- 2 On an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 2i| \leq |z + 2 - i|$ and $0 \leq \arg(z + 1) \leq \frac{1}{4}\pi$. [4]



[4]

[illegible]

- 4 The complex number u is defined by $u = \frac{3 + 2i}{a - 5i}$, where a is real.

(a) Express u in the Cartesian form $x + iy$, where x and y are in terms of a . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given that $\arg u = \frac{1}{4}\pi$, find the value of a . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 (a) Given that

$$\sin\left(x + \frac{1}{6}\pi\right) - \sin\left(x - \frac{1}{6}\pi\right) = \cos\left(x + \frac{1}{3}\pi\right) - \cos\left(x - \frac{1}{3}\pi\right),$$

find the exact value of $\tan x$.

[4]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

(b) Hence find the exact roots of the equation

$$\sin\left(x + \frac{1}{6}\pi\right) - \sin\left(x - \frac{1}{6}\pi\right) = \cos\left(x + \frac{1}{3}\pi\right) - \cos\left(x - \frac{1}{3}\pi\right)$$

for $0 \leq x \leq 2\pi$.

[2]

[illegible]

- 6 The parametric equations of a curve are

$$x = \sqrt{t} + 3, \quad y = \ln t,$$

for $t > 0$.

- (a) Obtain a simplified expression for $\frac{dy}{dx}$ in terms of t . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence find the exact coordinates of the point on the curve at which the gradient of the normal is -2 . [3]

.....

.....

.....

.....

.....

.....

.....

.....

- 7 The variables x and θ satisfy the differential equation

$$\frac{x}{\tan \theta} \frac{dx}{d\theta} = x^2 + 3.$$

It is given that $x = 1$ when $\theta = 0$.

Solve the differential equation, obtaining an expression for x^2 in terms of θ . [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 8 (a) By sketching a suitable pair of graphs, show that the equation

$$\sqrt{x} = e^x - 3$$

has only one root.

[2]

- (b) Show by calculation that this root lies between 1 and 2.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Show that, if a sequence of values given by the iterative formula

$$x_{n+1} = \ln(3 + \sqrt{x_n})$$

converges, then it converges to the root of the equation in (a).

[1]

.....

.....

.....

.....

.....

.....

.....

- (d) Use the iterative formula to calculate the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



- [4]

[illegible]

- (b) Using the substitution $x = \sqrt{u}$, or otherwise, find by integration the exact area of the shaded region bounded by the curve, the x -axis and the line $x = 3$. [5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

10 Let $f(x) = \frac{24x + 13}{(1 - 2x)(2 + x)^2}$.

(a) Express $f(x)$ in partial fractions.

[5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- (b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) State the set of values of x for which the expansion in (b) is valid. [1]


.....

.....

.....

.....

11



In the diagram, $OABCDEFG$ is a cuboid in which $OA = 3$ units, $OC = 2$ units and $OD = 2$ units. Unit vectors $\mathbf{i}$, $\mathbf{j}$ and $\mathbf{k}$ are parallel to OA , OD and OC respectively. M is the midpoint of EF .

- (a) Find the position vector of M . [1]

.....

.....

.....

The position vector of P is $\mathbf{i} + \mathbf{j} + 2\mathbf{k}$.

- (b) Calculate angle PAM . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.