

Cambridge International AS & A Level

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/03

Paper 3 Pure Mathematics 3

For examination from 2020

SPECIMEN PAPER

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

BLANK PAGE

- 1 Find the set of values of x for which $3(2^{3x+1}) < 8$. Give your answer in a simplified exact form. [3]

[illegible]

- 2 (a) Expand $(1 + 3x)^{-\frac{1}{3}}$ in ascending powers of x , up to and including the term in x^2 , simplifying the coefficients. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) State the set of values of x for which the expansion is valid. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3 (a) Sketch the graph of $y = |2x - 3|$.

[1]

- (b) Solve the inequality $3x - 1 > |2x - 3|$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 The parametric equations of a curve are

$$x = e^{2t-3}, \quad y = 4 \ln t,$$

where $t > 0$. When $t = a$ the gradient of the curve is 2.

(a) Show that a satisfies the equation $a = \frac{1}{2}(3 - \ln a)$. [4]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, providing a template for handwriting practice or general writing. There are no margins, text, or other markings on the page.

- (b)** Verify by calculation that this equation has a root between 1 and 2. [2]

[illegible]

- (c) Use the iterative formula $a_{n+1} = \frac{1}{2}(3 - \ln a_n)$ to calculate a correct to 2 decimal places, showing the result of each iteration to 4 decimal places. [3]

This image shows a full page of white paper with horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- 5 (a) Show that $\frac{d}{dx}(x - \tan^{-1}x) = \frac{x^2}{1+x^2}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Show that $\int_0^{\sqrt{3}} x \tan^{-1}x \, dx = \frac{2}{3}\pi - \frac{1}{2}\sqrt{3}$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 The complex numbers $1 + 3i$ and $4 + 2i$ are denoted by u and v respectively.

(a) Find $\frac{u}{v}$ in the form $x + iy$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State the argument of $\frac{u}{v}$. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

In an Argand diagram, with origin O , the points A , B and C represent the complex numbers u , v and $u - v$ respectively.

- (c) State fully the geometrical relationship between OC and BA . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Show that angle $AOB = \frac{1}{4}\pi$ radians. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 7 (a) By first expanding $\cos(x + 45^\circ)$, express $\cos(x + 45^\circ) - \sqrt{2}\sin x$ in the form $R\cos(x + \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$. Give the value of R correct to 4 significant figures and the value of α correct to 2 decimal places. [5]

[illegible]


(b) Hence solve the equation

$$\cos(x + 45^\circ) - \sqrt{2} \sin x = 2,$$

for $0^\circ < x < 360^\circ$.

[4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting or typing. There are no margins, text, or other markings on the page.



- [illegible]

- (b) Calculate the angle between the directions of $\overrightarrow{ON}$ and $\overrightarrow{CM}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Show that the length of the perpendicular from M to ON is $\frac{3}{5}\sqrt{5}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



- [6]

[illegible]

- (b)** Using the substitution $u = \sin x$, find the area of the shaded region bounded by the curve and the x -axis. [4]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school handwriting practice paper. The lines are evenly spaced and run across the entire width of the page. There are no margins, text, or other markings present.

- 10** In a chemical reaction, a compound X is formed from two compounds Y and Z .

The masses in grams of X , Y and Z present at time t seconds after the start of the reaction are x , $10 - x$ and $20 - x$ respectively. At any time the rate of formation of X is proportional to the product of the masses of Y and Z present at the time. When $t = 0$, $x = 0$ and $\frac{dx}{dt} = 2$.

- (a)** Show that x and t satisfy the differential equation

$$\frac{dx}{dt} = 0.01(10 - x)(20 - x). \quad [1]$$

[illegible]

- (b) Solve this differential equation and obtain an expression for x in terms of t . [9]

[illegible]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) State what happens to the value of x when t becomes large. [1]

.....

.....

.....

.....

.....

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.