Q1. Which change would alter the value of the equilibrium constant (K_p) for this reaction?

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

A Increasing the total pressure of the system. ⬭

B Increasing the concentration of sulfur trioxide. ⬭

C Increasing the concentration of sulfur dioxide. ⬭

D Increasing the temperature. ⬭

(Total 1 mark)

Q2. This question is about the reaction given below.

$$CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$$

Enthalpy data for the reacting species are given in the table below.

Substance	CO(g)	$H_2O(g)$	$CO_2(g)$	$H_2(g)$
$\Delta H_f^{\ominus}$ / kJ mol^{-1}	−110	−242	−394	0

Which one of the following statements is **not** correct?

A The value of K_p changes when the temperature changes.

B The activation energy decreases when the temperature is increased.

C The entropy change is more positive when the water is liquid rather than gaseous.

D The enthalpy change is more positive when the water is liquid rather than gaseous.

(Total 1 mark)

Q3. The equation for the combustion of butane in oxygen is

$$C_4H_{10} + 6\ \frac{1}{2}\ O_2 \rightarrow 4CO_2 + 5H_2O$$

The mole fraction of butane in a mixture of butane and oxygen with the minimum amount of oxygen required for complete combustion is

A 0.133

B 0.153

C 0.167

C 0.200

(Total 1 mark)

Q4. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

The mole fraction of SO_3 in the equilibrium mixture is

A 0.250

B 0.350

C 0.440

D 0.700

(Total 1 mark)

Q5. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

With pressures expressed in MPa units, the value of the equilibrium constant, K_p, is

A 4.90

B 6.48

C 9.07

D 16.8

(Total 1 mark)

Q6. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

Possible units for the equilibrium constant K_p include

A no units

B kPa

C Mpa^{-1}

D kPa^{-2}

(Total 1 mark)

Q7. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

At equilibrium in the same vessel of volume 1.80 dm^3 under altered conditions, the reaction mixture contains 0.0700 mol of $SO_3(g)$, 0.0500 mol of $SO_2(g)$ and 0.0900 mol of $O_2(g)$ at a total pressure of 623 kPa. The temperature in the equilibrium vessel is

A 307 °C

B 596 K

C 337 °C

D 642 K

(Total 1 mark)

Q8. The following information concerns the equilibrium gas-phase synthesis of methanol.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa.
The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
CO(g)	−110	198
H_2(g)	0	131
CH_3OH(g)	−201	240

Possible units for the equilibrium constant, K_p, for this reaction are

A no units

B kPa

C MPa^{-1}

D kPa^{-2}

(Total 1 mark)

Q9. The following information concerns the equilibrium gas-phase synthesis of methanol.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa.
The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
CO(g)	−110	198
H_2(g)	0	131
CH_3OH(g)	−201	240

The mole fraction of hydrogen in the equilibrium mixture is

A 0.242

B 0.485

C 0.653

D 0.970

(Total 1 mark)

Q10. The following information concerns the equilibrium gas-phase synthesis of methanol.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa.
The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
CO(g)	−110	198
H_2(g)	0	131
CH_3OH(g)	−201	240

With pressures expressed in MPa units, the value of the equilibrium constant, K_p, under these conditions is

A 1.37

B 1.66

C 2.82

D 4.80

(Total 1 mark)

Q11. The following information concerns the equilibrium gas-phase synthesis of methanol.

$CO(g) + 2H_2(g) \rightleftharpoons CH3OH(g)$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa.
The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
CO(g)	−110	198
$H_2(g)$	0	131
$CH_3OH(g)$	−201	240

Which one of the following statements applies to this equilibrium?

A The value of K_p increases if the temperature is raised.

B The value of K_p increases if the pressure is raised.

C The yield of methanol decreases if the temperature is lowered.

D The yield of methanol decreases if the pressure is lowered.

(Total 1 mark)

Q1. An experiment was carried out to determine the equilibrium constant, K_c, for the following reaction.

$$CH_3CH_2COOH + CH_3CH_2CH_2OH \rightleftharpoons CH_3CH_2COOCH_2CH_2CH_3 + H_2O$$

A student added measured volumes of propan-1-ol and propanoic acid to a conical flask. A measured volume of concentrated hydrochloric acid was added to the flask, which was then sealed.

After 1 week, the contents of the flask were poured into water and the solution was made up to a known volume.
This solution was titrated with standard sodium hydroxide solution.

(a) Explain how the student could determine the amount, in moles, of propan-1-ol added to the flask.

.....

.....

.....

.....

(2)

(b) The titration described above gives the total amount of acid in the equilibrium mixture.
Explain how, by carrying out a further experiment, the student could determine the amount of propanoic acid in the equilibrium mixture.

.....

.....

.....

.....

(2)

(c) In a repeat experiment, the student failed to seal the flask that contained the equilibrium mixture.

Explain why this error would lead to the student obtaining an incorrect value for the equilibrium constant K_c

.....

.....

.....

.....

(2)
(Total 6 marks)

Q2. Many chemical processes release waste products into the atmosphere. Scientists are developing new solid catalysts to convert more efficiently these emissions into useful products, such as fuels. One example is a catalyst to convert these emissions into methanol. The catalyst is thought to work by breaking a H–H bond.

An equation for this formation of methanol is given below.

$$CO_2(g) + 3H_2(g) \rightleftharpoons CH_3OH(g) + H_2O(g) \quad \Delta H = -49 \text{ kJ mol}^{-1}$$

Some mean bond enthalpies are shown in the following table.

Bond	C=O	C–H	C–O	O–H
Mean bond enthalpy / kJ mol^{-1}	743	412	360	463

(a) Use the enthalpy change for the reaction and data from the table to calculate a value for the H–H bond enthalpy.

H–H bond enthalpy = ... kJ mol^{-1}

(3)

(b) A data book value for the H–H bond enthalpy is 436 kJ mol^{-1}.

Suggest **one** reason why this value is different from your answer to part (a).

.....

.....

.....

(1)

(c) Suggest **one** environmental advantage of manufacturing methanol fuel by this reaction.

.....

.....

.....

(1)

(d) Use Le Chatelier's principle to justify why the reaction is carried out at a high pressure rather than at atmospheric pressure.

.....

.....

.....

.....

.....

.....

(3)

(e) Suggest why the catalyst used in this process may become less efficient if the carbon dioxide and hydrogen contain impurities.

.....

.....

.....

(1)

(f) In a laboratory experiment to investigate the reaction shown in the equation below, 1.0 mol of carbon dioxide and 3.0 mol of hydrogen were sealed into a container. After the mixture had reached equilibrium, at a pressure of 500 kPa, the yield of methanol was 0.86 mol.

$$CO_2(g) + 3H_2(g) \rightleftharpoons CH_3OH(g) + H_2O(g)$$

Calculate a value for K_p
Give your answer to the appropriate number of significant figures.

Give units with your answer.

K_p = ... Units = ...

(7)
(Total 16 marks)

Q3. The equation for the combustion of butane in oxygen is

$$C_4H_{10} + 6\tfrac{1}{2}O_2 \rightarrow 4CO_2 + 5H_2O$$

The mole fraction of butane in a mixture of butane and oxygen with the minimum amount of oxygen required for complete combustion is

A 0.133

B 0.153

C 0.167

C 0.200

(Total 1 mark)

Q4. This question is about the reaction given below.

$$CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$$

Enthalpy data for the reacting species are given in the table below.

Substance	CO(g)	$H_2O(g)$	$CO_2(g)$	$H_2(g)$
$\Delta H_f^{\ominus}$ / kJ mol^{-1}	−110	−242	−394	0

Which one of the following statements is **not** correct?

A The value of K_p changes when the temperature changes.

B The activation energy decreases when the temperature is increased.

C The entropy change is more positive when the water is liquid rather than gaseous.

D The enthalpy change is more positive when the water is liquid rather than gaseous.

(Total 1 mark)

Q5. Which change would alter the value of the equilibrium constant (K_p) for this reaction?

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

A Increasing the total pressure of the system. ⬭

B Increasing the concentration of sulfur trioxide. ⬭

C Increasing the concentration of sulfur dioxide. ⬭

D Increasing the temperature. ⬭

(Total 1 mark)

Q6. The manufacture of methanol can be achieved in two stages.

(a) In the first stage, methane and steam react according to the following equation.

$$CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g) \qquad \Delta H^{\ominus} = +210 \text{ kJ mol}^{-1}$$

Discuss, with reasons, the effects of increasing separately the temperature and the pressure on the yield of the products and on the rate of this reaction.

(6)

(b) In the second stage, carbon monoxide and hydrogen react according to the following equation.

$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$

A 62.8 mol sample of carbon monoxide was added to 146 mol of hydrogen. When equilibrium was reached at a given temperature, the mixture contained 26.2 mol of methanol at a total pressure of 9.50 MPa.

Write an expression for the equilibrium constant, K_p, for this reaction. Calculate a value for K_p at this temperature and give its units.

(8)
(Total 14 marks)

Q7. A sealed flask containing gases **X** and **Y** in the mole ratio 1:3 was maintained at 600 K until the following equilibrium was established.

$$X(g) + 3Y(g) \rightleftharpoons 2Z(g)$$

The partial pressure of **Z** in the equilibrium mixture was 6.0 MPa when the total pressure was 22.0 MPa.

(a) (i) Write an expression for the equilibrium constant, K_p, for this reaction.

.....

.....

(ii) Calculate the partial pressure of **X** and the partial pressure of **Y** in the equilibrium mixture.

*Partial pressure of **X*** .....

.....

*Partial pressure of **Y*** .....

(iii) Calculate the value of K_p for this reaction under these conditions and state its units.

Value of K_p .....

.....

.....

Units of K_p .....

(6)

(b) When this reaction is carried out at 300 K and a high pressure of 100 MPa, rather than at 600 K and 22.0 MPa, a higher equilibrium yield of gas **Z** is obtained.

Give two reasons why an industrialist is unlikely to choose these reaction conditions.

Reason 1 .....

Reason 2 .....

(2)
(Total 8 marks)

Q1. Sulphur dioxide and oxygen were mixed in a 2:1 mol ratio and sealed in a flask with a catalyst.
The following equilibrium was established at temperature T_1

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \qquad \Delta H^{\ominus} = -196 \text{ kJ mol}^{-1}$$

The partial pressure of sulphur dioxide in the equilibrium mixture was 24 kPa and the total pressure in the flask was 104 kPa.

(a) Deduce the partial pressure of oxygen and hence calculate the mole fraction of oxygen in the equilibrium mixture.

Partial pressure of oxygen .....

Mole fraction of oxygen .....

.....

.....

(3)

(b) Calculate the partial pressure of sulphur trioxide in the equilibrium mixture.

.....

(1)

(c) Write an expression for the equilibrium constant, K_p, for this reaction. Use this expression to calculate the value of K_p at temperature T_1 and state its units.

Expression for K_p .....

.....

Calculation .....

.....

.....

Units .....

(4)

(d) When equilibrium was established at a different temperature, T_2, the value of K_p was found to have increased. State which of T_1 and T_2 is the lower temperature and explain your answer.

Lower temperature.....

Explanation .....

.....

(3)

(e) In a further experiment, the amounts of sulphur dioxide and oxygen used, the catalyst and the temperature, T_1, were all unchanged, but a flask of smaller volume was used.

Deduce the effect of this change on the yield of sulphur trioxide and on the value of K_p.

Effect on yield of SO_3 .....

Effect on K_p .....

.....

(2)
(Total 13 marks)

Q2. The following information concerns the equilibrium gas-phase synthesis of methanol.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa. The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^\ominus$ / kJ mol^{-1}	$S^\ominus$ / J K^{-1} mol^{-1}
CO(g)	−110	198
H_2(g)	0	131
CH_3OH(g)	−201	240

The mole fraction of hydrogen in the equilibrium mixture is

A 0.242

B 0.485

C 0.653

D 0.970

(Total 1 mark)

Q3. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

At equilibrium in the same vessel of volume 1.80 dm^3 under altered conditions, the reaction mixture contains 0.0700 mol of $SO_3(g)$, 0.0500 mol of $SO_2(g)$ and 0.0900 mol of $O_2(g)$ at a total pressure of 623 kPa. The temperature in the equilibrium vessel is

A 307 °C

B 596 K

C 337 °C

D 642 K

(Total 1 mark)

Q4. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^\ominus$ / kJ mol^{-1}	$S^\ominus$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

Possible units for the equilibrium constant K_p include

A no units

B kPa

C Mpa^{-1}

D kPa^{-2}

(Total 1 mark)

Q5. The following information concerns the equilibrium gas-phase synthesis of methanol.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa. The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^\ominus$ / kJ mol^{-1}	$S^\ominus$ / J K^{-1} mol^{-1}
CO(g)	−110	198
$H_2(g)$	0	131
$CH_3OH(g)$	−201	240

Possible units for the equilibrium constant, K_p, for this reaction are

A no units

B kPa

C MPa^{-1}

D kPa^{-2}

(Total 1 mark)

Q6. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

The mole fraction of SO_3 in the equilibrium mixture is

A 0.250

B 0.350

C 0.440

D 0.700

(Total 1 mark)

Q7. This question relates to the equilibrium gas-phase synthesis of sulphur trioxide:

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

Thermodynamic data for the components of this equilibrium are:

Substance	$\Delta H_f^{\ominus}$ / kJ mol^{-1}	$S^{\ominus}$ / J K^{-1} mol^{-1}
$SO_3(g)$	−396	+257
$SO_2(g)$	–297	+248
$O_2(g)$	0	+204

This equilibrium, at a temperature of 585 K and a total pressure of 540 kPa, occurs in a vessel of volume 1.80 dm^3. At equilibrium, the vessel contains 0.0500 mol of $SO_2(g)$, 0.0800 mol of $O_2(g)$ and 0.0700 mol of $SO_3(g)$.

With pressures expressed in MPa units, the value of the equilibrium constant, K_p, is

A 4.90

B 6.48

C 9.07

D 16.8

(Total 1 mark)

Q8. (a) The gaseous reactants **W** and **X** were sealed in a flask and the mixture left until the following equilibrium had been established.

$2W(g) + X(g) \rightleftharpoons 3Y(g) + 2Z(g) \qquad \Delta H = -200 \text{ kJ mol}^{-1}$

Write an expression for the equilibrium constant, K_p, for this reaction.
State one change in the conditions which would both increase the rate of reaction and decrease the value of K_p. Explain your answers.

(7)

(b) Ethyl ethanoate can be prepared by the reactions shown below.

Reaction 1

$CH_3COOH(l) + C_2H_5OH(l) \rightleftharpoons CH_3COOC_2H_5(l) + H_2O(l) \qquad \Delta H^{\ominus} = -2.0 \text{ kJ mol}^{-1}$

Reaction 2

$CH_3COCl(l) + C_2H_5OH(l) \rightarrow CH_3COOC_2H_5(l) + HCl(g) \qquad \Delta H^{\ominus} = -21.6 \text{ kJ mol}^{-1}$

(i) Give one advantage and one disadvantage of preparing ethyl ethanoate by **Reaction 1** rather than by **Reaction 2**.

(ii) Use the information given above and the data below to calculate values for the standard entropy change, $\Delta S^{\ominus}$, and the standard free-energy change, $\Delta G^{\ominus}$, for **Reaction 2** at 298 K.

	$CH_3COCl(l)$	$C_2H_5OH(l)$	$CH_3COOC_2H_5(l)$	$HCl(g)$
$S^{\ominus}$/JK^{1}mol^{1}	201	161	259	187

(8)
(Total 15 marks)

Q9. When a mixture of 0.345 mol of PCl_3 and 0.268 mol of Cl_2 was heated in a vessel of fixed volume to a constant temperature, the following reaction reached equilibrium.

$$PCl_3(g) + Cl_2(g) \rightleftharpoons PCl_5(g) \qquad \Delta H^{\ominus} = -93 \text{ kJ mol}^{-1}$$

At equilibrium, 0.166 mol of PCl_5 had been formed and the total pressure was 225 kPa.

(a) (i) Calculate the number of moles of PCl_3 and of Cl_2 in the equilibrium mixture.

Moles of PCl_3 .....

Moles of Cl_2 .....

(ii) Calculate the total number of moles of gas in the equilibrium mixture.

.....

(3)

(b) Calculate the mole fraction and the partial pressure of PCl_3 in the equilibrium mixture.

Mole fraction of PCl_3 .....

.....

Partial pressure of PCl_3 .....

.....

(3)

(c) (i) Write an expression for the equilibrium constant, K_p, for this equilibrium.

.....

.....

(ii) The partial pressures of Cl_2 and PCl_5 in the equilibrium mixture were 51.3 kPa and 83.6 kPa, respectively, and the total pressure remained at 225 kPa. Calculate the value of K_p at this temperature and state its units.

.....

.....

.....

(4)

(d) State the effect on the mole fraction of PCl_3 in the equilibrium mixture if

(i) the volume of the vessel were to be increased at a constant temperature,

.....

(ii) the temperature were to be increased at constant volume.

.....

(2)
(Total 12 marks)

Q10. At high temperatures, SO_2Cl_2 dissociates according to the following equation.

$SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$ $\Delta H^{\ominus} = +93 \text{ kJ mol}^{-1}$

When 1.00 mol of SO_2Cl_2 dissociates, the equilibrium mixture contains 0.75 mol of Cl_2 at 673 K and a total pressure of 125 kPa.

(a) Write an expression for the equilibrium constant, K_p, for this reaction.

.....

.....

(1)

(b) Calculate the total number of moles of gas present in the equilibrium mixture.

.....

(2)

(c) (i) Write a general expression for the partial pressure of a gas in a mixture of gases in terms of the total pressure.

.....

.....

(ii) Calculate the partial pressure of SO_2Cl_2 and the partial pressure of Cl_2 in the equilibrium mixture.

Partial pressure of SO_2Cl_2 .....

.....

Partial pressure of Cl_2 .....

.....

(5)

(d) Calculate a value for the equilibrium constant, K_p, for this reaction and give its units.

.....

.....

.....

(3)

(e) State the effect, if any, of an increase in temperature on the value of K_p for this reaction.
Explain your answer.

Effect on K_p .....

Explanation .....

.....

(2)

(f) State the effect, if any, of an increase in the total pressure on the value of K_p for this reaction.

.....

(1)
(Total 14 marks)

Q1. The following information concerns the equilibrium gas-phase synthesis of methanol.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g)$$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa. The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^\ominus$ / kJ mol^{-1}	$S^\ominus$ / J K^{-1} mol^{-1}
CO(g)	−110	198
H_2(g)	0	131
CH_3OH(g)	−201	240

With pressures expressed in MPa units, the value of the equilibrium constant, K_p, under these conditions is

A 1.37

B 1.66

C 2.82

D 4.80

(Total 1 mark)

Q2. The following information concerns the equilibrium gas-phase synthesis of methanol.

$CO(g) + 2H_2(g) \rightleftharpoons CH3OH(g)$

At equilibrium, when the temperature is 68 °C, the total pressure is 1.70 MPa. The number of moles of CO, H_2 and CH_3OH present are 0.160, 0.320 and 0.180, respectively.

Thermodynamic data are given below.

Substance	$\Delta H_f^\ominus$ / kJ mol^{-1}	$S^\ominus$ / J K^{-1} mol^{-1}
CO(g)	−110	198

$H_2(g)$	0	131
$CH_3OH(g)$	−201	240

Which one of the following statements applies to this equilibrium?

A The value of K_p increases if the temperature is raised.

B The value of K_p increases if the pressure is raised.

C The yield of methanol decreases if the temperature is lowered.

D The yield of methanol decreases if the pressure is lowered.

(Total 1 mark)

Q3. Hydrogen and carbon monoxide were mixed in a 2:1 mole ratio. The mixture was allowed to reach equilibrium according to the following equation at a fixed temperature and a total pressure of 1.75×10^4 kPa.

$$2H_2(g) + CO(g) \rightleftharpoons CH_3OH(g)$$

(a) The equilibrium mixture contained 0.430 mol of carbon monoxide and 0.0850 mol of methanol.

(i) Calculate the number of moles of hydrogen present in the equilibrium mixture.

.....

(ii) Hence calculate the mole fraction of hydrogen in the equilibrium mixture.

.....

.....

.....

(iii) Calculate the partial pressure of hydrogen in the equilibrium mixture.

.....

.....

.....

(5)

(b) In a different mixture of the three gases at equilibrium, the partial pressure of carbon monoxide was 7550 kPa, the partial pressure of hydrogen was 12300 kPa and the partial pressure of methanol was 2710 kPa.

(i) Write an expression for the equilibrium constant, K_p, for this reaction.

.....

(ii) Calculate the value of the equilibrium constant, K_p, for the reaction under these conditions and state its units.

K_p .....

.....

Units .....

(3)

(c) Two isomeric esters **E** and **F** formed from methanol have the molecular formula $C_6H_{12}O_2$

Isomer **E** has only 2 singlet peaks in its proton n.m.r. spectrum.

Isomer **F** is optically active.

Draw the structures of these two isomers.

*Isomer **E***

*Isomer **F***

(2)
(Total 10 marks)

Q4.

Summarised directions for recording responses to multiple completion questions			
A (i), (ii) and (iii) only	**B** (i) and (iii) only	**C** (ii) and (iv) only	**D** (iv) alone

Which of the following statements about a catalyst is / are true?

(i) It speeds up the forward reaction and slows down the reversere action.

(iii) It increases the proportion of molecules with higher energies.

(iii) A homogeneous catalyst usually acts in the solid state.

(iv) It does not alter the value of the equilibrium constant.

(Total 1 mark)